



2808944844

REFERENCE ONLY**UNIVERSITY OF LONDON THESIS**Degree **MD**Year **2006**Name of Author **SANBHU****Sarbjinder
Singh****COPYRIGHT**

This is a thesis accepted for a Higher Degree of the University of London. It is an unpublished typescript and the copyright is held by the author. All persons consulting the thesis must read and abide by the Copyright Declaration below.

COPYRIGHT DECLARATION

I recognise that the copyright of the above-described thesis rests with the author and that no quotation from it or information derived from it may be published without the prior written consent of the author.

LOANS

Theses may not be lent to individuals, but the Senate House Library may lend a copy to approved libraries within the United Kingdom, for consultation solely on the premises of those libraries. Application should be made to: Inter-Library Loans, Senate House Library, Senate House, Malet Street, London WC1E 7HU.

REPRODUCTION

University of London theses may not be reproduced without explicit written permission from the Senate House Library. Enquiries should be addressed to the Theses Section of the Library. Regulations concerning reproduction vary according to the date of acceptance of the thesis and are listed below as guidelines.

- A. Before 1962. Permission granted only upon the prior written consent of the author. (The Senate House Library will provide addresses where possible).
- B. 1962 - 1974. In many cases the author has agreed to permit copying upon completion of a Copyright Declaration.
- C. 1975 - 1988. Most theses may be copied upon completion of a Copyright Declaration.
- D. 1989 onwards. Most theses may be copied.

This thesis comes within category D.

☐

This copy has been deposited in the Library of _____

☐

This copy has been deposited in the Senate House Library, Senate House, Malet Street, London WC1E 7HU.

TO INVESTIGATE

USING MORPHOMETRIC AND

IMMUNOHISTOCHEMICAL CRITERIA

THE RELATIONSHIP BETWEEN

PROSTATIC INTRA-EPITHELIAL NEOPLASIA

AND PROSTATE CANCER.

UMI Number: U592409

All rights reserved

INFORMATION TO ALL USERS

The quality of this reproduction is dependent upon the quality of the copy submitted.

In the unlikely event that the author did not send a complete manuscript and there are missing pages, these will be noted. Also, if material had to be removed, a note will indicate the deletion.



UMI U592409

Published by ProQuest LLC 2013. Copyright in the Dissertation held by the Author.
Microform Edition © ProQuest LLC.

All rights reserved. This work is protected against
unauthorized copying under Title 17, United States Code.



ProQuest LLC
789 East Eisenhower Parkway
P.O. Box 1346
Ann Arbor, MI 48106-1346

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

**AN ORIGINAL THESIS SUBMITTED FOR THE
DEGREE OF DOCTOR OF MEDICINE (MD) TO
THE UNIVERSITY OF LONDON**

BY

SARBJINDER SINGH SANDHU

BSc (Hons), MBBS, FRCS (Eng.)

Department of Surgery

Royal Free Hospital School of Medicine.

Abstract:

Prostatic Intraepithelial Neoplasia (PIN) is a putative precursor of prostate carcinoma (CaP). PIN is defined as an abnormal proliferation where the cellular changes exhibit the cytological changes of cancer, but retain a basal cell layer. Using radical prostatectomy specimens we have studied the relationship between PIN and carcinoma of the prostate.

Method:

This study was divided into two parts:

- 1) Step sections from 76 consecutive radical prostatectomy specimens were examined prospectively. Using planimetry software the total volume of the gland, the volume of cancer, the volume of benign tissue, and the volume of PIN was calculated. These volumes were compared to serum PSA.
- 2) The expression of p53, bcl2, bax_{alpha}, and Ki-67 was measured using semi quantitative immunohistochemistry in 92 hormone naïve specimens as well as specimens from 14 patients who had received androgen ablation. Benign prostate specimens obtained from 19 patients were used as controls. This was carried out retrospectively.

Results:

The mean weight of the prostate gland was 61.3 (range = 26 – 173) g, the mean calculated total volume of the gland was 50.9 (range 19.4 –155.5) mls, the mean volume of cancer was 2.5 (range = 0.09 – 15) mls, the mean volume of PIN was 0.53 (range = 0 - 2.7) mls, and the mean volume of benign tissue in each gland was 47.5 (range = 13.3 – 153.4) mls. The mean value for the PSA= 9.1 (range = 3-45) ng/ml.

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

The calculated volume of the gland correlated with the weight of the gland ($p<0.0005$, $r=0.9353$). The cancer volume also had an inverse correlation with the benign and the total volume of the gland ($p<0.001$). The volume of PIN correlated with the total volume of the gland.

Prostate weight, volume and cancer volume independently correlated with serum PSA. The volume of PIN did not correlate with serum PSA. 1 gram of benign tissue correlated with 0.07 ng/ml of serum PSA, and one gram of malignant correlated with 0.8 ng/ml of serum PSA.

The expression of p₅₃ increased in a stepwise manner, 0/19 benign, 5/92 PIN, and 9/92 CaP, but the increase was not significant. Bcl₂ was expressed in 75/92 PIN specimens, 27/92 Ca P and 0/19 benign specimens ($p<0.05$). Bax_{alpha} expression was ubiquitous, but staining intensity increased when comparing benign with PIN or CaP ($p<0.05$). Ki-67 expression was absent in benign tissue and expressed in 91/92 PIN specimens, and 92/92 with CaP ($p<0.05$). In the subset of patients treated with androgen ablation areas of cancer were compared with specimens from the hormone naïve group. P₅₃ was expressed in 4/14 pts ($p=0.07$), bcl-2 expression occurred in 9/14 ($p<0.05$), bax_{alpha}-staining intensity decreased ($p<0.05$), and the number of cell expressing Ki-67 decreased ($p<0.05$).

Conclusions:

PIN volume correlates directly with gland volume, the volume of cancer correlates inversely with the volume of benign tissue. 1 gram of malignant tissue correlates with eleven times as much serum PSA when compared to benign tissue. The volume of PIN does not have any relationship with serum PSA.

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

P₅₃ expression is infrequent and may not be important in the early genesis of CaP. Bcl₂ expression increases from benign to PIN and decreases in CaP. It is expressed in the luminal cells, and confers relative immortality, resistance to cytotoxics and androgen ablation. Therefore, PIN luminal cells may have the survival advantage required for progression from PIN to CaP, and are relatively resistant to intervention. These cells may also represent the precursors of androgen resistant clones. The increased intensity of bax_{alpha} in CaP, allows it to be amenable to therapeutic intervention. Ki-67 positive cells shift from basal to luminal as one progresses from benign to malignant tissue. This implies increasing cell production in areas of malignant tissue.

Androgen ablation leads to a decrease in Ki-67 staining, an increase in the proportion of specimens expressing bcl₂ and p₅₃, and a decrease in the staining intensity for bax_{alpha}. Therefore LHRH decreases the production of cells, may use p₅₃-mediated pathways for apoptosis and by up regulating the expression of bcl₂ may cause the selection of androgen resistant immortal clones.

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

CONTENTS	Page
Abstract	3
List of Tables	12
List of Figures	14
Acknowledgements	16
 Chapter 1: Introduction	 18
1.1 Embryology and Foetal Pre-Pubertal Histology	18
1.2 Gross Anatomy of the prostate	19
1.3 Lowsley's (lobar) anatomy	25
1.4 McNeal's (Zonal) Classification	26
1.5 Histology	29
1.6 Prostate Cancer	30
1.61 Epidemiology	31
1.62 Aetiology	31
1.63 Presentation	32
1.64 Diagnosis	33
1.65 Staging of prostate cancer.	38
1.66 Management: Watchful waiting	47
1.67 Management: Active Surveillance	49
1.68 Management: Radiotherapy -external beam	50
1.69 Management: Radiotherapy -brachytherapy	51
1.70 Management: Radical prostatectomy	52

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

Chapter 2: Prostatic Intraepithelial Neoplasia	58
2.1 Prostatic Intraepithelial Neoplasia	58
2.2 Incidence of high-grade PIN in benign prostate tissue (biopsy and TURP).	64
2.3 Location of PIN	66
2.4 PIN and serum PSA levels	66
2.5 PIN and Radiotherapy	67
2.6 Clinical Significance of PIN	67
2.7 Evidence linking PIN and Cancer	68
2.8 Differential diagnosis of PIN	71
 Chapter 3: P₅₃, Bcl₂, Bax-_{alpha}, and Ki-67	 73
3.1 P ₅₃	73
3.11 Function of p ₅₃ - Guardian of the genome	76
3.12 Evidence supporting wild type p ₅₃ as a tumour suppressor gene	78
3.13 Mechanism of action of p ₅₃	79
3.14 Inactivation of p ₅₃ function	80
3.15 Detection of p ₅₃ gene and product	81
3.16 Immunohistochemical analysis	81
3.17 Molecular analysis (DNA sequencing)	84
3.18 Serological analysis	84
 3.20 P ₅₃ and human cancers	 86
3.21 P ₅₃ as an independent marker for tumour detection and early diagnosis	86
3.22 P ₅₃ as an independent marker for tumour prognosis	87
3.23 P ₅₃ as an independent marker for tumour progression	87

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

3.30	P ₅₃ and Prostate Cancer	89
3.31	P ₅₃ in patients who have undergone surgical treatment	90
3.32	P ₅₃ and watchful waiting	98
3.33	P ₅₃ and radiotherapy	99
3.34	P ₅₃ and pre-treatment biopsies	102
3.35	P ₅₃ and metastatic prostate cancer	104
3.36	The effect of castration on p ₅₃ .	106
3.37	Site of p ₅₃ mutations	108
3.38	Racial differences	109
3.4	Bcl ₂	126
3.41	Bcl ₂ and prostate cancer	133
3.42	Bcl ₂ and radical prostatectomy	134
3.43	Bcl ₂ and radical radiotherapy	136
3.44	Bcl ₂ and watchful waiting	137
3.45	Bcl ₂ and recurrence	137
3.46	Bcl ₂ and prostate biopsies	138
3.47	Androgen deprivation and Bcl ₂	139
3.48	Bcl ₂ and high grade prostatic intra epithelial neoplasia	141
3.49	Bcl ₂ and metastases	141
3.50	Bcl ₂ and bax	142
3.60	Bax	148
3.61	Incidence of Bax	149
3.62	Bax and radiotherapy for prostate cancer	149
3.63	Bax and high grade prostatic intra epithelial neoplasia	150
3.70	Ki 67	151

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

3.71	Why Study Cell Proliferation?	152
3.72	Cell Cycle	153
3.73	MIB-1	154
3.80	Ki-67 and Prostate Cancer	155
3.81	Ki-67 expression and outcome in patients with radical prostatectomy	156
3.82	Ki-67 expression and outcome in patients with radical radiotherapy	157
3.83	Ki-67 expression and outcome in patients treated with androgen withdrawal	159
3.84	The effect of androgen ablation on Ki-67	159
3.85	Ki-67 and PIN	160

Chapter 4: Aims 164

Chapter 5: Materials and Methods 165

5.1	Methods	165
5.2	Fixation	166
5.3	Cutting the specimen	166
5.4	Scanning the slices	167
5.5	Manipulating the image	169
5.6	Calculation of total prostate, PIN and tumour volume	173
5.7	Shrinkage factor	174
5.8	Slice thickness	177
5.9	Preparation of specimens	177
5.10	Dewaxing	178
5.11	Staining methods	178
5.12	Streptavidin-biotin Method	179

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

5.13	Staining Procedure	180
5.14	Endogenous Peroxidase Activity	180
5.15	Antibodies	181
5.16	Dilution of antibodies	181
5.17	Incubation times	182
5.18	Antigen retrieval (micro waving)	183
5.19	Controls	184
5.20	Principles of Interpretation of immunostaining	186
5.21	Non-specific Background Staining	187
5.22	Scoring of immunostaining	188
5.23	Background staining	189
5.24	Staining Patterns	189
5.30	Statistics	190
Chapter 6:	Results	191
6.1	Morphometric Studies.	192
6.2	Immuno-histochemical determination of p ₅₃ , bcl ₂ , bax-alpha, and PSA in benign prostate tissue, PIN, and prostate cancer.	212
Chapter 7:	Discussion	218
7.1	Relationship between total volume of gland, volume of PIN, benign volume of gland, and volume of cancer.	218
7.2	Relationship between volume of PIN, volume of benign tissue, volume of cancer and serum PSA.	220
7.3	P ₅₃ in benign, PIN and malignant prostate tissue	227
7.4	Bcl ₂ in benign, PIN and malignant prostate tissue	230

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

7.5	Bax in benign, PIN and malignant prostate tissue	232
7.6	Ki-67 in benign, PIN and malignant prostate tissue	234
Chapter 8: Conclusion		236
Chapter 9: Publications and Presentations		246
9.1	Presentations resulting from this work.	246
9.2	Publications resulting from this work.	247
Appendix		248
References		266

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

List of Tables:

1	Age Specific Reference Ranges for PSA.	37
2	Staging of Localised Prostate Cancer.	46
3	Localised Prostate Cancer deferred therapy	48
4	Prostatic intraepithelial neoplasia and cancer	63
5	Incidence of PIN	65
6	The incidence of p ₅₃ in patients with prostate cancer	110
7	The relationship between p ₅₃ and watchful waiting	113
8	P ₅₃ and outcome in patients who have had radical prostatectomy	114
9	P ₅₃ and outcome in patients who have had radiotherapy	118
10	P ₅₃ in patients treated with hormonal manipulation	119
11	P ₅₃ and outcome in patients with extra prostatic disease	120
12	P ₅₃ in pre op biopsies and outcome	121
13	Aberrant p ₅₃ and prostatic intraepithelial neoplasia	122
14	Aberrant p ₅₃ and metastatic prostate cancer	123
15	Miscellaneous papers relating p ₅₃ and prostate cancer	125
16	Bcl ₂ expression and outcome in patients with radical prostatectomy	143
17	Bcl ₂ expression and outcome in patients treated with radical radiotherapy	145
18	Bcl ₂ expression in pre operative biopsies and outcome	146
19	The relationship between bcl-2 expression and metastasis	147
20	Ki-67 (MIB1) expression and outcome in patients with radical prostatectomy	161
21	Ki-67 (MIB1) expression and outcome in patients with radical radiotherapy	162
22	Ki-67 expression in patients treated with androgen withdrawal	163

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

23	A summary of the antibodies used the incubation times, and the controls.	186
24	Percentage of p ₅₃ positive cells per slide	214
25	Intensity of p ₅₃ staining	214
26	Percentage of bcl ₂	214
27	Intensity of bcl ₂ staining	214
28	Percentage of Ki67 positive cells per slide	215
29	Intensity of bax staining	215
30	Intensity of PSA staining	215
31	Summary of results obtained by immunohistochemistry	216
32	Comparison of the benign, PIN, and cancer	217

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

List of Figures:

1	A radical prostatectomy specimen	17
2	McNeals Zonal Classification.	28
3	General Norman Schwarzkopf	30
4	The digital rectal examination	33
5	MRI of the prostate	40
6	Lymph nodes obtained by surgical dissection	42
7	A positive bone scan	43
8	The continuum of change from benign to malignant	60
9	Tufting PIN (low power)	61
10	Micropapillary PIN (low power)	61
11	Cribriform PIN (low power)	61
12	Flat PIN (low power)	61
13	Cover of Science Dec 1993.	75
14	After fixation the prostate is sliced and scanned	168
15	Areas of PIN and cancer.	170
16	Composite image of prostate	172
17	H&E stained slices for comparison with slices from fig 14	176
18	The age distribution of the patients	192
19	Stage of specimens	193
20	Distribution of weight of gland	194
21	Distribution of volume of glands	195
22	Correlation between calculated volume and weight of gland	197
23	Distribution of Gleason score	198
24	The distribution of PSA in the 76 patients studied for tumour volume	199
25	The distribution of cancer volume	200
26	The distribution of PIN volume in the 76 prostates studied	201
27	Serum PSA and benign volume of prostate.	203

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

28	Serum PSA and volume of prostate cancer	204
29	Serum PSA and PIN volume.	205
30	PIN volume and cancer volume	206
31	PIN volume versus total calculated volume	207
32	Cancer volume versus reciprocal of total volume	208
33	Cancer volume versus Gleason score	210
34	PIN volume versus Gleason score	211
35	PIN stained with PSA x 40	226
36	Cancer stained with PSA x 40	226
37	PIN stained with P ₅₃ x 40	229
38	Cancer stained with P ₅₃ x 40	229
39	Bcl ₂ positive staining PIN x 40	231
40	Bcl ₂ positive staining cancer x 20	231
41	Bax positive staining PIN x 40	233
42	Bax positive staining cancer x 40	233
43	Ki-67 positive staining PIN x 40	235
44	Ki-67 positive staining cancer x 40	235

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

Acknowledgements:

Histopathology Department

Tara Walker, Consultant Histopathologist, Queen Alexandra Hospital,
Portsmouth.

Rosalind Sim, BSc FIBMS, Dept of Pathology, Royal Free Hospital School of
Medicine.

Linda More, Dept of Pathology, Royal Free Hospital School of Medicine.

Linda Garwood, Dept of Pathology, Royal Free Hospital School of Medicine.

Department of Urology

Ross Knight, Research Registrar, Royal Free Hospital School of Medicine.

Clare Wilson, Research Nurse, Royal Free Hospital School of Medicine.

Library

David Evans, Librarian, Royal Free Hospital

Elaine Smith, Librarian, Royal Free Hospital

Statistics

Richard Morris, Department of Statistics, Royal Free Hospital

David St George, Department of Statistics, Royal Free Hospital

Roger A'Hern, Department of Statistics, Royal Marsden Hospital.

Morphometry

Chris Thrasivoulou, Senior Scientist in the Department of Anatomy, Royal
Free Hospital School of Medicine.

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

Financial Assistance:

I would like to thank the South Essex Medical Education and Research Trust for their financial assistance with this thesis.

Supervisors:

I am indebted to my supervisors Michael Jarmulowicz, Consultant Histopathologist and Honorary Senior Lecturer at the Royal Free Hospital School of Medicine and Amir V Kaisary, Consultant Urological Surgeon and Honorary Senior Lecturer at the Royal Free Hospital School of Medicine, for their inspiration, support, and encouragement, without which this piece of work would not have started and would never have finished.

In addition, I would like to thank Mr Kaisary, for guiding me through my formative years and supporting me in my urological career, he has set aspirational bench marks which one day I may reach.

Chapter 1: Introduction

1.1 Embryology and Foetal Pre-Pubertal Histology

The male foetus under the control of the product of the SRY gene from the Y chromosome produces the SRY protein. The SRY protein initiates a developmental cascade which results in the formation of the testis, male genital ducts, and associated glands, the male external genitalia, and the entire constellation of male secondary sex characteristics.

In the third month the indifferent pelvic urethra is transformed. The prostate begins development from the mesenchyme surrounding the urogenital sinus¹². Its development depends upon dihydrotestosterone, which is produced from foetal testosterone by the enzyme 5-alpha reductase in the urogenital sinus⁸⁵. Epithelial buds invaginate from the posterior urogenital sinus on either side of the verumontanum. Simultaneously the mesonephric (Wolfian) ducts develop into the seminal vesicles, epididymis, vas deferens, and ejaculatory ducts, which are stimulated by foetal testosterone. By 16 weeks of gestation the basic structure of the prostate is complete.

The glandular buds, in the foetal prostate, develop as solid outgrowth of cells. Small lumina develop lined by cuboidal or columnar epithelium¹³³. These glands are simple tubular structures without significant branching, and are

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

lined by multiple layers of immature cells with round nuclei and scant cytoplasm.

1.2 Gross Anatomy of the prostate.

The non-pathological prostate in the post pubescent male weighs 12-20g and is the shape of an inverted cone. The size of the prostate enlarges with age, in those that are over 40, the average prostate is the size of a walnut and measures approximately 4.0-4.5cm (transverse diameter), 2.5-3.0 cm (a-p diameter), and 3.0 – 4.0cm (cephal-caudal diameter).

Prostatic Intraepithelial Neoplasia and Prostate Cancer.




Fig.1: A prostatectomy specimen obtained from a retropubic radical prostatectomy.

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

A pseudo capsule that is indistinct from the surrounding fascial tissue surrounds the prostate. The surgical capsule is not a histological entity, but consists of an extension of prostatic parenchyma with dense fibrous tissue and smooth muscle that connect with the muscular layers of the prostatic urethra and cannot be separated from the prostate without tearing the glandular tissue. This tissue is thicker anterior and thins as it surrounds the prostate laterally and posterior. The pseudo-capsule has a number of areas of weakness:

- i. The insertion of the neurovascular bundles
- ii. The insertion of the ejaculatory ducts
- iii. The insertion of the internal sphincter
- iv. At the apex and in the region of the levator ani
- v. And at the base where the bladder impinges upon the prostate.

At the apex, skeletal muscle is often mixed with prostatic glands and so the boundary is often obscure. This vagueness as to what constitutes the edge of the prostate has important implications for assessing extra prostatic extension by carcinoma or whether the capsule has been cut into.

The prostate is superior to the levator ani and is separated anteriorly from the pubic symphysis and the pubic bones by the veins of Santorini, fat, lymphatic, nerves and fascial tissues. Laterally the prostate is bound by the obturator internus. Posteriorly areolar tissue and Denonvillier's fascia separate the

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

prostate from the rectum. Cranially the prostate is attached to the bladder; the paired seminal vesicles are situated on the postero-superior aspect of the prostate separating the bladder from the rectum.

The caudal tip of the prostate, the apex, is where the urethra exits the prostate. The anterior aspect of the prostate is attached to the pubic symphysis and the pubis bones by the puboprostatic ligaments. The posterior surface of the prostate contains a central furrow the median sulcus. The prostatic urethra runs from the bladder to the prostate via the horse shoe shaped internal sphincter, which is at the base of the gland. It exits the horseshoe shaped external sphincter at the apex of the prostate where it becomes the membranous urethra. The prostatic urethra does not follow a straight line as it traverses the gland but bends anteriorly approximately 35 degrees at the verumontanum. There are two bilaterally paired ejaculatory ducts that pass from the junction of the medial vas and the lateral seminal vesicles, and anteriorly and inferiorly as they join the prostatic urethra at the verumontanum.

The internal sphincter is in the region of the proximal prostatic urethra and consists of urethral musculature with incorporation of both collagen and elastic fibres. The external sphincter at the level of the levator ani is composed of striated muscle. It surrounds the membranous urethra and extends into the prostate.

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

There are two significant groups of vessels in close relation to the prostate, the neurovascular bundles (described by Walsh) and Santorini's plexus. Santorini's plexus has minimal perforating vessels involving the prostate capsule. Its major feeding vessels are the deep vein of the penis, but there are some small tributaries from the prostate. The neurovascular bundle, as its name implies, is a mixture of the arteries, veins and nerves. These are bilateral, paired and run from the apex of the gland towards the tips of the seminal vesicles and the bladder. The bundles perforate the prostate capsule, and so allow a conduit for the spread of tumour at these potential sites of capsular weakness. The bundles were initially thought to be discrete structures at the base of the gland that could be dissected off. However recent work has suggested that this may be the case in the foetus, but in the adult as the prostate grows the ejaculatory nerves along with the prostatic fascia form a concave curtain around the prostate²³⁴.

During surgery for the prostate the neurovascular bundles have a major implication with respect to the management of cancer and the potential morbidity associated with the operation. The other arterial structures are:

- i) A basilar branch of feeding vessels course inferior to the seminal vesicles, enter the prostate at the base of the gland along the ejaculatory

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

ducts, and then courses to the level of the verumontanum.

- ii) An apical branch, which enters the inferior portion of the prostate and courses to and feeds the surgical capsule.
- iii) Urethral and inner gland flow is from a combination of the anterolateral and posterolateral vessels, coursing through the outer gland.

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

1.3 Lowsley's (lobar) anatomy:

The traditional description of the prostate divided it into five major lobes²³¹.

- i) The anterior lobe, which is between the prostatic urethra and the anterior aspect of the prostate. This area arises from the ventral walls of the urethra.
- ii) The median or middle lobe, which is defined as the area between the prostatic urethra anteriorly and the ejaculatory ducts posteriorly. This area arises from the rectal surface of the urethra and lies above the orifices of the ejaculatory ducts.
- iii) The posterior lobe which extends anteriorly from the ejaculatory ducts to the posterior surface of the gland. This area arises from the urethra below the ejaculatory ducts.
- iv) and v) the two lateral lobes which are relatively symmetrical and arise from the lateral walls of the urethra.

The Lowsley classification is used everyday in clinical urology, however it is limited in its use as a means of interpreting histological anatomy.

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

1.4 McNeal's (Zonal) Classification:

This separates the prostate into the following four zones²⁴⁸:

- i) The peripheral (outer – posterior and lateral)
- ii) The central,
- iii) The preprostatic region which encompasses the periurethral ducts and the transition zone
- iv) The anterior fibro muscular stroma.

The anterior fibro muscular stroma consists of 33% of the prostate; it contains very few glands and consists of smooth muscle and dense fibrous tissue. The anterior fibro muscular stroma blends with the internal sphincter of the bladder neck proximally and the striated muscle of the external sphincter at the apex. The central zone surrounds the ejaculatory ducts with its apex at the veru montanum; this region is thought to originate from the mesonephric (Wolfian) duct, similar to the seminal vesicle. The peripheral and central zones are glandular (acinar), the majority of the prostate is composed of the peripheral region (75%). The peripheral zone is distal to the central zone and corresponds to a horseshoe shaped structure extending posteriorly, posteriolaterally and laterally. The majority of tumours arise in the peripheral zone, and the majority of benign disease occurs in the transition zone of the preprostatic region. The periurethral region consists of cylindrical smooth muscle

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

sphincter surrounding the urethra and extending from the verumontanum to the bladder neck. The transition zone consists of ducts, which extend laterally in the region of the verumontanum. The transition zone in health consists of only 5% of the normal prostate but it may grow considerably in benign glandular stromal hyperplasia.

This zonal classification is often simplified to a two-zone concept i.e.

- i) inner (transition) zone, and
- ii) outer (central and peripheral) zone.

Prostatic Intraepithelial Neoplasia and Prostate Cancer.




Figure2: McNeals Zonal Classification.

The transition zone surrounds the urethra proximal to the ejaculatory ducts. The central zone surrounds the ejaculatory ducts, and the peripheral zone consists of apical, posterior and lateral regions of the prostate²⁴⁸.

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

1.5 Histology:

The prostate consists of stromal (smooth muscle cells, fibroblasts and endothelial cells) and epithelial cells. The epithelial cells line the alveoli. The epithelial cells consist of columnar secretory cells, basal cells and neuroendocrine cells. The tall columnar cells are secretory and are attached to each other by cadherins and to the basement membrane by integrins attached to laminin and fibronectin. These cells are terminally differentiated.

The basal cells lie beneath the secretory cells, and lie parallel to the basement membrane. The cells are elongated and are less differentiated than secretory cells. Basal cells are thought to be the progenitors of secretory cells. Basal cells are absent in adenocarcinoma of the prostate.

Neuroendocrine cells are present within the prostatic ducts and acini, and in the urothelium of the prostatic urethra ³. These cells may function by endocrine, paracrine, neurocrine, and lumencrine mechanisms. They play an important regulatory role, both during growth and differentiation of the prostate and also in the secretory process of the mature gland.

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

1.6 Prostate Cancer



Fig 3: General Norman Schwarzkopf on the cover of Time, April 1996, having been diagnosed with prostate cancer.

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

1.61 Epidemiology

In 2002, 8277 deaths occurred due to carcinoma of the prostate in England and Wales- the second commonest cancer related cause of death²⁸². Data from the Office of Population Censuses and Surveys (OPCS) has recorded an increase in the incidence from 34.0 per 100,000 in 1980 to 79.3 per 100,000 in 2002. The OPCS believe that the increase in the incidence is not only due to better detection but also the underlying prevalence of the prostatic carcinoma is increasing. An ageing population coupled with an increased awareness amongst the medical profession and the lay public are also contributory factors. However an obvious disparity exists between the number of patients with carcinoma of the prostate and the number that succumb to the disease.

1.62 Aetiology

A number of risk factors have been implicated, age is the most important, 95% of cases occur between 45 and 89 years, the median age of presentation being 72; 90% of deaths occur after 65 at a median age of 77 years. Post-mortem studies have revealed 40% of 90-year-old patients have carcinoma of the prostate. Race is a risk factor, blacks are affected more than whites and Oriental's have the lowest prevalence. The highest prevalence is amongst black men in Washington DC⁷¹.

An individual's risk is increased three fold if a first-degree relative has

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

carcinoma of the prostate. A group from the University of Utah is undertaking genealogy studies with the large Mormon population in Utah²⁵² and a group at the Royal Marsden is studying familial prostate cancer¹¹¹. A high fat diet has also been implicated, as have androgens. Androgens are essential for gland proliferation and in experimental models induce carcinoma of the prostate. There is no clear relationship between serum testosterone levels and prostatic carcinoma but patients with low levels have a poor response to treatment.

1.63 Presentation

Carcinoma of the prostate presents principally in four ways. Patients present with symptoms of bladder outlet obstruction (this used to be the majority), secondly an incidental finding upon histopathological examination of prostatic chips, thirdly patients may present with metastatic disease, and finally screening whether it is specifically for carcinoma of the prostate or a routine examination. Among men with newly diagnosed carcinoma of the prostate one third have distant metastasis, 10-15% have locally extensive tumour (about half have lymph node metastasis), and 50-60% have clinically localised disease (a third have lymph node metastasis). Thus two-third of newly diagnosed patients have disease which is not localised and so is amenable only to palliation³⁴⁰.

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

1.64 Diagnosis



Fig 4: The digital rectal examination still remains important.

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

Three primary diagnostic tests exist to initiate histological or cytological tissue sampling: digital rectal examination (DRE), trans-rectal ultrasound (TRUS), and prostate specific antigen (PSA). The clinical impression of the examining urologist correlates well with the positive predictive value of DRE. On biopsy samples from clinically normal prostates a positive predictive value of 2% was calculated, compared to 83% on those with marked induration³⁶³.

Transrectal ultrasound is the commonest modality used to image the prostate. It is of limited benefit with a distinctly palpable lesion. However when the DRE is unremarkable and the PSA is raised the positive predictive value of a TRUS guided biopsy is 16.4%. If the PSA is normal and the DRE is abnormal it is 13%, and if both tests are unremarkable it falls to 9.3%⁸⁸. If DRE is abnormal and a digitally guided biopsy result is benign then in the same cohort a TRUS guided biopsy will be positive in 53% of patients¹⁶⁴. The addition of sextant biopsies further improves the detection rate¹⁶⁵ and more recent work has shown that increasing the number of biopsies as a factor of the gland volume leads to further improvements in detection rate^{104;237}. Up to six sets of biopsies may be necessary to rule out cancer of the prostate³²⁴.

Prostate specific antigen is the best marker for prostatic carcinoma and probably the best tumour marker currently available. It is an established

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

immunohistochemical marker and is used to monitor patients. It can also be used to aid staging and diagnosis, and proponents of screening strongly advocate its use as a screening tool. PSA is a serine protease secreted into the ductal system of the prostate where it is thought to catalyse the liquefaction of the seminal coagulum after ejaculation. However, it does not explain the clotting and liquefaction of semen, which is so important for reproduction. Any process that disrupts the normal architecture of the prostate allows diffusion of PSA into the stroma; here it gains access to the blood stream via the stromal micro vasculature. Elevated levels of serum PSA are detected in prostatitis, infarcts, benign prostatic hypertrophy, transiently after prostatic manipulation and with prostatic adenocarcinoma. Cancer produces less PSA per cell than benign epithelium but the greater number of malignant cells and the stromal disruption associated with cancer account for the clinically important elevation in serum PSA levels ³⁴. The higher the grade of the cancer, the less differentiated the cells and the less PSA is produced with Gleason grade 5 tumours contributing relatively little PSA to the serum PSA⁵

A number of PSA parameters have been described to improve its sensitivity, specificity, and clinical utility. The normal range of PSA is less than 4.0ng/ml, however in 20% of patients with newly diagnosed prostatic carcinoma the PSA levels are less than 4.0 ng/ml. PSA may increase with age as a result of benign hypertrophy. Therefore, age specific reference ranges (Table 1) have been

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

introduced and appear to increase the sensitivity of PSA in men younger than 60 years and the specificity in men older than 60 ²⁹⁰.

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

Age yr.	40	45	50	55	60	65	70	75	80
PSA ng/ml	2.0	2.4	2.8	3.3	3.8	4.5	5.3	6.2	7.2

TABLE 1: Age-specific reference ranges for PSA.

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

PSA density, i.e. serum PSA divided by the volume of the prostate, was introduced to adjust for contribution of benign prostatic hypertrophy to serum PSA levels, it was found to be a better predictor of pathological stage than the Gleason grade^{26:347}. Others have not found this parameter useful^{51:290}.

A PSA velocity, i.e. the rate of increase of PSA over time, greater than 0.75 ng/ml per year is an accurate predictor of cancer identifying 72% of cancer cases with a specificity of 90%⁶⁹. In cohort studies serum PSA correlates with cancer volume, tumour grade and pathological stage^{300 34}, but it is not reliable on an individual basis.

If PSA is used as the initial test in asymptomatic men more than 50 years of age, then with the aid of TRUS guided biopsies 22% of men with a PSA of 4.0 - 9.9 ng/ml are diagnosed as having carcinoma of the prostate. This figure increases to 67% in men with PSA > 10.0ng/ml⁷².

Any of the three primary investigations can lead to a tissue biopsy, but PSA is the most sensitive single parameter⁷² and the positive predictive value of DRE and TRUS can be increased with a raised PSA.

1.65 Staging of Prostate Cancer.

Once the diagnosis of prostatic carcinoma is made, staging is carried out with

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

radiological imaging, a nuclear bone scan and TRUS. As part of the WHO consensus conference on imaging it was concluded that MRI is superior to other modalities in the assessment of local tumour stage, particularly if an endorectal coil was used ^{167:242}. The majorities of lesions arise in the peripheral zone and are seen as low signals on T₂-weighted images. The accuracy of MRI in differentiating invasive from localised disease is well established. A peripheral zone defect of 1cm or greater with an ill defined border and of low signal proved to be 100% sensitive ³⁰⁶ and 54% specific ³⁰⁹ for extra capsular spread (fig. 5).

Prostatic Intraepithelial Neoplasia and Prostate Cancer.



Fig 5a



Fig5b

Fig 5: Trans abdominal MRI of the prostate, a large left sided cancer can be seen in the peripheral zone. Fig 5a shows a T2 cancer and Fig 5b a T3 lesion. Although the images obtained with transrectal MRI were initially better, improvement in MRI technology has allowed extra corporal MRI to supersede this.

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

The value of MRI in detecting nodal disease was limited to evaluation of lymph node size and its accuracy, i.e. 80-90%, has not exceeded that of CT³¹. Any node greater than 1.5cm is likely to be malignant. However more recent work with ultra small particles of iron oxide – ferumoxtran-10, in pelvic malignancies has shown that the sensitivity may be increased with no loss of specificity³²³. It remains to be seen if this is of any benefit in the detection of pelvic lymph nodes in patients with prostate cancer.

Ultrasound imaging of the prostate via the abdominal approach is unreliable, the use of the transrectal probe has allowed excellent high resolution images of the peripheral zone where 70% of the carcinomas arise²⁴³, however it should be remembered that 24% of prostatic tumours are isoechoic⁹⁵. Transrectal ultrasound detection of local extension is operator dependent and the accuracy varies from 60 to 100%^{322:339}.

The utility of CT is diminishing. It may have a use in situations where MRI scans are inconclusive, e.g. the assessment of para-aortic nodes.

Prostatic Intraepithelial Neoplasia and Prostate Cancer.




Fig 6: Lymph nodes obtained by surgical dissection.

Prostatic Intraepithelial Neoplasia and Prostate Cancer.




Fig 7: A positive bone scan in a patient with multiple metastasis.

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

The axial skeleton is the commonest site of haematogenous metastasis. Bone metastasises are assessed with scintigraphy (fig.7) that can point to the need for plain radiographs. Routine radiographic surveys have no place in imaging for skeletal involvement; it is costly, time consuming and involves a significant dose of radiation. Furthermore there must be a 50% loss of bone density before a lesion is visible²²¹ and 23% of patients judged to be free of bone metastasises by radiography have metastatic disease by scintigraphy³⁰³.

The mainstay of assessing chest involvement is a chest radiograph²⁴².

The level of serum prostate specific antigen is determined by prostate cancer volume and by the percentage of high-grade cancer¹⁸⁸. Therefore, the use of serum PSA coupled with tumour grade and clinical stage has been advocated by Partin et al as a staging tool³⁰². In a series of pelvic lymphadenectomies if the PSA < 6ng/ml and the Gleason score < 5 all the pelvic lymph nodes were tumour free (n=142), but if the PSA < 10 ng/ml and the Gleason score < 6 then 1% of the pelvic lymph adenectomies were positive (n=388)²⁸¹.

Once the localised tumour has been staged using the TMN classification, (Table2) the patient is a candidate for one of the three management regimes available:

- i) watchful waiting

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

ii) active surveillance,

iii) radiotherapy - external beam

brachytherapy

iii) radical prostatectomy- retro pubic,
perineal, and
laparoscopic.

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

TABLE 2: Staging of Localised Prostate Cancer (UICC 2002).

T	Primary Tumour
TX	Primary tumour cannot be assessed
T0	No evidence of primary tumour
T1	Clinically inapparent tumour not palpable or visible by imaging.
T1a	Clinically inapparent tumour not palpable.
T1b	Tumour incidental histological finding in more than 5% of the tissue.
T1c	Tumour identified by needle biopsy.
T2	Tumour confined within the prostate or invading the apex.
T2a	Tumour involves one half of one lobe or less.
T2b	Tumour involves more than half of one lobe, but not both lobes.
T2c	Tumour involves both lobes.
T3	Tumour extends through the prostatic capsule.
T3a	Unilateral or bilateral extra capsular extension.
T3b	Tumour invades the seminal vesicles.
N0	No regional lymph node metastasis.
M0	No distant metastasis.
Nx	Regional lymph nodes not sampled.
N0	No positive regional lymph nodes.
N1	Regional lymph node metastasis
MX	Distant metastasis cannot be assessed.
M0	No distant metastasis.
M1	Distant metastasis.

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

1.66 Management:

Watchful Waiting.

Patients are initially followed without treatment and receive subsequent treatment for symptoms e.g. androgen ablation for metastatic disease or transurethral resection for outlet obstruction. Limited data is available from 4 studies about the outcome of watchful waiting (Table 3).

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

TABLE 3:

Localised prostate cancer, deferred therapy: Progression, Metastatic Rate, and Prostate Cancer Death.

Author	No. of patients.	Mean age	5 years/%		
			Prog.	Met.	Death
George ¹²⁶	120	75	84	11	20
Johansson ¹⁸⁴	223	72	31	9	8
Whitmore ⁴¹⁵	75	67	69	37	0
Adolfsson ⁴	122	68	55	14	1

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

These studies demonstrate that within 5 years 31-84% of patients will progress and that metastasis will occur in 9 to 37% of patients. A pooled analysis of six non randomised studies, which included three of the four studies listed in table 3^{4:184:415}, concluded at 10 years the disease specific survival for low grade disease (grade 1+2) was 87% with a metastasis free survival rate of 81% for grade 1, and 58% for grade 2. The figures for high-grade (grade 3) disease were 34 % and 26% respectively⁸².

A more recent study from Denmark of 2,570 patients with newly diagnosed clinically localised prostate cancer treated with watchful waiting; radical prostatectomy was not performed in Denmark before 1995, and external beam radiotherapy was only rarely used; demonstrated an annual mortality of 13%, with an increased in mortality for all ages at follow up. Follow up was at 10-14 yrs, 15-19 yrs, and greater than 20 years after diagnosis⁴⁹. Additionally the results of the Medical Research Council immediate versus deferred treatment study for carcinoma of the prostate have demonstrated more rapid progression and death in the deferred treatment group for all stages⁹⁴.

1.67 Management: Active surveillance.

The rationale behind this management option is that the majority of patients have a disease which is indolent and will not progress, and to treat these patients with curative intent would lead to unnecessary morbidity without any

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

survival benefit. Therefore active surveillance, to paraphrase Whitmore, attempts to separate the pussycats from the tigers. Patients with localised organ confined disease are followed up with three monthly reviews including PSAs DREs, and repeat biopsies, as part of a study protocol, no intervention is required unless there is evidence of disease progression – progression in PSA, Gleason score, or clinically.

If there is evidence of progression or it is felt that the patient is not suitable (high Gleason score) for active surveillance, the patient is referred for curative therapy. This is distinct from watchful waiting, which relies on the use of palliative intervention for symptomatic progression^{157:297}.

1.68 Management: Radiotherapy - External Beam:

The Stanford series consists of 1119 patients (from October 1956 to December 1990) treated with external beam irradiation derived from a mid to high-energy linear accelerator (range, 6-25 mV). 673 patients had disease confined to the prostate¹⁴. These authors found survival to be inversely proportional to clinical stage and pathological grade. At 15 years the survival rates for the best prognosis tumours were equivalent to that of an age matched cohort i.e. 50% at 15 years, however the poorer prognosis tumours had a survival rate of 18% at 15 years. Age had no influence on outcome.

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

More recently the use of adjuvant androgen blockade along with external beam radiotherapy has been championed. In a randomised study, androgen ablation was started at the time of the first dose of irradiation and continued for three years after. In patients with locally advanced disease, there was a survival advantage and decreased local progression in the group on androgen ablation therapy^{37,38}.

1.69 Management: Radiotherapy - Brachytherapy:

In this form of radiotherapy, radioisotopes are inserted directly into the prostate. It has been used for the management of other malignancies arising in organs accessible to this type of delivery system including the brain and the gynaecological tract. The Seattle group has data from 634 consecutive patients who have undergone brachytherapy³⁶, with 10 year follow up available on 125 patients¹⁴⁰. The initial results appear to be similar to those obtained for external beam radiotherapy or surgery. Brachytherapy is not a benign form of management. 5-10% of patients develop short term urinary retention³⁶, the Seattle group report incontinence rates of less than 1% and potency rates of 75 to 90% after treatment³⁶. Other series have not been so flattering reporting 40% of patients developing urinary incontinence, with 18% reporting having used a pad in the last week³⁷⁸. The risk of incontinence appears to increase with time and 83% of men who had had a previous TURP were incontinent³⁷⁸. The incidence of total impotence is 45%, with 68% of men having an erection

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

inadequate for penetration³⁷⁸.

1.70 Management: Radical Total Prostatectomy.

Surgery is curative only if the entire tumour is removed. Two approaches are advocated for open surgery, retro-pubic or perineal, the results are comparable. Precise control of bleeding from the dorsal vein complex allows anatomic dissection of the apex, and thus identifies the branches of the pelvic plexus which innervate the corpora and so preserves sexual function⁴⁰⁹. Incontinence (6-10%) and impotence (30%) are serious side effects.

Laparoscopic radical prostatectomy is slowly gaining favour in the United Kingdom. Richard Gaston developed the initial technique in Bordeaux; however this remains an unpublished series. The procedure was popularised by Guillonnet and Vallencien¹⁴³⁻¹⁴⁵. The Montsouris group advocated a transperitoneal approach. This technique is based on the primary incision of the peritoneum above the recto-vesicle cul-de-sac followed by dissection of the seminal vesicles. This approach does not adhere to the basic laparoscopic tenet of replicating the open operation. Therefore, an alternative extra peritoneal approach has been developed³⁹.

Whichever approach is adopted, open or laparoscopic, perineal or retro pubic, transperitoneal or extra peritoneal, oncological cure coupled with acceptable

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

functional results remains the imperative. To this end, data from the Virginia Mason Clinic for patients with localised disease revealed a 15 year crude survival of 64% and after 10 years the disease specific mortality was 2% with the cause specific survival curve plateau at 17 years with no further disease specific mortality ¹²⁸. The group from Duke, Durham demonstrated a significant difference in the failure and death rates between patients with organ confined, specimen confined, and margin positive disease, i.e. failure rates of 12%, 30%, 60%, and death rates of 8%, 12%, and 30% at 10 years respectively ³⁰⁴. A more recent series has shown that the grade of the tumour also affects the rate of progression¹¹³. Therefore cancer control after surgery is affected by tumour grade, and whether the tumour was organ-confined, specimen-confined, or not confined to the specimen ¹¹³. PSA can also be a measure of recurrence however biochemical PSA failure may not manifest itself as clinical progression in the patient ¹¹³.

Other novel approaches have also been advocated, such as high intensity focussed ultrasound ³⁵ and cryotherapy²⁸, but long term data is awaited.

Advocates of screening in Britain suggest there may be a potential benefit in terms of survival if screening is adopted ²⁹⁸, however there is considerable opposition to this in this country. In the USA early detection programmes detect organ-confined disease in a younger cohort of patients ³⁴¹. Long-term

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

follow up of these patients is required to determine if there is an improvement in survival. A study comparing two large fixed cohorts of male Medicare patients, Seattle $n = 94,900$ and Connecticut $n = 120,621$, exposed to different screening regimens concluded that more intensive screening over a eleven year period is not associated with lower prostate cancer specific mortality²³². A recent randomised study from Denmark compared surgery with watchful waiting and initially demonstrated a significant improvement in cancer specific survival in the surgery arm but no improvement in overall survival¹⁶⁹, a further update has now shown a decrease in local progression and a decrease in the development of distant metastasis in the patients randomised to surgery³³.

Using the SEER (surveillance, epidemiology, and end results) data base a study of 104,577 patients with localised prostate cancer concluded that 1) treatment is associated with lower disease specific and overall mortality, and 2) advanced age and black race are markers of disease specific and overall mortality. No conclusion was drawn about the optimal treatment modality²⁰⁶.

A second study using the SEER database and an intention to treat analysis i.e. patients with lymph node dissection and radiotherapy or radiotherapy alone were assigned to the radiotherapy group, and those with a lymph node dissection alone or radical prostatectomy were classified into the prostatectomy group. The advantage of this type of analysis is that it includes

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

patients in the prostatectomy group who have been deselected by pathological staging. The disadvantages are that co-morbidity and tumour-related variables, e.g. tumour size and PSA, are not controlled for. The data suggested that treatment was most useful for high grade tumours, patients with low grade tumour did better than the general population, and the data for surgery may be flattering because surgeons were selecting best patients when analysing data by treatment received ²³³.

The Prostate Cancer Clinical Guidelines Panel has recently published its report on the management of clinically localised prostate cancer ²⁵⁷. The report concluded that the published data to date was inadequate for a valid comparison of the treatment options and, all patients with newly diagnosed, clinically localised cancer should be informed of all the treatment options. The appropriate patient for total prostatectomy should have organ confined disease, an expected longevity that is longer than the natural history of the cancer i.e. 15 years, no significant surgical risk factors and a preference to undergo surgery. The major advantage is the potential for removal of the cancer. The downside includes genitourinary dysfunction and disease progression. The patients most likely to benefit from radiation therapy would have a relatively long life expectancy. No significant risk factors for radiation toxicity and a preference for radiation therapy. In favour of radiotherapy is the fact it is well tolerated and a potential for cure exists. The adverse effects include radiation

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

cystitis, proctitis, and erectile dysfunction. The prostate is still in place and so disease progression may occur. Patients with a short life expectancy and/or a low-grade tumour are likely to be suitable for surveillance. Disease progression is likely with only a marginal compromise to disease specific survival at 5 and 10 years²⁵⁷.

The absence of hard data is a constant source of embarrassment to all health professionals involved in the management of localised carcinoma of the prostate. The Department of Veterans Affairs and the National Cancer Institute's Cancer Therapy Evaluation Programme Co-operative Study (407)- the Prostate Cancer Intervention Versus Observation Trial (PIVOT) - is a randomised trial designed to determine the benefits of early intervention with radical prostatectomy compared to surveillance⁴¹⁸. PIVOT will enrol 2000 patients over a three year period and follow up will be for a minimum of 12 years. Radiotherapy has not been included as a third treatment arm because of sample size, cost, and feasibility. This study should hopefully answer some of the questions.

In the United Kingdom, the three arm randomised ProtecT (prostate testing for cancer and treatment) randomised study is currently recruiting¹⁰⁷. This study has demonstrated the negative connotation patient's place on the phrase watchful waiting, patients interpreted this as no treatment, and has replaced

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

this with the acceptable concept of active monitoring.

The peculiar biology of this disease has contributed to the lack of a consensus about the management of localised prostate cancer. The median time of detection of this disease is 72 years, the median age of death due to prostate cancer is 77 years, since the average life expectancy is 74.2 years, and it is evident that the most men will die in spite of the disease rather than because of it.

Chapter 2: Prostatic Intraepithelial Neoplasia.

The increasing focus of the public has generated considerable interest in the management of prostate cancer. This in turn has resulted in screening programmes being established; not only have these programmes detected prostate cancer but they have detected an increasing number of men with atypical epithelial proliferation the most common being prostatic intra-epithelial neoplasia.

2.1 Prostatic intra-epithelial neoplasia

Bostwick and Brawer⁴⁶ introduced the term Prostatic Intraepithelial Neoplasia (PIN) in 1987, and although initially it was divided into three grades it rapidly evolved into a two tier system – high and low grade¹³⁷. Low grade PIN is of no clinical relevance and the generic term PIN exclusively refers to high grade PIN. In a model analogous to Vogelstein's for colorectal cancer, it has been proposed that PIN refer to the precancerous end of the morphological spectrum. PIN is defined as an abnormal proliferation, where the cellular changes within the prostatic ducts, ductules and acini exhibit the cytological changes of cancer i.e. progressive loss of the markers of secretory differentiation, increasing nuclear and nucleolar abnormalities, increasing proliferative index, neovascularisation, genetic instability, and variation in

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

DNA, but retain a basal cell layer (intact or fragmented). PIN is a spectrum of atypical cytological features ranging from minimal change to those that are indistinguishable from prostate cancer. The most striking features are nuclear and nucleolar enlargement, and are often regarded as diagnostic of PIN. The absence of a basal cell layer distinguishes cancer from PIN. The generic term PIN usually refers exclusively to high grade PIN. Low grade PIN is no longer thought to be clinically relevant.

Prostatic Intraepithelial Neoplasia and Prostate Cancer.




Figure 8: The continuum of change from benign to malignant proposed by Bostwick and Brawer ⁴⁶. The loss of the basal cell layer is pathognomonic of cancer, but the the cytological changes of cancer occur in prostatic intraepithelial neoplasia which has an intact basal cell layer. The presence of the basal cell layer can be confirmed with the use of antibodies against high molecular weight cytokeratins.

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

There are four main architectural types of high grade PIN: tufting, micro papillary, cribriform, and flat ⁴⁵.




Fig 9: Tufting PIN (low power).




Fig10: Micropapillary(low power).



Fig 11: Cribriform PIN (low power)




Fig 12: Flat PIN (low power).

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

These different architectural types do not appear to have any clinic pathological significance. PIN spreads through the prostatic ducts in three different patterns ⁴⁴:

- i) Neoplastic cells replace the normal luminal secretory cells,
- ii) Direct invasion of the duct wall,
- iii) Pagetoid spread, the cells grow between the secretory and basal cells.

Clinical studies suggest that PIN precedes carcinoma by ten years, with low-grade PIN first emerging in men in the third decade ³³¹ and the peripheral zone is the most common location for PIN. The frequency of PIN in prostates with adenocarcinoma is much greater than in prostates without cancer ³¹¹.

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

	Low-grade PIN (PIN 1)	High grade PIN (PIN 2 and 3)	Cancer
Architecture	Epithelial cell crowding and stratification, with irregular spacing.	More crowding and stratification. Four patterns: tufting, micro papillary, cribriform, and flat	Crowding with stratification
Cytology: Nuclei	Enlarged with size variation	Enlarged with size and shape variation.	Hyper chromatic nuclei with macro nucleoli.
Cytology: Chromatin	Normal	Increased density and clumping	Increased density and clumping
Cytology: Nucleoli	Rarely prominent	Frequently large and prominent	Large and prominent
Basal Cell Layer	Intact	May show some disruption	Absent
Basement Membrane	Intact	Intact	Absent

Table 4: Summary of relationship between prostatic intraepithelial neoplasia and cancer.

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

2.2 Incidence of high-grade PIN in benign prostate tissue (biopsy and TURP)

The incidence of PIN varies in benign needle biopsy specimens between 0.7% and 16.5%. In TURP specimens it is between 2.8% and 3.2%. The incidence of PIN is lower in screened populations than in patients presenting for urological assessment.

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

Ref./Year	No of Patients	Tissue Specimen	Patient	Incidence of PIN /%
²¹⁸ /1989	256	14 G needle biopsy	Non-screening	11
²⁵³ /1991	330	18 G needle biopsy	Screening	5.2
³²¹ /1994	163	18 G needle biopsy	Non-screening	8.6
⁴⁸ /1995	200	18 G needle biopsy	Non-screening	16.5
⁴⁸ /1995	200	18 G needle biopsy	Non-screening	9.5
²¹⁶ /1996	1275	18 G needle biopsy	Non-screening	4.4
⁴¹⁷ /1997	439	18 G needle biopsy	Non-screening	5.5
³⁶⁰ /1997	79	14 G needle biopsy	Non-screening	7.6
³⁰⁵ /1997	148	18 G needle biopsy	Non-screening	14.1
⁸⁰ /1997	1009	18 G needle biopsy	Non-screening	1.5
¹²⁴ /1997	158	TURP	Non-screening	3.2
²⁹⁶ /1997	570	TURP	Non-screening	2.8
³¹⁸ /1998	2219	18 G needle biopsy	Non-screening	4.0
¹⁷² /1998	388	18 G needle biopsy	Non-screening	8.0
³²⁰ /1998	795	18 G needle biopsy	Non-screening	7.4
²⁹⁴ /1998	62537	18 G needle biopsy	Non-screening	4.1
²⁸⁷ /1999	15753	18 G needle biopsy	Non-screening	3.9
¹⁶⁶ /1999	8763	18 G needle biopsy	Screening	.7
³³² /2000	3891	18 G needle biopsy	Non-screening	5.4
⁸ /2000	485	18 G needle biopsy	Non-screening	6.8
¹⁷⁰ /2001	1474	18 G needle biopsy	Screening	4.7
¹²⁰ /2001	1961	18 G needle biopsy	Screening	4.7
²⁸⁴ /2004	1086	18G needle biopsy	Non-screening	4.0
²⁶⁷ /2005	1188	18G needle biopsy	Non-screening	2.7

Table 5: The incidence of PIN according to recent studies.

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

The incidence and extent of PIN appears to increase with age, it is more prevalent and extensive in patients of African origin¹²⁰. PIN is also detected a decade earlier in men of African origin compared to Caucasians¹²⁰.

2.3 Location of PIN

PIN lesions are often multifocal, and bilateral, as is prostate cancer. PIN and prostate cancer are principally located in the peripheral zone of the gland. In radical prostatectomy specimens with cancer, high grade PIN is present in the transition and peripheral zone simultaneously in 36% of cases, and is found exclusively in the transition zone in 1% of cases³¹¹.

2.4 PIN and serum PSA levels

This is a contentious area; a study of 81 men undergoing surgery for benign disease found that in men with PIN alone the mean serum PSA fell between the levels for men with benign and malignant disease⁵⁰. The same authors found similar results for patients with BPH, PIN, and cancer on TRUS sextant biopsy⁵². However, Ronet et al³²⁶ and Alexander et al⁶ did not confirm this finding. In a recent study of 81 consecutive men who have undergone radical retro pubic prostatectomy it was shown that PIN did not contribute to total or percent free PSA levels²⁶⁹. Using grid counting to calculate the volume of tumour, PIN, and benign disease, in a study of 194 patients PIN volume was

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

shown to correlate to PSA, this result was interpreted by the authors as an artefact and they concluded that PIN did not contribute to it⁶. Immunohistochemical studies reveal that the expression of PSA in PIN is less than that observed in BPH and cancer²⁴⁹. The explanation for a lower PSA in patients with PIN is that PIN is an abnormality within pre-existing ducts and acini, and cytoplasmic secretory products would be expected to wash downstream rather than escape into the stroma and blood vessels.

2.5 PIN and Radiotherapy.

A study of 29 patients who underwent salvage prostatectomy after radiotherapy demonstrated PIN in only two of the specimens. After radiotherapy the basal cell layer in PIN becomes more prominent and the presence of cytoplasmic blebs distinguishes secretory cells from basal cell.⁷⁷.

2.6 Clinical Significance of PIN

High grade prostatic intraepithelial neoplasia has a high predictive value as a marker for adenocarcinoma, Davidson et al evaluated this in a retrospective case control study of 100 patients with needle biopsies with high grade PIN and 112 biopsies without PIN matched for clinical stage, patient age, and serum PSA. Adenocarcinoma was identified in 35% of the subsequent biopsies with PIN compared with 13% in the controls. The likelihood of adenocarcinoma increased with time, 32% at one year, and 38% after one year.

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

The presence of high grade PIN and a raised PSA are associated with adenocarcinoma in 51% of cases in repeat biopsies ¹⁹². Others studies have found cancer detection rates of 28% ²¹⁶ and 47% ³⁵⁰.

The incidence of prostate cancer after an initial diagnosis of PIN increases if a patient has repeat biopsies²⁰⁷. In a series of 245 patients with an initial diagnosis of PIN, cancer was diagnosed in 24.5% of patients on the initial repeat biopsy and a total of 32.2% after the second repeat biopsy ²⁰⁷.

Recent work has also highlighted the importance of second biopsies with a diagnosis of PIN. In a cohort of 190 men diagnosed with PIN and having serial repeat biopsies the overall incidence of prostate cancer on repeat biopsies was 30.5%, but if there was a second biopsy with PIN detected, the risk of cancer increased to 41%, compared to only 18% if the second biopsy had benign tissue¹³⁵.

PIN is often associated with moderately to poorly differentiated adenocarcinoma of the prostate; it is not often associated with low-grade prostate cancer or cancer arising in the transition zone ²⁶³.

2.7 Evidence linking PIN and Cancer.

There is no disruption of the basal cell layer in benign disease, where as in

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

cancer the basal layer is absent. In PIN there is partial disruption. There is increased expression of the proteolytic enzyme type IV collagenase in cancer and PIN compared to benign disease²⁶⁴.

The number of stromal micro vessels in PIN is greater than in benign or BPH tissue, but less than in adenocarcinoma. The micro vessels in PIN are shorter than those in benign epithelium, with irregular contours and open lumina, an increase in the number of endothelial cells, and a greater distance from the basement membrane²⁶⁴.

The doubling time of PIN is at an intermediate level when compared to BPH and cancer²⁶⁵ and there is a progressive increase in the number of mitotic figures and apoptotic bodies from BPH through to PIN and finally cancer²⁶⁶.

The cell kinetics of PIN have been studied. It has been shown that PIN, which is not associated with carcinoma, shows a down regulation in apoptosis, which is common with prostate cancer, but distinct from PIN which is associated with prostate cancer²⁰⁰

Many markers of differentiation are down regulated in PIN and cancer, e.g. neuroendocrine cells⁴⁷, prostatic acid phosphatase, and prostate specific⁹⁶ antigen. Benign prostate has most cells with neuroendocrine differentiation.

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

Others are up regulated e.g. fatty acid synthetase²⁷⁷, glycoprotein A-80¹³⁶, glycoprotein TAG-72⁵⁴, and Lewis X and Lewis Y antigen²⁷⁶.

Using micro array analysis hepsin is up regulated in PIN, and prostate cancer but is absent in benign tissue¹⁰¹, in hormone refractory disease it has an intermediate level of staining. Hepsin has been shown to be a predictor of relapse¹⁰¹. PIM1 kinase was up regulated in all stages of prostate cancer but absent in PIN and benign disease¹⁰¹.

Transgenic mice have been used to study the pathogenesis of prostate cancer. The most extensively used model is the TRAMP (transgenic adenocarcinoma mouse prostate) model^{130 138}. The simian virus 40 (SV40) early region tumour antigens have the ability to induce transformation in vivo. The large T antigen (Tag) interacts with the retinoblastoma gene (Rb) and p₅₃, and the small t antigen interacts with a protein phosphatase. The loss of wild type Rb and p₅₃ was used in this model. Probasin was used to target the SV40 to the prostate. Probasin encodes for an androgen and zinc related protein specific to the dorsolateral epithelium of the prostate. A probasin SV40 Tag construct was used to target the prostate of mice¹³⁸. The expression of SV40 early genes in the prostate of the mice results in mice displaying progressive forms of prostate disease. The mice develop prostatic hyperplasia at 10 weeks and invasive adenocarcinoma by 18 weeks of age. By 10 to 12 weeks TRAMP

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

mice reproducibly displayed evidence of high grade PIN¹²⁹. By 30-36 weeks all TRAMP mice developed metastases¹³⁰. This model has therefore facilitated in the understanding and characterisation of the sequence of events, which may occur in the initiation, and progression of prostate cancer. This model is however open to criticism, since in the human form of the disease benign prostatic hyperplasia does not progress to adenocarcinoma.

2.8 Differential diagnosis of PIN

There are several benign conditions that can mimic PIN; these include atypia induced by inflammation, infarction, or radiation; transitional cell metaplasia; basal cell hyperplasia; clear cell cribriform hyperplasia; normal ejaculatory duct and seminal vesicle epithelium. Malignant lesions, which have to be distinguished from PIN, include high grade transitional cell carcinoma involving the prostate ducts and acini, and cribriform acinar, and cribriform ductal prostatic carcinoma²⁹⁹.

Despite its major impact as the most common cancer in men, prostate cancer is relatively poorly understood at the genetic level. It is hypothesised that, as with other neoplasms, mutations of proto-oncogene and tumour suppressor genes contribute in some way to the genesis of prostate cancer. The use of molecular markers to supplement clinical information concerning the biological aggressiveness of a carcinoma may allow better selection for

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

different management strategies. Alternatively, they may be used after surgery to stratify men into groups whereby those with poorer prognosis may receive adjuvant treatment, perhaps providing an advantage of improved disease specific survival.

Chapter 3: P₅₃, Bcl₂, Bax- α , and Ki-67.

3.1 P₅₃

The p₅₃ gene is located on the short arm of chromosome 17 (17p13.1). The gene is composed of 11 exons. The p₅₃ gene product is a 53Kda (393-amino acid) nuclear phosphoprotein²⁰³, which inhibits the advancement of cells from the G1 to the S phase. Lane and Crawford first identified this in 1979²¹⁴.

The DNA tumour virus SV40 expresses two antigens, small t and largeT. These are essential for the virus to transform cells. Lane and Crawford demonstrated that the T antigen binds to a 53 Kd host protein, and precipitated this out. They postulated that this protein was not virally coded and that it could be detected in cells transformed by other viruses, radiation, chemical carcinogens, or even in untransformed cells. They further precipitated the protein from polyoma virus transformed cell line. The authors postulated that this protein may have an important role in the modulation of the transformed state, and it may normally act as a regulator of cellular functions related to growth. Lane and Crawford concluded that it is of prime importance to determine the level of this protein in untransformed cells or normal tissue and to see if it is induced by other carcinogens²¹⁴. The importance of this discovery was not clear early on, since these viruses do not cause human

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

cancers.

From its humble beginnings, the influence of p₅₃ rapidly grew. The number of publications with p₅₃ in the leading role increased exponentially every year from 1979. In 1993 p₅₃ was named as Science magazines Molecule of the Year²⁰³. In the editorial Dr Koshland commented that the early detection of p₅₃ may lead to treatment by known methods such as surgery, and stressed the importance of basic research - “Basic research does not mean of no practical value”²⁰³. This is also the aspiration of this piece of research.

Prostatic Intraepithelial Neoplasia and Prostate Cancer.



Fig13: Cover of Science Dec 1993.

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

A mutant p_{53} gene synthesises a defective p_{53} protein that lacks the ability to regulate cell proliferation. The p_{53} protein is found in small quantities in normal cells, but in large quantities, 5 to 100-fold increase, in transformed cells both in vivo and vitro. Inactivation of p_{53} is the single most common mutation in human cancer²²⁴. The presence of p_{53} nuclear reactivity is a predictor of recurrence after definite local therapy i.e. radiotherapy or radical prostatectomy¹⁴². P_{53} and Ki-67 have been shown to be independent prognostic markers of recurrence²⁷⁴.

3.11 Function of p_{53} -Guardian of the genome²¹³.

The wild type p_{53} is capable of forming a homo oligomer, which binds to specific DNA sequences and act as a transcription factor. Insults to cellular DNA, such as exposure to ultraviolet light, or radiation results in an increased levels of the p_{53} protein in normal cells leading to arrest in the G1phase (the growth phase of the cell cycle after mitosis and prior to DNA synthesis), allowing repair of the DNA prior to replication of the DNA in the S-phase of the cell cycle. If DNA repair cannot be accomplished, programmed cell death (apoptosis) may occur. Thus p_{53} is thought to ensure genomic integrity. The p_{53} gene does not directly establish integrity of the genome but acts as a

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

transcription factor to increase expression of other genes that will initiate the cell cycle arrest and/or apoptosis. Apoptosis is genetically programmed cell death program, which functions as a suicide pathway wherein individual cells kill themselves when triggered to do so. Apoptotic death is distinct from necrotic death.

It has been suggested that p₅₃ regulates the transcription of at least two sets of genes; one is responsible for transient cell arrest in G1 and the other controls the initiation of apoptosis. Both of these processes eliminate potential oncogenic mutations, either by DNA repair or by inducing damaged cells to commit suicide.

This response has been lost in many tumour cells as their p₅₃ genes have either been mutated or had its activity neutralised through the production of proteins that bind to it and neutralise it. Lack of p₅₃ function in tumour cells is thought to be responsible for the observed genomic instability of these cells as manifested by aneuploidy and the ability to undergo gene amplification. However, the p₅₃ gene does not seem to be essential for normal development but lack of p₅₃ function confers an elevated risk of developing cancer – Li Fraumeni syndrome (see next section).

Initially the p₅₃ gene was thought to be an oncogene but later studies showed

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

that the p₅₃ clones that had been studied were in fact mutated versions of the gene, and these mutant alleles have properties which are different from those of the wild type p₅₃ gene. Current evidence supports its classification as a tumour suppressor gene.

3.12 Evidence supporting wild type p₅₃ as a tumour suppressor gene:

- 1) The murine wild type p₅₃ gene can suppress the transformation of rat embryo fibroblasts in cell culture by known oncogenes e.g. *ras* and adenovirus E1B. The introduction of wild type gene or cDNA into a transformed cell culture stops cell growth at the G1 phase of the cell cycle following mitosis¹³⁴.
- 2) The wild type p₅₃ gene is capable of reverting the transformed phenotype of human cancer cells and osteosarcoma cells in an animal model²⁸⁰.
- 3) Mutations and allelic loss of the p₅₃ gene have been associated with tumours from a wide variety of human organs and tissues, e.g. colon⁶⁷, lung¹⁸¹, oesophagus⁷⁰, breast¹⁸, ovary²²³, brain³⁹⁵, and haematopoietic tissue.
- 4) The Li-Fraumeni familial cancer syndrome⁴⁰⁰, characterised by diverse mesenchymal and epithelial neoplasms at multiple sites – breast and adrenocortical carcinomas, sarcomas of soft tissue and bone, brain tumours and leukaemia, is associated with the presence of germ line

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

mutations in the p₅₃ gene⁴⁰⁰.

- 5) Transgenic mice deficient for p₅₃ or carrying altered p₅₃ gene are significantly more prone to the spontaneous development of a variety of neoplasms⁴¹⁶.
- 6) Oncoproteins from several DNA tumour viruses e.g. the SV40 large T antigen, the adenovirus E1B, and papillomavirus E6 protein, were found to functionally inactivate p₅₃ by binding to or degrading the wild type p₅₃ protein⁸⁷.

3.13 Mechanism of action of p₅₃.

- 1) Up regulation of transcription of the bax gene and down regulation of bcl₂²⁶¹.
- 2) Up regulation of IGF-BP3 gene expression. IGF-BP3 blocks IGF induced signalling by binding to IGF. Inhibiting IGF promotes apoptosis⁶⁴.
- 3) Up regulation of FAS/APO-1 gene expression. The cell surface receptor Fas mediates apoptosis²⁷⁹.
- 4) Activation of G proteins. Cdc42 is a GTPase, and p₅₃ promotes apoptosis through up regulation of cdc42 mRNA and protein, up regulation was followed by cytoplasmic to membrane translocation of this G protein. One action of Cdc42 is phosphorylation and so inactivation of Bcl₂³⁸⁶.

3.14 Inactivation of p₅₃ function

A mutation of p₅₃ gene is the most common mechanism of inactivation of p₅₃ function¹⁶⁸. DNA mutation can result from external environmental factors or internal errors in mechanisms involved in nucleic acid metabolism³⁶¹. In the first case, the mutagenic agent determines the nature of the lesion. In the second case, mutations appear to be spontaneous. Analyses of the mutational spectrum of p₅₃ in human tumours revealed that the majority of p₅₃ mutations in human cancers are missense mutations, giving rise to an altered protein⁷⁵. At least four mutation "hot spots," have been identified in a variety of human neoplasms. Mutations in these hot spot codons account for approximately 30% of all p₅₃ mutations.

In addition to mutation, inactivation of p₅₃ function can occur by other mechanisms, including binding of p₅₃ to another protein or increased degradation of p₅₃. The E1B protein of adenovirus binds to p₅₃ and prevents p₅₃ activated transcription. The E6 protein of some types of human papilloma virus (HPV) binds to p₅₃, causing p₅₃ to be marked for degradation¹⁷¹. Thus, HPV infection can lead to increase p₅₃ degradation. Only E6 proteins from those strains of HPV that have been strongly associated with anogenital cancers (HPV 16 and HPV 18 are the most common) result in accelerated p₅₃ degradation.

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

Another mechanism by which p₅₃ is inactivated is interaction with endogenous cellular proteins. In sarcomas, there is amplification of the MDM2 (mouse double minute) gene, located at chromosome 12q13-14, resulting in high levels of its 90-kD nuclear protein product²⁹¹. Binding of the MDM2 protein to p₅₃ is thought to inactivate its transcriptional activity²⁶². The binding occurs at a site distinct to the T Ag binding site, and in tumours where there is up regulation of MDM2, mutation of wild type p₅₃ is not necessary for transformation²¹³.

3.15 Detection of p₅₃ gene and product

The methods currently used in the detection of p₅₃ alterations are outlined below. Different methods have different advantages and disadvantages, which are discussed.

3.16 Immunohistochemical analysis

In 1982 Benchimol et al²³ demonstrated by radio-immunological assay that the p₅₃ protein was specifically over-expressed in transformed cells and 'undetectable' in normal cells. Numerous studies have confirmed these observations and proved that p₅₃ protein accumulation is a consequence of its stabilisation. Wild type p₅₃ (normal) product has a short half-life of about 6-20 min and normally does not accumulate in amounts detectable by conventional immunoprecipitation or immunohistochemical methods¹⁸. Mutations in the

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

gene result in a product (mutant p₅₃) with a much longer half-life of up to 6 hours²²⁴. Therefore, the findings of intense nuclear staining with anti-p₅₃ antibody is indicative of a point mutated p₅₃¹⁸. Early data from direct DNA sequencing confirms that mutations were present in cases that over expressed the protein and were absent in tumours that did not over express the protein¹⁸
24:146

Immunostaining studies of tumour cells require antibodies with very high specificity, absence of cross-reactivity with another cellular protein, high affinity for the antigen regardless of the method of tissue fixation. Almost all archival tissue in histopathology departments all over the world is preserved by formaldehyde and embedded in paraffin blocks. This has led to the production of a new panel of monoclonal antibodies (MAbs) specific for human p₅₃, such as DO1, D07 and HR231^{219:407} which are very efficient on this type of material. All these monoclonal antibodies are currently available commercially.

At present, immunohistochemistry is the most widely used technique for assessing p₅₃ alterations in human cancers because it is simple and rapid³⁶¹. From the clinical standpoint p₅₃ immunohistochemistry appears to be the most practical and useful method for detection of p₅₃ alterations in pre-cancerous lesions and carcinomas. This technique allows precise localisation and

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

identification of the cells that exhibit p₅₃ alterations, however it cannot detect the complete loss of p₅₃ or non-sense mutations that introduce premature stop codons into the gene⁷⁵. Moreover point mutations in the p₅₃ gene are not the only mechanism by which the p₅₃ protein can be stabilised. The normal p₅₃ protein can be stabilised by the action of viral and cellular oncoproteins. Treatment of cells with DNA-damaging agents (preoperative radio and chemotherapy) increases the level and stability of the p₅₃ protein as well¹⁵³. Therefore, interpretation of immunohistochemically positive staining with p₅₃ must include consideration of the mechanisms that underlie stabilisation of the p₅₃ protein.

A comparative study carried out on 19 archival specimens of colorectal cancer and comparing six commercially available antibodies – Bp53-12(BioGenex), 1801 (Oncogene Science), D07 (Novocastra), 240(Novocastra), CM1 (Novocastra), and Signet (Signet) with results obtained by PCR and double-stranded DNA sequencing demonstrated that D07 gave significantly the best results. When D07 was used with an antigen retrieval system and the cut of for high expression was a labelling index of 30% the sensitivity was 67%, the specificity was 90%, the positive predictive value was 86%, the negative predictive value was 75%, and the efficiency was 79%. If the cut of for the labelling index was lowered to 1% the sensitivity was 89%, the negative predictive value was 86%, specificity was 60%, and the positive predictive

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

value was 67%, with an efficiency of 74%¹³.

3.17 Molecular analysis (DNA sequencing)

DNA sequencing determines the exact mutational event that modified the p₅₃ gene. This knowledge is mainly of use in molecular epidemiological analysis. It is generally assumed that a decrease in genetic stability of the genome is involved in the accumulation of a multitude of genetic changes leading to the formation of a neoplastic cell. As mentioned before, the type of p₅₃ mutation depends on the event, which affects the DNA (external or internal). Genetic assay is the most accurate technique for identification of p₅₃ mutations but it is generally laborious and requires specially equipped molecular laboratories.

3.18 Serological analysis

Several studies have demonstrated the presence of p₅₃ auto antibodies in patients with carcinomas^{68:91;98:419}. The immune response was found only when p₅₃ accumulation was detected in the tumour cell, but mutation or over-expression did not automatically lead to a p₅₃ immune response and there was no correlation with the location of the mutation^{212:342}. Several laboratories have developed ELISA procedures. Despite this, serological analysis of p₅₃ alterations in human cancers is still in its infancy³⁶¹. Progress in this field will be of great importance, because this technique can be easily performed for

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

routine diagnosis and does not require tumour samples.

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

3.20 P₅₃ and human cancers

Mutations in the p₅₃ gene are now regarded as one of the commonest specific genetic changes found in human cancer⁷⁵. More than a half of human malignancies derived from the epithelial, mesenchymal, haematopoietic, and lymphoid tissues, as well as the central nervous system, analysed thus far, have an altered p₅₃ gene^{17:89;146:168}. This has led to a concentration of efforts to uncover the value of p₅₃ as an independent marker for tumour detection and diagnosis, prognosis and progression.

3.21 P₅₃ as an independent marker for tumour detection and early diagnosis

Mutated p₅₃ protein might be a good target for cancer detection and diagnosis. As discussed before this is primarily due to many missense mutations inducing changes which prolong the half-life of the p₅₃ protein allowing immunohistochemical detection. P₅₃ positive immunostaining has been observed in pre-cancerous lesions of the colorectum^{190:307} stomach³⁵⁷ bronchus^{288:362} larynx¹⁰⁵ oesophagus^{25:410} and in head and neck premalignancies⁴²⁰. Although p₅₃ mutation seems to occur at the earliest clinically detectable stages of the neoplastic process in some cancers, the fraction of tumours with an altered p₅₃ gene is generally higher in the late stage of tumours^{114:119}.

3.22 P₅₃ as an independent marker for tumour prognosis

Altered p₅₃ function has been examined as potential prognostic factors in a variety of cancers. An ever increasing number of studies are reporting that altered p₅₃ is an independent predictive factor of unfavourable prognosis. Aberrant forms of p₅₃ protein are correlated with more aggressive tumours, early metastasis and lower 5-year survival rates.

In breast cancer p₅₃ over expression (with either demonstrated or presumed mutation) has been associated with late stage, metastatic spread, and low progesterone receptor content ⁹⁷. In other studies it was associated with a shortened survival ^{182;327;336;359;373;390} and shortened metastasis free survival ³⁹¹. A consistent picture is now also emerging correlating the loss of p₅₃ function with shortened survival in patients with cancers of the colon ^{67;272;380;427} lung ^{181;268} oesophagus ⁷⁰ stomach ¹¹⁸ ovaries ²²³ and bladder ³⁴⁹ as well as soft tissue sarcomas ¹⁰⁸.

3.23 P₅₃ as an independent marker for tumour progression

The role of p₅₃ mutations in the progression of astrocytomas to glioblastomas ^{147;358}, in the evolution of chronic myelocytic leukaemia to blast crisis ^{119;408} and in primary as compared with metastatic cervical cancers ⁹² has been demonstrated. Genetic analysis showed the presence of p₅₃ mutant cells in low numbers in primary tumours and the largely mutant composition of the more

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

advanced tumours as well as their metastases^{147:358:408}. In the pathogenesis of colon carcinoma point mutations typically arise during the conversion of a benign adenoma to an invasive adenocarcinoma¹¹⁴. The molecular mechanisms that underlie tumour progression are thought to include the clonal expansion of cells that previously acquired a mutation in cancer-related genes. A cell that carries a specific change in a critical gene might acquire a selective growth advantage and become the dominant cell type as the tumour progresses. This hypothesis is also supported by the repeated discovery of the mutant p₅₃ preferentially in anaplastic and undifferentiated tumours^{106:183:189} e.g. in thyroid carcinomas in which the p₅₃ mutants are primarily confined to undifferentiated tumours. The colorectal tumour progression model continues to be the paradigm for all tumour progression.

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

3.3 P₅₃ and Prostate Cancer.

A literature review was carried out using the Medline database. The initial keyword used was p₅₃ and the second was prostate neoplasms. For the first keyword a total of 16282 references were detected and for the second 32852. These two searches were combined and more manageable 297 references were detected. These were further limited to English language and human and the number of references coned down to 280. The abstracts were reviewed and those that were felt not to be pertinent were omitted, a total of 90 references were reviewed in detail

10:11;19-21;29;32:42;55;56;60;63;65;66;74;75;78; 79;81;100;103;110;112;121;123;139;142;151;161;162;173-176;180;187;193;194;201;209;210;215;220;236;238;238-241;251;254;259;273;274;278;283;295;312;314;315;334;341;364;368;369;371;372;374;375;377;387;389;396;397;399;406;413;423;425;426

Fifteen years ago the first study linking the inactivation of the p₅₃ gene and prostate cancer was published¹⁸⁰. Since this study there has been an explosion in research activity associated with p₅₃ and prostate cancer. In this seminal study by Isaacs et al it was reported that mutations occurred in the p₅₃ gene in prostate cancer cell lines and human tissue specimens. Sequence analysis of exons 5-8 of the p₅₃ gene reveals that three of five prostate cancer cell lines contain mutations which alter the base sequence of this portion of the gene. One of two primary prostatic cancer specimens examined also contained a mutation in this region. Transfection of the wild-type p₅₃ gene versus a mutated p₅₃ gene into two cell lines with p₅₃ mutations results in reduced

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

colony formation. Immunocytochemical data indicate that prostate carcinoma cells expressing the transfected wild-type p₅₃ gene are growth arrested because they exhibit a reduced level of thymidine incorporation into DNA.

The first paper linking p₅₃ and prostate cancer specimens from a patient appeared in 1992 from the group from North Carolina¹¹². Using PCR techniques they detected p₅₃ mutations in a patient with T₄N₁M₀ prostate cancer who had initially been treated with a multimodality combination of radiotherapy and hormonal therapy. The patient 14 years after the initial treatment went on to have a TURP and radical cystoprostatectomy. The areas of the primary tumour and the lymph nodes were examined with PCR techniques. Although the mutations in p₅₃ were heterogeneous in the primary tumour, the same mutations were detected both in the primary tumour and the nodal deposits.

3.31 P₅₃ in patients who have undergone surgical treatment.

Radical prostatectomy has become the main stay for the surgical management of localised prostate cancer; however it is only successful if the patient is cured. P₅₃ has been advocated as a predictor of outcome. One of the earliest studies of the incidence of p₅₃ in patients with prostate cancer consisted of 150 patients. The study group consisted of 88 men who had had a radical prostatectomy, 52 cases had a TUR, and 10 cases had a TUR for palliation of

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

recurrent disease. P₅₃ was detected using immunohistochemistry in 17 of the 140 primary tumours, and in two patients with recurrence⁴². It is noteworthy that an antigen retrieval system was not used in this earlier study. Immunostaining was significantly related to stage. The immunohistochemical findings were confirmed in 14 cases by DNA amplification and screening between exons 4-9. 20 negatively staining cases were used as controls and no mutations were detected in these cases. The authors used the fact that single strand conformational polymorphism may not be 100% sensitive, and p₅₃ accumulation may occur without p₅₃ mutation as an explanation for the discrepancy in the results obtained between immunohistochemistry and single strand conformational polymorphism. The authors also commented upon the fact that compared to other tumours – breast, lung, colon, and oesophagus, the frequency of p₅₃ mutation and accumulation was rather low⁴².

In a retrospective study of 49 patients with clinically localised (T₁₋₂ N_x M₀, PSA < 20, bone scan negative) moderately differentiated (Gleason score 5-7) carcinoma of the prostate treated with radical prostatectomy and with a follow up of at least 5 years, Yang et al⁴²⁵ showed that p₅₃ determined by immunohistochemical was present in 10/16 patients which had a recurrence compared to 7/33 in the non-recurrent group. P₅₃ was significantly associated with cancer recurrence. It is interesting to note that in this report 22 out of the 49 patients that had extra capsular disease and the incidence of nodal

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

involvement is not mentioned.

In a study of 93 patients from Memorial Sloan-Kettering Cancer Centre, 19/93 (20%) demonstrated nuclear immunostaining¹¹. Of the 93 tumours, 48 were primary tumours, 29 were lymph nodes involved with tumour, and 16 were specimens from patients who had presented with pathological fractures. Of the 48 primary tumours, 9 were organ confined, 18 had seminal vesicle involvement, at the time of the operation 11 had lymph node involvement, and 10 were from patients with disseminated hormone refractory disease. Apart from the 10 patients with metastatic disease who had undergone palliative TURPs, the remaining 38 primary tumours were from patients who had undergone radical prostatectomy. 9 out of the 48 primary tumours stained positively for p₅₃, of these 9, 8 had Gleason scores above 8. Only one of the primary tumours with a Gleason score of less than 7 was observed to accumulate p₅₃. None of the specimens with organ confined disease stained positively for p₅₃, but 1 out of the 18 with seminal vesicle involvement, and 2 out of the 11 patients with pelvic lymph node metastasis stained positively for p₅₃. Eight out of the 16 bone specimens compared to two out of the 29 lymph nodes stained positively for p₅₃. Therefore p₅₃ accumulation significantly correlated with high Gleason score, stage, hormone refractory disease, and was more common in bone metastasis than lymph node metastasis. A study of 100 consecutive radical prostatectomy specimens confirmed the low incidence of p₅₃ mutations

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

in localised carcinoma i.e. one in 67 cases, compared to 5 out of 33 cases in locally invasive carcinoma ²⁷³. This study also demonstrated a good correlation between immunohistochemically detected p₅₃ over expression compared to p₅₃ mutations detected by polymerase chain reaction amplification and single strand conformation polymorphism. The DO-1 antibody detected 5 out of 6 mutations in p₅₃ and an additional mutation in exon 6 was detected by molecular biology ²⁷³.

A further study of 73 radical prostatectomy specimens demonstrated that 34 specimens stained positively for p₅₃ ¹⁶². It is interesting to note that only 28 out of the 73 prostatectomy specimens were in fact organ confined i.e. 39%. 24 out of the 28 pT₂ or less lesions were p₅₃ negative, thus staining for p₅₃ correlated with high stage. In addition p₅₃ staining correlated with increased PSA, Gleason score and increased tumour volume. This correlation is not surprising since high stage disease is associated with increased PSA, high Gleason score and high tumour volume. Follow up was not an aspect of this study and this paper did not comment upon whether p₅₃ staining was an independent marker for recurrence. A further study of 137 consecutive prostate specimens obtained by radical prostatectomy detected 11 specimens (8%) with p₅₃ over expression, ten out of the 11 specimens were pT3, nine out of the 11 were high grade i.e. Gleason score >7. P₅₃ over expression was associated with increased Ki-67 labelling index implying rapid tumour

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

proliferation⁶⁰.

A study of 85 prostate carcinoma specimens obtained from a mixture of radical prostatectomy specimens and TURP specimens analysed by immunohistochemistry and DNA analysis demonstrated only 3 specimens with aberrant p₅₃ expression i.e. 3.5%⁴⁰⁶. No information was given regarding the stage of the disease; the Gleason score of the specimens was 6, 7, and 8. The authors concluded that the presence of aberrant p₅₃ appears to be an infrequent event. The infrequency of p₅₃ in localised prostate cancer was further confirmed on a study of tissue obtained from 37 patients with localised carcinoma of the prostate and subsequently treated with external beam radiotherapy, only one of the 37 specimens stained positively for p₅₃¹⁵¹. This study went on to compare immunohistochemistry with PCR using both positive and negative samples and found complete concordance. These authors concluded that the presence of p₅₃ mutation was a late step in the progression of prostate cancer and was a late step in the progression of prostate cancer and associated with advanced disease, dedifferentiation, and the acquisition of androgen independence.

The studies so far mentioned in this section have largely concentrated upon detecting the presence of p₅₃, and correlating it with known markers of poor outcome, i.e. high Gleason score, high tumour volume/stage. The group from

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

Walter Reed Army Medical Centre, Washington, then published the outcome of their series of radical prostatectomy operations ^{19:20:274}. The series consisted of 175 patients who had had radical prostatectomy over a seven-year period and who they had archival material available on. Using immunohistochemistry 114 of the 175 patients (65%) were positive for p₅₃, a much higher proportion than previously detected. Expression correlated with higher stage, Gleason score, and nuclear grade. P₅₃ positivity did correlate with a higher incidence of recurrence, i.e. 51.1% in p₅₃ positive patients versus 22% in p₅₃ negative patients. This was an independent prognostic marker ¹⁹. Over expression of bcl₂ was detected in 47 patients (26.9%), the combination of p₅₃ positive and bcl₂ positive was highly significant at predicting recurrence when compared to p₅₃ negative and bcl₂ negative patients. Similar results have been reported by others, in a series of 40 radical prostatectomy specimens, 32 (80%) stained for p₅₃ and of these 20 (62.5%) progressed, none of the p₅₃ negative patients progressed. P₅₃ immunoreactivity was an independent predictor of recurrence ³⁷⁴. In a British series of 76 radical prostatectomies carried out by the Bristol group with a mean follow-up of 38 months, aberrant p₅₃ was detected in 39 (57%) of patients. 19 of these patients had recurrence of prostate cancer, where as in the group with normal expression of p₅₃ only three out of 23 patients had evidence of recurrent disease. Therefore the authors concluded that aberrant p₅₃ does predict recurrence ⁵⁵. In a further series of 76 radical prostatectomies, aberrant p₅₃ was detected in 18 patients (20% positive

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

staining) and this significantly correlated with poor survival in a multivariate analysis²⁰⁹.

A heterogeneous series of patients treated with radical prostatectomy, some having had LHRH therapy preoperatively, from the Memorial Sloan-Kettering demonstrated a strong link between p₅₃ over expression and recurrence²⁹⁵. However they also found that some tumours expressed p₂₁. There was no relationship between p₂₁ and p₅₃ status, this is interesting since p₂₁ is thought to be a downstream messenger of p₅₃, and one would expect only patients with wild type p₅₃ to have expression of p₂₁. The authors postulated that p₂₁ may also be activated by a p₅₃ independent pathway²⁹⁵.

A study from New South Wales carried out on 263 consecutive radical prostatectomy attempted to quantify the relationship between p₅₃ and outcome after surgery³¹². The authors compared the number of cell with aberrant p₅₃ with outcome, and also compared clustering of p₅₃ i.e. the presence of a cluster of p₅₃ positive tumour cells within a 200 x magnification field, with outcome. They concluded that their data supported the relationship between outcome and p₅₃ immunohistochemistry at thresholds of no nuclear accumulation versus any accumulation; and the presence of clusters of p₅₃ positive cells versus their absence; and between >20% positive cells and <20%staining. In predicting early death from prostate cancer p₅₃ positivity in >20% of nuclei defines a

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

group of patients with highly aggressive disease, all six of the patients that died during the follow up period of the study (4.9 to 123 months, mean 55.5months) fell in to this group³¹².

A recent study investigated the usefulness of p₅₃ in locally advanced prostate cancer²²⁰. This study group consisted of 72 patients with a median follow up of 5.4 years (range .5 to 6.4). Biochemical failure, local failure, or systemic failure occurred in 29 of the patients. The group compared the usefulness of p₅₃ positive staining at the 10% level with Gleason score, PSA, pathological stage, volume of extraprostatic tumour extension, and total volume of tumour. Using a multivariate analysis, which contained p₅₃, only p₅₃ predicted outcome, and when it was added to the statistical model for predicting outcome it improved the predictive ability of the model²²⁰.

Other studies have not shown p₅₃ to be so useful. A study evaluating the role of p₅₃, Ki-67, and apoptotic index in predicting recurrence in a series of 47 patients demonstrated that 16 of the specimens had clustered immunopositivity for p₅₃. 17 of the 47 patients had evidence of disease recurrence at 5 years and the conclusion of these authors was that although both p₅₃ and apoptotic index may predict recurrence using a multivariate analysis only the apoptotic index reaches significance in the final logistic regression³⁶⁸. A further study of 208 consecutive radical prostatectomy specimens, 75 with localised intracapsular

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

disease, 123 extra capsular locally invasive, and 8 of these had lymph node involvement. P₅₃ at the >5% level was detected in 79 specimens, and although this did correlate with tumour volume, p₅₃ expression levels were not associated with differences in recurrence free survival¹⁹⁴. A study of 134 patients also showed that p₅₃ was not helpful; this report did not indicate the method of treatment³⁹⁷.

3.32 P₅₃ and watchful waiting

The Scandinavian countries have adopted watchful waiting as their mainstay for the management of asymptomatic carcinoma of the prostate. The rationale for this has been discussed previously (see section 1.66). In a study of 186 patients diagnosed by TURP and treated with watchful waiting 163 patients died, 79 from prostate cancer, during the follow up. Nineteen patients stained for aberrant p₅₃, the mean survival for p₅₃ positive patients was 52 months and for p₅₃ negative patients it was 123 months. There was a significant correlation between tumor stage, tumor grade, metastasis, age and p₅₃ staining. However in a Cox multiple regression analysis of tumor stage, tumor grade, metastasis, age and p₅₃, p₅₃ lost its significance. The authors concluded that the prognostic value of p₅₃ appears to depend upon its association with advanced stage and high grade, and they did not feel it would be a good prognostic marker in early prostate cancer³⁶⁹.

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

A further study from Scandinavia of 221 men diagnosed with carcinoma at TURP and followed up for a median of 15 years showed that all tumors expressed some degree of staining for p₅₃. There was a significant association between p₅₃ staining, local tumor stage and grade. The patients that were highly immunoreactive for p₅₃ did significantly worse than the patients with low immunoreactivity for p₅₃⁴³.

3.33 P₅₃ and radiotherapy.

Radiation causes cellular DNA damage which, if irreparable can lead to apoptosis in a cell expressing wild type p₅₃. Cells with an aberrant p₅₃ can not enter cell cycle arrest at the G₁/S or G₂/M checkpoints, as would cells expressing the wild type p₅₃ gene. In a series of 33 patients who had failed radiotherapy and had pre and post radiotherapy specimens no significant difference in p₅₃ immunostaining was found between the failures (67%) and the successes (75%) of radiation therapy¹⁷³. However, the sample size was small and the follow up period was short at 3 years. A further study of 38 patients also failed to show any relationship between p₅₃ over expression and recurrence³¹⁵.

A large study was commissioned under the auspices of the Radiation Therapy Oncology Group. The protocol, 8610, consisted of cytoreduction pre and peri radiotherapy for locally advanced prostate cancer. One hundred and twenty

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

nine patients were analysed in this sub set analysis with a median follow up of 5 years (range 1.17 – 7.14 years). 23 patients stained for aberrant p₅₃ with > 20% positive nuclei, 12 in the radiotherapy alone arm and 11 in the radiotherapy plus cytoreduction group. Using Cox regression analysis abnormal p₅₃ expression was an independent prognostic indicator for significantly shortened survival, increased incidence of distant metastases, and a reduced progression free survival¹³⁹.

A study of 55 patients with recurrent disease despite radiotherapy, who then underwent salvage radical prostatectomy, p₅₃ staining was detected in 50 specimens⁷⁹. Over expression of p₅₃ was associated with an increased proliferative activity as measured by Ki-67 labelling index. It is not clear from this study whether the over expression of p₅₃ developed overtime or was always present. Additionally radiotherapy can lead to stabilisation of the wild type p₅₃ and so this may be detected by immunohistochemistry. An older study of 18 radiotherapy failures patients also showed a high level of p₅₃ accumulation (72% - 13 patients)³¹⁰. In this study group the authors had pre radiotherapy tissue specimens of 10 patients and in these specimens there was also a high level of p₅₃ expression, implying longstanding p₅₃ alteration. The authors also tested the validity of p₅₃ immunoreactivity with PAb 1801 in ten post irradiation patients, five with positive staining. Using single strand conformational polymorphism they could only demonstrate abnormalities

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

between exons 5 to 8 in three out of the five p₅₃ positive staining patients. None of the negative staining specimens demonstrated any abnormalities. The authors attributed this discrepancy to less than 100% sensitivity of single strand conformational polymorphism, false positive staining with immunohistochemistry, or the p₅₃ mutations may lie outside exons 5 to 8³¹⁰.

The alteration in the phenotype and the genotype of recurrent tumours after definitive therapy was investigated in 32 patients¹⁴². 18 patients had recurrence after radical prostatectomy, and 13 had recurrence after radiotherapy. In both groups of patients there was a significant increase in the Ki-67 labelling index, and p₅₃ nuclear reactivity in tissue obtained from recurrences when compared with primary lesion. The nuclear reactivity was of the clustered type, which may represent a more aggressive phenotype⁴²⁵. There was also an increase in the Gleason scores of the recurrent lesions in the radiotherapy group. This would suggest that a more aggressive phenotype. The authors suggested a number of explanations for this; tumour cells are genetically unstable and therefore the first cells to relapse are the ones with a selective growth advantage. An alternative explanation may be by removing the primary tumour, a growth inhibitor is removed. There is therefore growth acceleration of the residual tissue as is the case in the Lewis lung carcinoma model²⁸⁹.

3.34 P₅₃ and pre-treatment biopsies.

The ability to detect preoperatively those patients that would benefit from radical surgery would be a considerable aid to the surgeons. In a series of 76 patients who had undergone radical prostatectomy and had preoperative TRUS guided biopsies it was shown that the risk of relapse was 24% higher if aberrant p₅₃ was detected on the pre-operative biopsies. Using a multivariate analysis the presence of aberrant p₅₃ and Gleason score > 7 was shown to be independent predictors of relapse in patients undergoing radical prostatectomy⁵⁵.

A study comparing biopsies with 106 radical prostatectomy specimens that had neoadjuvant androgen ablation prior to definitive surgery did not show a correlation between preoperative expression of p₅₃ and outcome⁷⁴. In this study, with respect to radical prostatectomy specimens, the aberrant p₅₃ was detected in 32 cases prior to antigen retrieval and in 77 cases after antigen retrieval. Only 77 biopsies were available for analysis and after antigen retrieval only eight were positive. The expression of aberrant p₅₃ in radical prostatectomy specimens correlated with poor outcome, however in a multivariate analysis this was not an independent prognostic factor. Biopsy expression of p₅₃ did not relate to outcome⁷⁴. The increase in the levels of p₅₃ may be explained by the effect of androgen ablation on the expression of p₅₃.

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

A retrospective review of the pre-treatment biopsies of 49 patients who underwent definitive radiotherapy for the treatment of localised carcinoma was carried out. A surrogate end point for relapse was used; this was a PSA nadir of 1.0ng/ml. The biopsies of 17 patients stained positively for p₅₃, of these patients 15 (88.2%) did not reach a PSA nadir of 1.0. 32 patients stained negatively for p₅₃, of these 23 reached the surrogate end point. Logistic regression analysis showed that p₅₃ was an independent predictor of relapse after definitive radiotherapy³⁴¹. The results of this study could be interpreted as showing that increased p₅₃ is common in patients with advanced possibly extraprostatic disease not amenable to radiotherapy.

A comparison of pre-treatment biopsies and survival was carried out on 60 patients³⁷¹. This was a retrospective study, with the majority of patients not receiving any pre-treatment hormonal manipulation. The group of patients consisted of 36 patients with T₂ or below disease and 18 patients with stage T3-T4 disease. Six patients were Tx, i.e. not staged; the local staging was carried out with digital rectal examination. 15 patients underwent lymph node dissection, and five of these were positive. 59 patients had had a bone scan which was negative. P₅₃ immunostaining was carried out with an antigen retrieval system, overall positive staining occurred in 15 tumours, surprisingly and in contrast to other studies a greater proportion of lower stage tumours stained positively. The survival of the p₅₃ positive group appeared to be better

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

than the p₅₃ negative group. The authors concluded that no survival disadvantage was seen in patients with p₅₃ immunoreactive tumours, implying aberrant p₅₃ does not confer radio resistance and that alternative pathways to apoptosis may be more important. This study is incomplete in many ways. There is no mention of PSA, and no imaging of the lymph nodes was carried. It is not incomprehensible to suspect that many of these patients may have advanced disease since in the 15 patients suitable for radical surgery a third had disease beyond the prostate and mean survival of the patients was only 75 months (range 8.5 – 172).³⁷¹.

3.35 P₅₃ and metastatic prostate cancer

A study of ten lymph node specimens with prostate cancer showed that only one stained positively for p₅₃⁴⁰⁶. A further study of 56 patients with metastatic lymph node disease and a median follow up of 3.8 years was carried out²³⁸. In this study 26 patients had material available from both the primary tumour and lymph nodes, a further 15 had material available from the primary tumours only, and 15 patients had only lymph node samples. P₅₃ was detected in 5 primary tumours, and in 7 lymph node specimens. The median Ki-67 labelling index was 8.4% in the primary tumours and 8.7% in the lymph nodes. The median Ki-67 labelling index was significantly higher in the p₅₃ positive patients. Using a multivariate analysis the Ki-67 labelling index and Gleason scores of the primary tumour were independent prognostic factors, in patients

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

with positively staining lymph nodes the presence of bone metastasis was also a prognostic factor²³⁸. Although patients staining positively for p₅₃ and bcl₂ tended to have a poorer outcome this did not reach significance²³⁸. A study of 36 primary prostate cancers, 17 patients with nodal involvement, and 14 cases of bone metastasis, demonstrated a greater proportion of the specimens obtained from metastasis showed aberrant staining for p₅₃; i.e. only ten of the primary tumours, ten of the lymph nodes with metastasis, and six of the bone metastasis¹⁷⁴. Interestingly there was no correlation between p₅₃ expression in the primary tumour and metastasis.

The difference in p₅₃ staining between treatment naïve, metastatic and hormone refractory prostate cancer was investigated in 61 patients¹⁶¹. The 61 patients consisted of 27 patients with recurrence after either radiotherapy or hormonal intervention or both and tissue was obtained by TURP and in one case by prostate biopsy; 8 patients with metastatic lymph node disease and tissue was obtained from the lymph nodes; and 27 primary untreated tumours and tissue was prostate biopsies. The authors demonstrated an increase in p₅₃ in the hormone refractory group i.e. 20 out of the 26 cases (76%) stained positively for p₅₃ compared to 4 out of the 8 (50%) metastatic, and 6 out of 27 (22%) in the untreated group. The authors concluded that this study demonstrated an increase in p₅₃ from untreated primary tumour to hormone refractory disease¹⁶¹. However there are several points in this paper, which

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

are not that clear. The group of patients with refractory disease was rather heterogeneous, and it is not clear whether all the patients in the relapse group actually relapsed using PSA criteria or needed a TURP for non-malignant outlet obstruction. The abstract of the paper claims that 94% of the patients in this group were p₅₃ positive, where as in the body of the paper only 76% i.e. 20 out of 26 stain positively for p₅₃. In addition it is worth noting that five patients in the relapsed group had cytoplasmic staining for p₅₃, the authors graded these as positive¹⁶¹.

A study of bone marrow biopsies obtained from the iliac crests of 43 patients with hormone refractory metastatic carcinoma of the prostate showed that 22 (51%) exhibited immunohistochemical positivity for p₅₃²⁴⁵. Eight cases exhibited positive staining for bcl₂ and p₅₃, and six cases were positive for just bcl₂. The expression of either of these proteins did not influence outcome and so it was concluded that these genetic mutations are not essential for the fatal progression of prostate cancer. There must be multiple pathways of tumour progression in prostate cancer²⁴⁵.

3.36 The effect of castration on p₅₃.

In a retrospective study, Stattin et al studied the effect of castration on 28 patients with carcinoma of the prostate³⁷². The 28 patients consisted of 15 responders and 13 non-responders, 3 patients were initially stained for p₅₃ and

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

further 3 responders stained after LHRH therapy. P₅₃ immunoreactivity did not correlate with outcome after castration³⁷².

Increased p₅₃ was also detected in-patients who had had neoadjuvant LHRH therapy prior to radical prostatectomy. All the patients in this cohort who had expression of p₅₃ relapsed. The authors concluded that the finding of an aberrant p₅₃ could be due to the advanced stage at which the patients presented and were selected for treatment using neoadjuvant LHRH, since an altered p₅₃ may lead to resistance to castration induced apoptosis and so disease relapse. Therefore the presence of an aberrant p₅₃ may be an early event in the evolution of hormone refractory disease in clinically localised prostate cancer²⁹⁵. This relationship between aberrant p₅₃ after neoadjuvant therapy and relapse was further commented upon, when 19 out of 21 patients with p₅₃ clustering relapsed within 36 months of definitive treatment³¹².

This relationship between the presence of aberrant p₅₃ detected immunohistochemically after LHRH treatment was further strengthened in a subset analysis of 263 radical prostatectomies carried out in New South Wales in Australia³¹². 39 patients in this cohort of 263 had received neoadjuvant LHRH treatment prior to definitive surgery. 23 patients in this group of 39 had relapsed. A total of 21 patients were p₅₃ cluster positive, and 90% (19 patients) of the cluster positive patients relapsed. P₅₃ cluster status was the

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

strongest predictor of recurrence in a multivariate analysis comparing p₅₃, pretreatment PSA, Gleason score, and pathological stage³¹².

In a study of 26 patients with prostate cancer that had relapsed despite first line therapy – radiotherapy or hormonal or both, a total of 20 patients demonstrated p₅₃ staining and in 4 of these the staining was cytoplasmic¹⁶¹. The significance of cytoplasmic staining of a nuclear protein is at present unclear. However the authors counted these tumors as p₅₃ positive.

The relationship between aberrant p₅₃ expression and androgen receptor gene amplification was studied in 30 patients with hormone refractory disease. Comparing primary untreated tumours with hormone refractory specimens obtained from the same patient, the immunostaining for p₅₃ increased from 17% to 40% respectively. The tumours with aberrant p₅₃ expression were more likely to have androgen receptor gene amplification, suggesting that tumours with an aberration in p₅₃ were more likely to become hormone refractory and so survive in lower androgen concentrations²⁰¹.

3.37 Site of p₅₃ mutations

Micro dissected DNA selectively extracted from paraffin-embedded sections of prostate cancer and PIN has shown a high frequency of p₅₃ mutations in the lesions dissected from the peripheral zone lesions when compared to those that

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

were dissected from the transition zones. For the cancer lesions the incidence was 33.3% in the peripheral zone versus 4% in the transition zone, and for PIN the incidence was 14.7% and 5.6% respectively. In areas of benign tissue adjacent to the PIN or cancer there were mutations detected in p₅₃. The point mutations detected in PIN and cancer were not identical the authors attributed this to the polyclonality of prostate cancer. In addition the authors found a higher incidence of p₅₃ mutation in T₃ carcinomas of the prostate when compared to T₂ lesions. The authors conclude by stating that their findings indicate that the p₅₃ gene mutations are involved in prostatic carcinogenesis and explain why the non-transition zone is the predominant site of cancer³⁷⁷.

3.38 Racial differences

A heterogeneous group of 62 archival samples of prostate tumor obtained from Asians of non-Japanese origin were examined⁸¹. The specimens were obtained by TURP and the patients were staged with a CT scan of the abdomen and pelvis, only 2 patients' expressed p₅₃, both of these patients had metastatic disease. The authors concluded that the expression of p₅₃ mutation in Asian cohorts is significantly lower.

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

Table 6 (cont.): The incidence of p53 in patients with prostate cancer.

²⁷⁸	1994	Rad Pros tumour (28), Rad Pros PIN (16) metastatic primary (16), lymph nodes (18).	78	28	?	16+18	IHC	2/28 localised carcinoma p53+, 1/16 PIN p53+, 9/16 metastatic p53+, 10/18 lymph nodes p53 +.	Increasing p53 in pts with mets.
³²	1994	Microdissection of benign lesions close to malignant tissue	21	6	12	3	PCR SSCP	1/21 benign lesion close to Gleason 7 lesion showed alteration in p53.	P53 alteration may be an early change in the genesis of prostate cancer.
¹⁶¹	1995	TURP + Biopsies	61	20	26 +27	8	IHC PCR	6/27 treatment naïve p53+, 20/26 androgen refractory p53+, 4/8 metastatic were p53+.	Progression in p53 alteration from untreated to hormone refractory disease.
¹⁵¹	1995	TURP + biopsies.	37	26	11	0	IHC PCR	1/37 p53 + P53 positive on IHC confirmed by PCR. 7 negative patients had PCR, no p53 mutation.	IHC is reliable, and correlates well with PCR. The authors concluded p53 mutations detected by IHC are a late step in the progression of prostate cancer and are associated with advanced disease, dedifferentiation, and the acquisition of androgen independence.
³¹⁰	1996	Salvage prostatectomy (15) or TURP (3) for relapse post DXT.	18	?	?	?	IHC + PCR	13/18 (72%) + in relapse group, 20 % + in control (Non irradiated advanced Ca P)	P 53 aberration common in patients with relapse post DXT.

Table 6: The incidence of p53 in patients with prostate cancer.

Ref.	Year of pub	Type of specimen	No of cases	No with intracapsular disease	No with extra capsular disease.	No with metastasis	Method	Results of P53 staining	Conclusion
⁴⁰⁶	1994	85 prostate (Rad pros + biopsies)+ 10 lymph nodes	85 + 10	?	?	10	IHC + Polymerase chain reaction – single stranded polymorphism (PCR-SSCP)	3/85 p53 + in primary prostate cancer and 1/10 p53 + in lymph nodes.	P53 mutations appear to be of low frequency in prostate cancer.
¹⁶²	1994	Rad pros	73	28	45		IHC	4/28 localised p53 + 31/45 extracapsular p53 +	P53 correlated with more advanced tumour stages, with higher Gleason grades, and with larger total and high-grade tumour volumes.
¹¹	1994	Radpros, TURP, Primary, lymph nodes, Mets	93 (48 prostates 29 lymph nodes, 16 bone mets).	9	18	11 rad pros in pts with lymph node involvement 10 TURP 29 lymph node 16 Bone mets	IHC	0/9 p53+ in localised 1/18 p53+ in extra capsular 2/11 p53+ in pts with lymph node+ 6/10 p53+ in hormone refrac CaP 2/29 p53+ in lymph node met 8/16p53+ in bone mets.	P53 correlates with high grade, hormone refractory tumours, and bone mets.

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

Table 6 (cont.): The incidence of p53 in patients with prostate cancer.

⁴²³	1997	Rad TURP. Pros. Biopsy	36	22	?	14	IHC	5/22 (23%) p53 positive localised. 1/14 (7%) metastatic positive.	Follow up only 12 months. no relationship between outcome and p53 status
²⁷³	1997	Radical prostatectomy	100	67	33	29	IHC PCR-SSCP	1/67 localised p53 + 5/33 extra capsular p53 + 5/6 mutations detected by IHC 6/6 detected by PCR - SSCP.	P53 mutations are a rare late event in prostate cancer.
⁶⁶	1997	Biopsy TURP	40	15	24	21		T ₁₋₂ =5/15 p53 + T ₃₋₄ =14/24 p53 + 17/19 p53+ pts progressed 12/21 p53 - pts progressed	P53 positive tumours higher grade and quicker progression. No correlation between p53 and p21 or waf1.
³⁹⁷	1998	Not Stated	134	51	27	56	IHC	11/51 p53+ Localised 7/27 p53+ Extra capsular 15/56 p53+ With extra prostatic disease.	P53 did not predict outcome (Mode of treatment not stated)
⁷⁹	1999	Salvage Radical prostatectomy in post-DXT relapse.	55	?	?	?	IHC	50/55 P53 +	High incidence of p53+ in patients who relapsed post DXT.
⁸¹	2000	62 TURP	62	13	4	36	IHC PCR	2 p53+	Lower p53 expression in Asians
³⁵⁶	2004	Rad Prost	98	?	?	?	IHC	55/98 Positive by IHC	No prognostic data, but compared IHC with PCR. IHC generated 27/55 false positive. and 15/42 false negative.

Table 7: The relationship between p53 watchful waiting

Ref.	Year of pub	Type of specimen	No of cases	No with intracapsular disease	No with extra capsular disease.	No with metastasis	Method	Results of P53 staining	Conclusion
³⁶⁹	1996	TURP	186	159	27	23	IHC	T0 = 6% M0 = 9% T1 = 13% M1 = 22% T2 = 10% Mx = 14% T3 = 25% T4 = 38%	P53 correlates with stage, grade, metastasis, but in multiple regression analysis it is not an independent predictor.
⁴³	2000	TURP	221	131	90	60	IHC	P53 staining correlated with stage.	P53 was an independent adverse prognostic indicator.

Table 8: The relationship between p53 and outcome in patients who have had radical prostatectomy.

Ref.	Year of pub	Type of specimen	No of cases	No localised	No locally invasive	No lymph nodes/mets	Method	Results	Conclusion.
⁴²	1993	Rad Pros +TURP	150	38	36	69	IHC + PCR	Stage 0 1/12 p53 + Stage 1 0/26 p53 + Stage 2 2/36 p53 + Stage 3 8/36 p53 + Stage 4 8/33 p53 + Unstaged 0/7 p53 + {Treatment was radical prostatectomy (88), TURP (52), hormonal manipulation (10)}	19/150 p53 + P53 + related to stage.
¹⁹	1995	Rad Pros	139	57	81	1	IHC	28/57 localised p53 + 56/81 extracapsular p 53 +	P53 was an independent marker for progression.
⁴²⁵	1996	Rad Prost	49	29	20	?		10/16 with recurrence, compared to 7/33 in non-recurrent group.	P53 staining associated with recurrence.
²⁷⁴	1996	Rad Pros	162 mean follow up 4.5yrs	62	99	?	IHC	38/62 localised p53 + 73/99 extraapsular p53 +	In a multivariate analysis p53 and Bcl-2 independent prognostic markers. P53 + = 112 (69.1%), bcl-2 = 44 (27.2), Ki-67 = 62 (38.3%)
²⁰	1996	Rad Pros	175pts with mean fu 4.6yrs	174	1	?	IHC	113/174 localised p53 + 1/1 extracapsular p 53 +	In a multivariate analysis p53 and Bcl-2 independent prognostic markers. Bcl2+ = 26.9%, p53+ = 65%, multivariate analysis p53 and bcl2 independent predictors of disease.

Table 8 (cont.): The relationship between p53 and outcome in patients who have had radical prostatectomy.

⁶⁰	1996	Rad Pros	137	44	93	34	IHC	1/44 localised p53 + 10/93 extra capsular p53 + 3/34 node + pts were p53 + in the primary tumour.	P53 + tended to worse prognosis but not significance. Bcl2 pT3 (31%) > pT2 (5%) significant. Combined analysis of Ki67, LI, and Bcl-2 allowed the distinction of three groups with different clinical outcome. Prognosis was best in Bcl-2- negative tumours with low Ki67 LI, worst in Bcl-2- positive tumours with high Ki67 LI.
³⁷⁴	1996	Rad Pros	40	0	40	?	IHC	32/40 p53 +	20/32 p53 + progressed. P53 was an independent marker for progression.
²⁰⁹	1998	Rad Pros	76	42	34	6	IHC	18 >20% p53 +	>20% p53 + only variable associated with decreased long- term survival in multivariate analysis.
¹⁹⁴	1998	Rad Pros	208	75	123	8	IHC	79 p53 +	P53 and Rb did not predict recurrence, but p53 did correlate with tumour volume. MIB-1 and Bcl-2 correlated with progression.
¹⁴²	1998	Rad Pros	18	11	7	5	IHC	1/18 Primary p53 + 7/18 Recurrent p53+	Increased p53 in recurrences, genotype of recurrence is more aggressive.
³⁶⁸	1998	Rad prost	47	47	0	0	IHC	10/17 p53+ in relapse group @ 5yrs 6/30 p53+ in non-relapse group @ 5 yr.	In multivariate analysis only apoptotic index and clustered p53 added info. in regression model only AI reached stat sig.

Table 8 (cont.): The relationship between p₅₃ and outcome in patients who have had radical prostatectomy.

⁵⁵	1999	Rad Prost + pre-op biopsies.	76	37	39	?	IHC	39 p53+	P53 predicted recurrence. Aberrant p53 led to a 24% higher risk of relapse. P53 expression was more frequently observed in biopsy specimens? difference in fixation? difference in tumour due to multifocality.
²⁹⁵	1999	Rad Prost	86	51	35	?	IHC	6/86 >20% p53 +	P53 +ve relapsed quicker. No correlation between p53 + and preop PSA, tumour stage, or grade. Also stain for mdm2 and p21 which was positive in 50% of cases and postulated that p21 activation due to an alternative mechanism other than p53 activation.
³¹²	2000	Rad Pros	263	114	144	5	IHC	132 (50.2%) (nuclear clustering) >0 to <2% 33 (12.2%) >2 to <5% 35 (13.3%) >20% 34 (12.9%)	P53+ve (Clustering or nuclear accumulation) was an independent predictors of relapse, as was pre treatment PSA, tumour stage, tumour grade, and lymph node status. In pts treated with preop LHRH p53 clustering predicted relapse.
²²⁰	2000	Rad Pros in locally advanced disease.	72	0	70	23	IHC	28 >10% p53 +	P53+ predicted biochemical, local, & systemic failure.

Table 8 (cont.): The relationship between p₅₃ and outcome in patients who have had radical prostatectomy.

⁷⁴	2000	Rad Prost Biopsies after neoadjuvant treatment.	106	57	31	17	IHC	77/106 p53 + 8/77 biopsies p53 +	P53+ increased after hormonal Tx. and correlated with grade, stage, biochemical progression and short survival. P53+ from pre LHRH biopsies did not show any correlation.
³⁵⁶	2004	Rad Prost	98	?	?	?	IHC	55/98 Positive by IHC	No prognostic data, but compared IHC with PCR. IHC generated 27/55 false positive, and 15/42 false negative.
¹⁷⁷	2005	Rad Pros	52	14	38	0	IHC	20 p53 +	13 p53+ relapsed, 7 p53- relapsed (P<0.0001). p53+

Table 9: The relationship between p53 and outcome in patients who have had radiotherapy.

Ref.	Year of pub	Type of specimen	No of cases	No localised	No locally invasive	No with lymph nodes/mets	Method	Results	Conclusion.
³⁷¹	1996	Pre-DXT Biopsies.	60	36	18	5	IHC	Localised = 12/36 p53+. Locally invasive = 3/18 p53+	P53 correlated with stage, not grade. P53 did not predict relapse.
¹³⁹	1997	Pre-DXT Biopsies	129		129		IHC	23/129 (18%) with p53+ >20%. 6/23 p53+ progressed 27/106 p53- progressed 12/23 p53+ died at 4 yr. 43/106 p53- died at 4 yr.	P53 + prognostic indicator for survival, decreased progression free survival, decreased overall survival and distant metastasis. Independent of Gleason score and stage.
¹⁴²	1998	Pre-DXT biopsies & post-DXT biopsies.	13			13	IHC	Primary p53 + = 1/13 (8%) Recurrent p53 + = 7/13 (54%)	Genotype of recurrence is more aggressive than primary lesion.
¹⁷³	1998	Pre-DXT biopsies & post-DXT biopsies.	33 pt. who relapsed after DXT	?	?	?	IHC	P53 + pre DXT = 24/33 P53 + post DXT = 20/33	P53 did not increase post DXT, and could not predict DXT failures.
³⁴¹	1999	Biopsies	49	49			IHC	17/49 p53+ 32/49 p53-	15/17 p53+ did not reach PSA nadir of 1.0 23/32 p53- did reach PSA nadir of 1.0. Pre DXT p53 status predicts response to DXT.

Table 10: P₅₃ in patients treated with hormonal manipulation.

Ref.	Year of pub	Type of specimen	No of cases	No localised	No locally invasive	No mets	Method	P53	
⁴⁰⁵	1992	TURP = 44 Biopsy= 75 Open Spec = 18	137	51	82	36	IHC	T1 + T2 = 2% M0 = 2% T3 = 2% M1 = 11% T4 = 12%	Pts treated with hormonal manipulation (n=97), DXT, and surgery. High level staining correlates with early relapse, poor survival, and high grade.
³⁷²	1998	Biopsies before & after LHRH	28	7	21	0	IHC	3 (6 after LHRH) p53 +ve.	P53+ve could not predict response to LHRH treatment, but increased in responders.
²⁴¹	2000	Biopsies pre hormonal manipulation.	73	0	0	73	IHC	P53 + = 14/73 (19.2%). 6/14 P53+ progress 8/59 p53- progress	P53 positive patients correlated with early relapse.

Table11: The relationship between p₅₃ and outcome in patients with extra prostatic disease.

Ref.	Year of pub	Type of specimen	No of cases	No localised	No locally invasive	No with metastasis	Method	P53	Conclusion
²⁴⁵	1998	43 patients with hormone refractory disease.				43	IHC	22/43 (51%) p53+ in bone marrow.	Did not correlate with outcome
²³⁸	1998	Lymph nodes + Rad Prost	56	39	17	56 nodes 19Bone metastasis.	IHC	5/41 Primary tumours p53+ 7/41 lymph nodes p53+	P53+ had no prognostic significance.
⁷⁸	1999	Rad Pros + Lymph nodes	220	0	0	220	IHC	109 Rad Pros p53+ 83 Lymph nodes p53+ The majority of patients then had LHRH.	P53 expression in the primary tumour was concordant with p53 presence in nodes. The presence of p53 expression in the nodes was an adverse prognostic factor for survival.

Table 12: The relationship between p53 in pre op biopsies and outcome.

Ref.	Year	Specimen	Number	Method	Stage	Results	Conclusion
²³⁰	1997	Pre treatment biopsies Mean follow up 35.5 (range 1 –132 months). Treatment with Rad pros (49), hormonal therapy (89), or surveillance (8).	146	IHC	T ₁₋₂ N ₀ M ₀ =50, T ₃₋₄ N ₀ M ₀ =28, T ₁₋₄ N ₁ M ₀ =9, T ₁₋₄ N ₁ M ₁ =59.	T ₁₋₂ N ₀ M ₀ p53 + = 10/50. T ₃₋₄ N ₀ M ₀ p53 + =8/28. T ₁₋₄ N ₁ M ₀ + T ₁₋₄ N ₁ M ₁ p53 + = 22/68. P53+ = 10% staining +	P53 positive correlates with shorter progression free survival. Bcl2 positivity an independent indicator of poor prognosis.
⁶³	1998	Pre-tx biopsies. Follow up 5 years, heterogeneous treatment with Rad pros, DXT, hormonal therapy, TURP, or surveillance.	111	IHC	?	16% p53 positive. 20% bcl2 positive. 7.5% Ki67 positive.	P53 + correlated with high-grade tumour. Univariate analysis showed that high Gleason score, high percentage of tumour, high Ki-67 LI, and p53 positive purports high percentage of death. But multivariate high Ki-67 only independent predictor.
⁵⁵	1999	Pre treatment biopsies and radical prostatectomy specimens. Mean follow up38 months (range 6 to 114 months). Treatment with radical prostatectomy.	76	IHC	T ₁₋₂ N ₀ M ₀ =37, T ₃₋₄ N ₀ M ₀ =39.	19/22 with relapse are p53+ on biopsy. 20/40 relapse free are p53+ on biopsy.	Biopsy p53 + in multivariate analysis correlated with poor outcome. Aberrant p53 observed more often in biopsies than in radical prostatectomy specimens.
³⁶⁴	1999	Comparison of needle biopsies (129) with rad prostatectomy(199) specimens	129	IHC	?	Rad Pros 132/193 p53+ Needle biopsy 64/129 p53+	Immunostaining of biopsies not predictive of recurrence. Poor correlation between biopsy and rad prostatectomy specimens, but as the editorial suggests the biopsies were done at many institutions and this may be a technical problem.

Table 13: The relationship between aberrant p₅₃ and prostatic intra epithelial neoplasia.

Ref.	Year	Specimen	Number	Method	Stage	Results	Conclusion
¹⁷⁵	1995	Rad Pros	40	IHC	T ₁₋₂ N ₀ M ₀ =40	9/40 PIN p53+ 13/40 CaP p53+	Ca P & PIN from 7 common patients demonstrated intense p53 staining.
⁴²⁶	1998	Rad Pros (27) TURP (2)	29	PCR-SSCP		3/29 PIN p53 + 5/29 CaP p53 +	2 patients had mutations in both Ca P & PIN, but these were distinct mutations.
¹⁸⁷	1998	Rad Pros (9 pts with neoadjuvant LHRH)	44	IHC	T ₁₋₂ N ₀ M ₀ =26. T ₃ N ₀ M ₀ =18.	0/44 PIN p53 + 2/43 CaP p53 +	Some pts with LHRH, low incidence of p53.
³⁷⁷	2000	Rad Pros (14 pts with neoadjuvant LHRH)	27	PCR-SSCP	T ₁₋₂ N ₀ M ₀ =11. T ₃ N ₀ M ₀ =16.	6/27 PIN had p53 mutations 8/27 Ca had p53 mutations. In only 2 cases did both Ca and PIN have mutations. Both of these mutations were different.	Increase in p53 mutation with stage. Different mutations in p53 in PIN and Ca imply PIN arises from a different clone.

Table 14: The relationship between aberrant p₅₃ and metastatic prostate cancer.

Ref.	Year	Specimen	Number	Method	Stage	Results	Conclusion
³⁸⁷	1993	Primary tumours	68	IHC Flow Cytometry	?	9/68 p53+	P53 + tumours were higher stage & Gleason score, + progressed more rapidly.
¹⁰³	1994	TURP (20) + Lymph node(15)	35	IHC PCR-SSCP	?	2/20 TURP p53 + 2/15 Lymph node p53+	Total concordance between PCR-SSCP and IHC. P53 mutation is a late and infrequent event.
²⁹	1995	50 hormone sensitive + 34 hormone resistant	84	IHC PCR-SSCP	?	6/34 hormone res p53+ 19/50 hormone sens p53+	A significantly higher proportion of tumours with mutations were poorly differentiated compared with tumours without mutation (P < 0.04). p53 mutation is a late event in the progression of the disease
¹¹⁰	1995	Rad Pros (39). Rad Pros/TURP + mets/nodes (47)	86	IHC PCR-SSCP	T ₁₋₂ N ₀ M ₀ = 18 T ₃₋₄ N ₀ M ₀ = 21 T ₁₋₄ N ₁ M ₁ = 47	0/18 T ₁₋₂ N ₀ M ₀ p53+ 2/21 T ₃₋₄ N ₀ M ₀ p53+ 4/47 T ₁₋₄ N ₁ M ₁ p53+. & 11/47 mets/nodes p53+.	4/47 pts with mets had abnormal both in prostate and lymph nodes. 7/47 had mutation at mets only. P53 mutation associated with advanced stage.
¹⁷⁴	1995	36 primary cases. 17 positive lymph nodes. 14 bone mets	68	IHC	?	10/36 primary tumours positive 9/36 were Gleason score>7. 10/17 lymph nodes positive. 6/14 bone mets positive.	Metastatic tumours express more aberrant p53 than primary tumours.

Table 14 (cont): The relationship between aberrant p53 and metastatic prostate cancer.

²⁵⁰	1995	Rad Pros (29), frozen section of primary tumour (9), lymph node (12), and bone marrow (1).	51	IHC, PCR-SSCP	?	6/38 primary tumours had aberrant p53. 3/13 mets/nodes had aberrant p53.	Metastatic tumours display higher p53 than primary.
¹⁰	1996	Biopsy, TURP, Rad Pros. Treatment heterogeneous.	77	IHC	T ₁₋₂ N ₀ M ₀ =18, T ₃ N ₀ M ₀ =14, T ₁₋₄ N ₁ M ₁ =45	23/77 + for p53, 14 of the 23p53+ patients were refractory to hormonal manipulation. 37/77 + for bcl2.	23/77 + for p53, 37/77 + for bcl2. P53 staining corresponded to hormone resistance. In 14/21 (66%) of cases (p=.0012), increased p53 +bcl2 corresponded to hormone resistance in 13/16 (81.2%) of cases (p<.0001).
²⁴⁵	1997	Bone Marrow biopsy in pts treated with LHRH.	43	IHC	T ₁₋₄ N ₀₋₁ M ₁ =43	22/43 P53+	P53 expression did not correlate with tumour burden or outcome. Bcl2 expression was inversely correlated with p53 expression.
¹⁴²	1998	Rad pros + prostate fossa biopsy (18).	18	IHC	T ₁₋₂ N ₀₋₁ M ₀₋₁ =11, T ₃ N ₀₋₁ M ₀₋₁ =6, T ₄ N ₀₋₁ M ₀₋₁ =1	Primary p53 + = 1/18 Recurrent p53 + = 7/18	Genotype of recurrence is more aggressive than primary lesion.
²⁵⁴	1998	Bone Marrow biopsy in pts treated with LHRH.	17	IHC RT-PCR+SSCP	T ₁₋₄ N ₀₋₁ M ₁ =17	10/17 P53+ staining 8/10 abnormal on RT-PCR+SSCP	71%(12) of the bone marrow samples had aberrant p53; RT-PCR detected 2 missed by IHC.

Table 15: Miscellaneous papers relating p53 and prostate cancer.

Ref.	Year	Design of study	Specimen	Number	Method	Stage	Results	Conclusion
²¹⁵	1998	Detect autoantibodies for p53 in sera.	Sera	73	ELISA	T ₁₋₄ N ₀ M ₀ =62, T ₁₋₄ N ₀ M ₁ =11.	2/62, 0/11 + for antibodies.	2 patients with localised disease had antibodies; none of the patients with metastatic disease had antibodies. Both of these patients died in 2-6 months.
²⁴⁰	1998	Relationship between p53 and p21.	Archival specimens	60	IHC	T ₁₋₂ N ₀ M ₀ =16, T ₃₋₄ N ₀ M ₀ =10, T ₁₋₄ N ₀₋₁ M ₁ =34,	T ₁₋₄ N ₀ M ₀ =10/26 p53 +, T ₁₋₄ N ₀₋₁ M ₁ =23/34 p53+	P21waf1 downstream messenger of wttype p53 not activated by mutant p53. p53+/p21-poor prognosis, compared to p53+/p21+
²⁰¹	1999	Hormone refractory tumours (30) versus matched untreated primary (24).	Archival specimens	30	FISH IHC	T ₁₋₂ N ₀ M ₀ =6, T ₃₋₄ N ₀ M ₀ =24, T ₁₋₄ N ₀₋₁ M ₁ =10,	Primary tumours = 4/24 p53+. Hormone refractory = 12/30 p53+	P53 positive tumours show androgen receptor gene amplification, which may allow these tumours to grow in a low androgen milieu.
¹⁹³	2000	Relationship between p53 and 15-lipoxygenase in CaP	Fresh frozen	48	IHC	?	36/48 15-lipoxygenase +. 19/48 p53 +	Correlation between p53, 15-lipoxygenase, and Gleason score. Mutant p53 upregulates 15 lipoxygenase expression.
³⁷⁵	2000	Relationship between p53 and VEGF IN CaP	Archival specimens	55	IHC	T ₁₋₂ N ₀ M ₀ =31, T ₃₋₄ N ₀ M ₀ =24,	12 overexpression of P53	No correlation between p53 and VEGF

3.4 BCL₂

The Bcl₂ family of related proteins constitute important regulators in the control of apoptosis (programmed cell death) ⁷. There are over a dozen members of the bcl₂ family some suppress and some promote apoptosis. The bcl₂ gene is highly conserved. The bcl₂ gene was originally detected in a proportion of B-cell follicular lymphomas at the t (14:18) chromosomal translocation breakpoint ²⁰². The bcl₂ proto-oncogene is located on chromosome 18q21 and codes for a 26 kDa, 239 amino-acid protein with a hydrophobic carboxy terminal. Bcl₂ protein has been located to the mitochondrial membrane ¹⁶³, nuclear envelope, and the endoplasmic reticulum³⁸⁸. Bcl₂ may inhibit apoptosis by regulating the mitochondria permeability and the release of pro apoptotic enzymes. The pro apoptotic members of this family are cytosolic, following the initiation of apoptosis the pro apoptotic members undergo a conformational change and integrate into membranes. An additional characteristic of this family is their frequent ability to form homo as well as heterodimers, suggesting neutralising competition between these proteins.

In IL-3 dependent myeloid and lymphoid cell lines, bcl₂ inhibited apoptosis induced by IL-3 withdrawal without stimulating proliferation ⁴⁰¹. Implying therefore, that the malignant potential of bcl₂ is mediated by its ability to promote cell survival rather than proliferation. In benign prostatic tissue, the

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

basal cells express bcl₂ ¹⁶³.

In prostate cancer, hormone manipulation achieves its effect through activation of apoptosis. Bcl₂ is an inhibitor of apoptosis induced by a number of mechanisms – growth factor withdrawal, glucocorticoid withdrawal, heat shock, and chemotherapeutic agents.

The initial evidence demonstrating that bcl₂ imparts a growth advantage following androgen ablation was obtained using the hormone sensitive LNCaP cell lines, the cell lines were transfected with bcl₂ and androgen withdrawn from the growth medium. The native cell lines did not grow in the androgen free medium; where as the rate of growth of the transfected cell lines was not affected by androgen withdrawal ³¹³. This model was further developed in xenografts of these cell lines. The cells were injected into nude mice. Following castration, the bcl₂ expressing tumours exhibited a significant growth advantage. Therefore, bcl₂ enables prostate cancer cells to remain viable despite androgen ablation ²².

The relationship between androgen ablation and bcl₂ expression was studied in castrated animals at intervals of 1 to 10 days post castration. Messenger RNA for bcl₂ was extracted from the ventral lobe of the prostate. This study demonstrated that castration lead to an increase in the levels of mRNA for

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

bcl₂. The increase was greatest at ten days. This effect could be negated with subcutaneous injections of testosterone²⁴⁶. Therefore androgen ablation itself may lead to the selection of an androgen insensitive phenotype²⁴⁶.

It has been suggested, using gamma irradiated LNCaP cells that bcl₂ protein may inhibit the nuclear import of p₅₃ protein and so prevent apoptosis, the exact mechanism by which bcl₂ inhibits the transfer of cytoplasmic p₅₃ protein into the nucleus is unclear²¹. Structural studies of the bcl₂ protein demonstrate similarities to the pore-forming domains of bacterial toxins, which suggest a role in modulation of membrane function⁸³. This evidence suggests a gate keeper role for bcl₂.

A set point exists in most cells for the ratio of bcl₂ family dimmers, e.g. bcl₂: bax and this determines the sensitivity of a cell towards survival and apoptosis²⁹². Internal and external messengers can further regulate this ratio³³⁵. Bax may be negatively selected during progression of cancer implying that it may function in a tumour suppresser role independent of p₅₃³¹⁷.

Phosphorylation of bcl₂ at specific serine residues can lead to its inactivation and so can inhibit its anti-apoptotic functions¹⁵⁰. P₅₃ can mediate bcl₂ phosphorylation and apoptosis via activation of G proteins Cdc42. Cdc42 is a

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

GTPase, and p₅₃ promotes apoptosis through activation of this G protein³⁸⁶.

Phosphorylation of the bcl₂ protein also leads to decreased binding to bax.

The anti-cancer drug taxol induces bcl₂ phosphorylation and subsequent apoptosis of prostate cancer cell lines¹⁴⁹. Taxol also has an effect upon microtubule, it inhibits microtubule depolymerisation during the cell cycle³²⁹, Haldar et al have used this evidence to suggest that microtubule damage may lead to cell death through a mechanism that involves bcl₂ phosphorylation¹⁴⁸. Taxol therefore induces bcl₂ phosphorylation and cell death, whereas drugs, which damage DNA, do not induce bcl₂ phosphorylation but induce apoptosis through a p₅₃ dependant pathway. This has led Haldar et al to suggest that bcl₂ is the guardian of microtubule integrity¹⁴⁸. Microtubules are important in chromosome segregation and alteration could lead to genomic instability and cell death.

Antisense oligodeoxynucleotides (ODN) are chemically modified stretches of single stranded DNA that are complimentary to mRNA regions of a target gene. ODNs inhibit gene expression by forming RNA/DNA duplexes. Using antisense bcl₂ ODNs and the agent paclitaxel in LNCaP cell lines implanted into nude mice, Leung et al demonstrated that serum PSA levels mirrored tumour volume. Mean serum PSA levels returned to or were above precastration levels by 11 weeks post castration in mice treated with bcl₂

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

ODNs or paclitaxel alone, but remained 90% below pre-castration levels in mice treated with combined antisense ODNs and paclitaxel. Therefore the combination of antisense bcl_2 ODN and paclitaxel led to a decrease in tumour volume and delay in androgen independent progression²²². Similar results were obtained by Miyake et al except they also demonstrated using Shinogi tumour cells that antisense bcl_2 ODNs could be tumour specific and there was no reduction in bcl_2 expression in normal tissue^{132:255:256}.

Transgenic mice have been used to study the pathogenesis of prostate cancer. The most extensively used model is the TRAMP (transgenic adenocarcinoma mouse prostate) model. In this model probasin directed expression of SV40 early genes in the prostate results in progression from hyperplasia to metastatic disease by 28 weeks of age. This model was used to target bcl_2 to the prostate and generate the BxT (bcl_2 x TRAMP) mouse. The BxT mice displayed increased prostate size associated with decreased apoptosis, earlier development of PIN and invasive carcinoma. By 20 weeks 100% of the BxT mice exhibited evidence of prostate cancer compared to only 50% of the TRAMP mice. The rate and incidence of tumour metastases was the similar for the TRAMP and BxT mice. The transgenic bcl_2 did not result in malignant transformation of the prostate gland⁵⁸.

Transfection of bcl_2 into LNCap and PC-3 cell lines followed by exposure to

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

ionising radiation led to the bcl_2 transfected cells having decreased apoptosis although both the controls and the transfected cells did not form colonies after exposure to radiation ²¹¹. This elevation in bcl_2 resulted in a significant suppression of radiation induced apoptosis at 48 hrs, but after longer periods post irradiation even the bcl_2 expressing cells underwent apoptosis. This suggests that bcl_2 over expression does not provide a permanent protection against radiation induced apoptosis but rather it delays the onset of the process ²¹¹.

The angiogenic switch stimulates angiogenesis at a distinct time during tumorigenesis. Through the stimulation of angiogenesis and the inhibition of apoptosis tumour cells attempt to negate the effect of hypoxia. This was investigated using prostate cancer cell lines transfected with bcl_2 . These cell lines were subjected to hypoxic conditions in vitro. Cells with transfected bcl_2 expressed significantly more vascular endothelial growth factor (VEGF) than the controls ¹¹⁶. The cells were then injected subcutaneously into nude mice; the bcl -transfected tumours grew faster, expressed more VEGF and a greater micro vessel density ¹¹⁶. The bcl_2 transfected cells were implanted in mouse corneas, and six of the eight corneas implanted with transfected cells had evidence of neovascularisation compared to none of the controls. The authors of his study concluded that increased VEGF expression and the resulting increased vascularisation associated with this are essential components of the

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

in vivo growth advantage conferred to tumours expressing bcl_2 ¹¹⁶.

Bcl_2 over expression occurs in colorectal adenomas, cervical intra-epithelial neoplasia and other pre-malignant lesions. The basal cells of the prostatic epithelium probably house the prostatic stem cell population⁴⁰. These cells are resistant to androgen dependant apoptosis, where as secretory luminal cells require androgens and undergo apoptosis when deprived of androgens. The expression of bcl_2 is not uniform within the prostate, the transitional and peripheral zones exhibit strong expression for bcl_2 in the basal cells but not in the secretory luminal cells. The central secretory zone does express heterogeneous expression of bcl_2 ⁴¹. In a series of 216 foci of high grade PIN from 25 radical prostatectomy specimens, Bcl_2 expression was restricted to the basal cells in 169 specimens (78%), and was present in secretory luminal cells in 22%. No difference was detected in bcl_2 expression with and without adjacent cancer. There was a marked inverse relationship between the expression of the androgen receptor and bcl_2 . In the transition and the peripheral zones the secretory luminal epithelium strongly expressed the nuclear androgen receptor. The basal cells were unreactive for androgen receptor staining. High grade PIN expressing bcl_2 in the basal cells only stained for the androgen receptor in the luminal secretory cells. Where as high grade PIN over expressing bcl_2 in the luminal cells showed a marked reduction in expression of the androgen receptor. The inverse correlation between bcl_2

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

and androgen receptor expression was highly significant. It has therefore been postulated that the aberrant expression of bcl_2 in sub-sets of PIN may indicate resistance to androgen dependent cell death, and probably reflect reduced androgen-responsiveness of the dysplastic epithelium ⁴¹.

The development of androgen resistance can be associated with bcl_2 expression, it can also occur independently of bcl_2 . This has been demonstrated by work carried out on the Dunning AT-3 cell line that is highly malignant but does not express bcl_2 ¹²².

3.41 Bcl_2 and Prostate Cancer

A literature review was carried out using the Medline database. The initial keyword used was bcl_2 and the second was prostate neoplasm. For the first keyword a total of 6302 references were detected and for the second 33298. These two searches were combined and the number of references decreased to 160 references. The abstracts of these 160 citations were reviewed and a total of 50 pertinent references were reviewed in detail

10:15:16:20:21:21:22:27:41:43:55:58:60:66:7476:116:122:125:132:142:148:173:179:185:187:191:204:211:217:
222:227:235:238:239:245:246:255:256:274:308:315:341:348:364:372:381:382:385:385:394

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

3.42 Bcl₂ and Radical Prostatectomy.

To study the significance of bcl₂ a cohort of 137 consecutive radical prostatectomy specimens obtained from patients who underwent radical prostatectomy between 1978 and 1993 at the University of Basel were examined for over expression of bcl₂ and p₅₃⁶⁰. 40 patients were stage pT₂, 93 patients were pT₃, no stage was given for 4 patients, 103 patients were pN₀ and 34 were pN_{1/2}. Follow up data was available from 115 patients with a mean follow up of 5.2 years. Only eight patients had died, therefore survival data was not discriminatory, but progression data was available. 54 patients had received adjuvant local radiotherapy. Orchiectomy was performed upon six patients at the time of prostatectomy. Progression was documented in 37 patients based upon PSA increase in 29 patients, biopsy in 3 and a positive bone scan in 5. Bcl₂ over expression was associated with higher stage, shortened disease free survival. This study also stained for Ki-67 and p₅₃ expression. Progression analysis showed that patients with bcl₂ positive and Ki 67 positive tumours had the shortest progression free survival, and those with bcl₂ and Ki 67 negative tumours had the longest disease free survival⁶⁰.

The group from the Walter Reed Army Medical Centre, Washington carried out immunohistochemistry on 175 radical prostatectomy specimens for both p₅₃ and bcl₂²⁰. One hundred and seventy four of the patients had capsule confined disease. Over expression of bcl₂ was detected in 47 of 175 (26.9%)

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

patients, these patients had a statistically significant higher 5 year failure rate when compared to bcl₂ negative patients i.e. 67.0% versus 30.7%. Combining the bcl₂ positive and the p₅₃ positive patients, the failure rate was 75.3%, compared to a failure rate of only 20.4% if both bcl₂ and p₅₃ were negative²⁰. Similar results were obtained using a consecutive series of 208 radical prostatectomies and a mean follow up of 4 years. Twenty-eight cases (13%) expressed bcl₂; bcl₂ positivity was an independent prognostic marker in a multivariate analysis¹⁹⁴.

Specimens obtained from a heterogeneous series of 40 patients with prostate cancer were used to determine the association between disease progression and bcl₂, waf-1, and p₅₃. Twelve patients over expressed bcl₂, of these ten progressed; twenty-eight patients did not express bcl₂ of these 19 progressed. There was no correlation between the expression of bcl₂, clinical grade, stage and progression⁶⁶.

Apakhama and colleagues studied seventy seven patients with prostate cancer at various stages were followed up for a median of 30 months and the importance of p₅₃ and bcl₂ assessed. The age range was from 46 to 88 years. Forty-five patients (58.4%) had metastases, 37 patients stained positively for bcl₂. Seventeen of these patients stained positively for both p₅₃ and bcl₂, of these 16 had had hormonal manipulation, and 13 had hormone refractory

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

disease. These workers postulated that the combined detection of p₅₃ and bcl₂ may be useful in detection of hormone refractory disease ¹⁰.

A study of 235 patients, 172 with no metastases which had been treated by a number of modalities including seven with radical prostatectomy, 4 with radical radiotherapy, 22 with oestrogen, 61 with orchiectomy, 33 patients had a TURP and 82 patients were treated conservatively ²²⁷. 71% of the tumours were bcl₂ negative, 18% were weakly positive, and 11% were strongly positive. The expression of bcl₂ correlated with higher tumour stage, metastatic disease, and high grade and in a univariate analysis with poor survival, but in a multivariate analysis the expression of bcl₂ had no prognostic significance. In a study of 41 patients with lymph node metastases from a Japanese centre, bcl₂ expression was observed in both primary tumours and metastases from 14 tumours ²³⁸. Bcl₂ expression was found to be of no prognostic value.

3.43 Bcl₂ and Radical Radiotherapy.

A study of 43 patients who had been treated with definitive radiotherapy for prostate cancer with a follow up period of at least 3 years showed increased bcl₂ staining predicted relapse¹⁷³. A study of 54 patients, who were treated with radiotherapy, showed that increased staining of bcl₂ or p₅₃ predicted relapse ³⁴¹.

3.44 Bcl₂ and Watchful Waiting.

In the Scandinavian nations prostate cancer is often treated conservatively, a study of 221 men diagnosed by TURP chips and treated conservatively with a median follow-up of 15 years demonstrated that 114 cases over expressed bcl₂. This correlated with clinical stage and disease specific survival. If the tumours were further divided into those that were p₅₃ positive and bcl₂ negative compared to p₅₃ positive and bcl₂ positive, those in the first group did better than the second. But dividing the p₅₃ negative group into those that were bcl₂ positive versus those that were bcl₂ negative made no difference to disease free survival⁴³.

3.45 Bcl₂ and Recurrence.

The relationship between the primary tumour and localised recurrence after definitive treatment was studied in a series of recurrent tumours after definitive surgery (n=18) or radiotherapy (n=13). There was an increase in the expression of bcl₂ in recurrent tumours when compared to the primary lesion but this did not reach significance¹⁴². This was further studied in a series of 32 patients who had failed radiotherapy and had pre and post radiotherapy specimens, prior to radiotherapy 14 patients expressed bcl₂ and post radiotherapy 19 expressed bcl₂. In the group of patients that had no disease

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

recurrence post radiotherapy only one out of the 12 patients expressed bcl₂. This was highly significant. If both p₅₃ and bcl₂ were examined together significantly more patients had p₅₃ and bcl₂ immunopositivity following radiation failure, 48%, compared to 8% in those that were successful¹⁷³.

3.46 Bcl₂ and Prostate biopsies.

The role of prostate biopsies at predicting outcome was studied in 146 patients. Bcl₂ positivity was detected in 29 patients. Bcl₂ positivity was detected in 23.7% (18/76) of patients with localised disease, and in 15.7% (11 of 70) of patients with metastatic disease. Using multivariate analysis bcl₂ positivity was an independent predictor of outcome²³⁹.

The group from Bristol compared preoperative biopsy assessment of bcl₂, p₅₃, CD-44 and e-cadherin with the expression of these markers in the respective radical prostatectomy specimen⁵⁵. Radical prostatectomy and biopsy specimens were examined from a total of 76 patients; aberrant staining for bcl₂ was detected in eight preoperative biopsies and in 15 radical prostatectomy specimens. Brewster et al did not comment if preoperative biopsy staining for bcl₂ correlated with expression of bcl₂ in the radical prostatectomy specimens. Preoperative bcl₂ was not a significant predictor of outcome. Using Cox regression no direct relationship between bcl₂ over expression and p₅₃ protein accumulation could be established. Using a multivariate analysis post-

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

operative p₅₃ and bcl₂ were independent prognostic predictors of disease free survival⁵⁵.

Similarly, in a series of 117 preoperative samples from radical prostatectomies carried out at the Walter Reed Medical Centre the preoperative biopsies did not correlate with outcome³⁶⁴. Forty-nine patients had evidence of disease recurrence, and 18 preoperative samples had evidence of bcl₂ over expression. Comparison between biopsy and prostatectomy staining revealed no concordance. Biopsy samples could not predict outcome³⁶⁴.

Biopsies have also been used to predict outcome in-patients treated with definitive radiotherapy. In a series of 52 patients with pre radiotherapy biopsy material available for immunostaining, thirteen stained positively for bcl₂. Of these, 11 patients (84.6%) failed radiotherapy, in the bcl₂ negative group only 16 of the 39 patients (41.1%) failed radiotherapy. This result was significant³⁴¹.

3.47 Androgen deprivation and Bcl₂

The role of androgen ablation in changing the expression of bcl₂ and p₅₃ was studied using radical prostatectomy specimens obtained from 106 patients was compared with 71 pre treatment prostate biopsies⁷⁴. The patients prior to surgery had 3 months of LHRH treatment. Eleven of the 106 radical

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

prostatectomy specimens over expressed bcl₂ and nine of the 71 pre treatment biopsies expressed bcl₂. No correlation was found between the expression of bcl₂ in the biopsies and prostatectomies, four cases were positive in both samples, four only in the biopsies and three only in the prostates. No correlation existed with bcl₂ over expression and clinical outcome, preoperative PSA, or accumulation of aberrant p₅₃⁷⁴. However the data from this study must be viewed with caution since in studies comparing biopsies and radical prostate specimens, the expression of bcl₂ in material obtained from biopsies does not correlate with the expression of bcl₂ obtained from radical prostatectomy samples.

An earlier study comparing two groups of radical prostatectomies, one that had neoadjuvant hormonal therapy (n=28) with ones that had not (n=51), found increased expression of bcl₂ after neoadjuvant hormonal ablation³⁹⁴.

A study of 69 heterogeneous patients with prostate cancer demonstrated a trend towards increased bcl₂ expression post androgen deprivation¹⁶. However in this study the authors could not demonstrate any correlation between the expression of bcl₂ and apoptosis, p₅₃, or clinical outcome.

This effect of increased bcl₂ with castration was further investigated in series 15 responders and 13 non-responders of androgen ablation³⁷². The patients

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

had biopsies pre androgen ablation and one week after treatment. Both groups had an increase in bcl₂ expression associated with androgen ablation, but the responders had significantly increased bcl₂ expression as compared to the non-responders³⁷².

3.48 Bcl₂ and High Grade Prostatic Intra Epithelial Neoplasia (PIN).

High-grade prostatic intra epithelial (PIN) neoplasia is a putative precursor of prostate cancer, and the incidence of bcl₂ expression was studied in PIN using 43 radical prostatectomy specimens. Bcl₂ was over expressed in 1 of the 43 cancers (2.3%), and in 15 of the 43 PIN specimens (34.9%)¹⁸⁷. These authors suggested that the foci of bcl₂ positive PIN might not be the direct precursors of invasive cancer.

3.49 Bcl₂ and Metastases.

Prostate cancer often metastasises to the bone marrow. The accumulation of bcl₂ was studied in 43 bone marrow samples obtained from patients with bone metastases. All patients had been treated with at least hormonal ablation. Bcl₂ was detected in 14 of the 43 cases. Six of these cases exhibited only positive staining for bcl₂, and eight expressed both p₅₃ and bcl₂. The expression of bcl₂ protein did not significantly influence patient survival²⁴⁵.

The relationship between bcl₂ expression and metastasis was in hormone naïve

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

patients showed that 5 of 30 (17%) patients with lymph node disease had positive bcl₂ staining and 14 of 27 (52%) patients with bone metastasis had positive staining in bone marrow samples ¹²². Therefore bcl₂ expression increases with increasing stage.

3.50 Bcl₂ and Bax.

Prostate cancer is often managed with external beam radiotherapy. The prognostic importance of bcl₂ and bax were investigated in a series of 41 patients who had undergone external beam radiotherapy for the management of localised carcinoma of the prostate. 21 patients responded to radiotherapy and 20 failed to respond as determined by a PSA nadir of 1.0ng/ml. Bcl₂ expression was significantly higher in the non-responders and bax expression was higher in the responders. The ratio of bcl₂/bax was significantly higher in the non-responder when compared to the responders' ²³⁵.

Table16: The relationship between bcl₂ expression and outcome in patients with radical prostatectomy.

Ref.	Year of pub	Type of specimen	No of cases	No localised	No locally invasive	No mets	Method	Results/Conclusions
⁶⁰	1996	Radical Prostatectomy	137	40	93	34	IHC	T2 2/40 (5%) + T3 29/93 (31%)+ N0 25/104 (24%)+ N1/2 8/34 (24%)+ Bcl-2 expression increased with stage.
²⁰	1996	Radical Prostatectomy	175	T1=25 T2=149	T3=1	0	IHC	47/175 (27%) Bcl2 positive. Correlated with poor survival
¹⁰	1996	Heterogeneous	77			45	IHC	37(48%) positive for bcl2. 20/37 androgen resistant
⁶⁶	1997	Heterogeneous	40	T1=10 T2=5	T3=11 T4=13	21	IHC	12/40 (30%) Bcl-2 + 10/12 progressed. No correlation with bcl-2 + and clinical outcome
²²⁷	1997	Heterogeneous	235	T1= 94 T2 = 56	T3=96 T4=19	63	IHC	18% weakly positive 11% strongly positive. Bcl-2 correlated with poor survival, but of no an independent prognostic value.
¹⁹⁴	1998	Radical Prostatectomy mean follow up 4yrs.	208	111	97	8	IHC	28/208 (13%) + Bcl-2 correlated in a multivariate analysis with poor outcome.
²³⁸	1998	Biopsies and Radical Prostatectomies	41	?	?	41	IHC	14/41 (34%) bcl-2 +. No correlation with bcl-2 + and clinical outcome.
¹⁴²	1998	Radical Prostatectomy	18	11	7	5	IHC	Increased expression of bcl-2 in recurrence. Not significant.

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

¹⁴²	1998	Biopsies DXT	13	13	0	0	IHC	Increased expression of bcl-2 in recurrence. Not significant.
⁷⁴	2000	Radical Prostatectomy	106	58	31	17	IHC	11/106 + No correlation with bcl-2 + and clinical outcome
⁴²¹	2003	Radical Prostatectomy	70	70			IHC	3/70 + Increased expression of bcl2, predicted relapse.
³¹⁹	2005	Radical Prostatectomy	131	60	71	10	IC	13/131 bcl2+ Increased expression of bcl2, predicted relapse.

Table17: The relationship between bcl₂ expression and outcome in patients treated with radical radiotherapy.

Ref.	Year of pub	Type of specimen	No of cases	No localised	No locally invasive	No mets	Method	Results/Conclusions
²³⁵	1998	Pre DXT biopsies	41	41	0	0	IHC	Significantly higher bcl-2/bax ratio in radiation failures.
¹⁷³	1998	Biopsies and rad prostatectomy	43	43	0	0	IHC	1/12 bcl-2 + in the radiation success, 14/31 bcl-2 + in the radiation failures. Bcl-2 was found in significantly more radiation failures.
³⁴¹	1999	Pre-DXT biopsies	52	52	0	0	IHC	13/52 (25%) bcl-2 +. 11/13 bcl+ cases failed treatment with DXT.
³²⁸	2003	Needle biopsy	40	29	11	0	IHC	11/20 patients with recurrence overexpressed bcl2, 0/20 in the recurrence free group.

Table18: The relationship between bcl₂ expression in pre operative biopsies and outcome.

Ref.	Year of pub	Type of specimen	No of cases	No localised	No locally invasive	No mets	Method	Results/Conclusions
²³⁹	1997	Biopsies	146	50	28	68	IHC	29/146 bcl-2 +. In multivariate analysis an independent prognostic indicator.
³⁷²	1998	Pre and post castration biopsies in 15 responding patients and 13 patients not responding to hormonal therapy.	28	7	21	20	IHC	Bcl-2 expression increased significantly with androgen ablation, significantly more so in the responders.
³⁶⁴	1999	Pre op biopsies	117				IHC	18/117 (15%) bcl-2 +. No correlation with bcl-2 + and clinical outcome.
⁵⁵	1999	Pre op biopsies	65				IHC	8/65 bcl-2 (12%) + Pre-op Bcl-2 did not correlate with outcome.
⁷⁴	2000	Pre op biopsies	71					9/71 bcl-2 + No correlation with bcl-2 + and clinical outcome
⁴³	2000	TURP biopsies	221	125	96	60	IHC	Patients treated with watchful waiting. 114 (52%) cases + for bcl-2. Correlated with clinical stage and disease specific survival.

Table19: The relationship between bcl₂ expression and metastasis.

Ref.	Year of pub	Type of specimen	No of cases	No localised	No locally invasive	No mets	Method	Results/Conclusions
¹²²	1996	Lymph node (30) & Bone marrow(27)	57	0	0	57	IHC	5/30 lymph node bcl-2 + 14/27 bone marrow bcl-2 + Bcl-2 expression increases with stage.
²⁴⁵	1997	Bone marrow.	43	0	0	43	IHC	14/43 (32%) bcl-2 +. No correlation with bcl-2 + and clinical outcome.

3.60 BAX

Bcl₂ associated X protein, or bax is a proapoptotic molecule, which dimerises with the apoptotic Bcl₂. This dimerisation inhibits the action of bax. The dimerisation takes place at the homology regions and all family members share at least one of the four regions- BH1, BH₂, BH3 and BH4¹²⁴. The truncated form of Bax, expressing only BH3, is sufficient to induce apoptosis and antagonise the effects of ant-apoptotic members of the family¹⁵⁶. Induction of bax is promoted by the initiation of apoptosis. This causes the monomeric bax to translocate from the cytosol to the mitochondrial membrane; here it becomes an integral part of the membrane and cross link's as a homodimer, resulting in cell death by the release of cytochrome c from the mitochondria¹⁴¹. This ability to target the mitochondrial membrane is regulated by the terminal 20 amino acids. Bcl₂ inhibits bax-mediated apoptosis. The bax gene promoter is p₅₃ responsive and so p₅₃-binding results in transactivation of the bax gene. Therefore, bax up regulation is one method by which p₅₃ exerts its pro-apoptotic action²⁶¹.

A literature review was carried out using the Medline database. The initial keyword was prostate cancer and this revealed 35234 citations, the second keyword was bax and the citation count was 23802. When the two citations were combined the number decreased to 194. The abstracts were reviewed

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

and a total of 15 citations were selected as pertinent

9:81;101;117;122;132;187;199;204;226;230;235;235;313;330;422

3.61 Incidence of Bax

The incidence of bax was studied in 64 radical prostatectomy specimens. Bax was detected in every specimen²⁰⁴. In Asians the bax was present in 47 of 62 radical prostatectomy specimens i.e. 76%. In a univariate analysis bax expression predicted a significantly poorer survival⁸¹.

3.62 Bax and radiotherapy for prostate cancer.

A retrospective review of 41 patients who underwent radiotherapy with curative intent showed that the patients could be divided into two groups depending upon the PSA response and the presence of tumour in the post operative biopsies. This demonstrated no difference in bax immunoreactivity between responders and nonresponders²³⁵, but the staining of bcl₂ was increased in non responders and so the ratio of bcl₂/bax was higher in non-responders²³⁵. The importance of the bax: bcl₂ ratio was further assessed by using synthetic peptide sequences from the BH3 domain of bax, this resulted in binding between the BH3 peptides and bcl₂ and so blocking bcl₂/bax heterodimerisation. The excess of released bax led to an increase in apoptosis¹¹⁷.

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

Using adenovirus vectors, bax was over expressed in four different cell lines. In all the cell lines bax over expression caused cytochrome c release and initiation of the capsase mediated apoptosis pathway. Bax over expression by passed the anti apoptotic effect of bcl₂, and caused apoptosis in cell lines resistant to chemically induced apoptosis²²⁶. This finding has been confirmed by a second group²³⁰. Androgen withdrawal is known to cause apoptosis; the presence of androgens in androgen responsive cell lines causes attenuation in the expression of bax¹⁹⁹. Using an adeno virus construct, which contained the bax gene, linked to androgen responsive elements and with probasin as a promoter. Probasin is a prostate specific promoter, so that the molecule can be over expressed in prostate only. This model has shown that bax expression and so apoptosis could be made to occur in androgen dependant manner, both in vitro and in mice⁹.

3.63 Bax and high grade prostatic intraepithelial neoplasia.

In a heterogeneous group of 44 radical prostatectomy specimens, Bax expression can be detected in all cases of PIN¹⁸⁶.

3.70 Ki 67

The name of the Ki-67 protein is derived from its city of origin – Kiel, and the number of its original clone in a 96 well plate. The Ki-67 antigen is a nuclear protein that is present in proliferating human cell ¹²⁷. The genetic locus has been assigned to chromosome 10 and consists of 30,000 base pairs^{127:345}. Two isoforms of the protein have detected with masses of 320 and 359kD; both contain a large central region with 16 repetitive elements ³⁴⁵. Homology searches have failed to detect regions of similarity between Ki-67 and other proteins. Although the functional role of Ki-67 is unknown, it is expressed in all stages of the cell cycle (G1, S, G₂, and mitosis) except for G₀. Ki-67 immunoreactivity appears to be a valid method to measure proliferation ³⁴⁶. Ki-67 is a part of the nuclear matrix and of the chromosome scaffold. Initially the use of Ki-67 was limited to frozen section material, however the development of antigen retrieval systems has led to newly developed antibodies being applied to formalin fixed material ¹⁹⁶.

The proliferation rate is low in prostate cancer but it may be important in determining tumour progression. Proliferation assessed by mitotic rate ⁴⁰², S-phase fraction ^{392:404} and Ki-67 ^{159:247} has been related to outcome³⁷⁰. However at present the clinical significance is far from established. Using archival material and correlating this with outcome it has been shown that proliferation as assessed by Ki-67 immunoreactivity is higher in prostate

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

cancer than in benign tissue^{1:84:389}. Indeed the grade of the tumour is related to the expression of Ki-67^{155:260:285}. Prostate hyperplasia and well-differentiated tumours are negative for the expression of Ki-67²⁸⁵. The Ki-67 expression can be used as a marker for predicting outcome in prostate cancer^{1:155:370}. In prostate biopsies the expression of Ki-67 can predict local extra capsular spread but not metastasis²²⁹.

The cellular proliferation index is also a predictor of recurrence after definite local therapy i.e. radiotherapy or radical prostatectomy¹⁴². In patients with lymph node metastasis the Ki-67 labelling index of the primary tumour and the lymph nodes had an independent prognostic value where as p₅₃ and bcl₂ had no prognostic value¹⁴². Others have not found Ki-67 so useful³⁹⁷.

3.71 Why Study Cell Proliferation?

A major stimulus to the study of cell proliferation is the belief that quantification of this fundamental biological process would provide objective data with which to more effectively categorise tumours, predict prognosis and consequently result in tailoring therapy to individual patients. In the past a number of techniques have been used^{152:154}. For example, mitotic index estimation by light microscopic examination, in vivo (or in vitro) incorporation of thymidine analogues into DNA during S phase, flow cytometric assessment of DNA index. These methods have significant

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

disadvantages such as ethical, methodological problems and loss of tissue architecture. An alternative method which is simple, reliable, reproducible and allows retention of normal histology has been developed using antibodies that recognise, by immunohistochemistry, molecules whose expression is linked to the cell cycle^{152:244}.

3.72 Cell Cycle

The cell cycle describes a co-ordinated series of biochemical alterations that allow exact duplication of the DNA content in a cell (S phase) followed by precise segregation of the DNA into two sets and their subsequent partition into two daughter cells during mitosis (M phase). After mitosis the daughter cells enter a quiescent period of varying duration before commitment to a further cell cycle. The period between one M phase and the next S phase is called G₁. During G₁ phase there is growth and preparation of the chromosomes for replication. The period between S phase and the following M phase is called G₂. The cell undergoes preparation for replication during the G₂ phase. Regulated alteration in expression and function of regulatory and structural proteins occurs during the cell cycle. This cyclical alteration in proteins permits the use of antibodies to define antigens expressed in a cell cycle related manner.

Potential antigenic targets for the definition of cells within the cell cycle are

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

numerous. One of the most widely used antibody is Ki-67¹²⁷, which reacts with a nuclear non-histone protein of 395 and 345 kD present in all active parts of the cell cycle, i.e., G₁, S, G₂, and mitosis, but is absent in G₀^{127:208}.

The evidence for the link between the expression of the Ki-67 antigen and the cell cycle is overwhelming^{57:337}. The molecule begins to be expressed in mid G₁ increasing in level through S and G₂ and peaking in M¹²⁷. It is very rapidly catabolised at the end of M. There is good relationship between detectable expression of Ki-67 and growth fraction in several model systems^{244:346}. Furthermore, there appears to be no expression of Ki-67 during DNA repair processes unlike some other cell cycle associated molecules (for example PCNA). The function of Ki-67 remains unclear but it may represent a structural protein that maintains the higher order structure of DNA during the important events of mitosis³³⁷. The major drawback of this antibody is that its application is restricted to fresh or frozen tissues and the epitope does not survive routine histopathological fixation in formaldehyde.

3.73 MIB-1

Recently monoclonal antibodies, MIB 1-3 (MIB for Molecular Immunology Borstel), against recombinant parts of the Ki-67 antigen have been developed. These antibodies are true Ki-67 equivalents, as demonstrated by immunostaining of fresh specimens, biochemistry, and molecular biological

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

techniques^{73:195}. Formerly immuno-histochemistry on formalin-fixed, paraffin-embedded sections failed to stain for Ki-67. However, when dewaxed microwave oven-processed paraffin sections of formalin-fixed tissues were used, MIB 1 and MIB 3 gave strong nuclear staining of proliferating cells under a variety of normal and neoplastic conditions^{73:286}. This method is reproducible, easy to perform at low cost; no additional technical skill is needed and allows a much better recognition of cellular details and, therefore, a better identification of positive cellular subsets. This new method has been successfully applied to tissue sections stored for a long time. The assessment of cell kinetics through the detection of Ki-67 antigen is now possible on archival material.

By immunostaining tissue with a suitable antibody a proliferation index (P.I.) can be calculated as the ratio of nuclear area number positively stained to total nuclear area number in a fixed high power field.

3.80 Ki-67 and Prostate Cancer.

A literature review was carried out using the Medline database. The initial keyword used was prostate cancer and the second was Ki-67. For the first keyword a total of 4250 references were detected and for the second 37565. These two searches were combined and the number of references decreased to 145 references. This was further limited to English language abstracts and five

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

further abstracts were eliminated. The abstracts of these 140 citations were reviewed and a total of 43 pertinent references were reviewed in detail

1:15:30:43:59:62:63:84:90:109:115:142:155:158-160:187:194:197:228:229:238:241:260:274:275:285:286:295:

334:338:345:368:379:379:389:393:394:397:398:403:411:414

3.81 Ki-67 expression and outcome in patients with radical prostatectomy.

In a series of 180 radical prostatectomies with a follow up of 1 to 9 years, MIB1 scoring was high in 168 specimens and absent or low in 12 specimens. The five-year recurrence free survival correlated with MIB1 expression. The authors concluded that MIB 1 expression could be used to determine a subset of patients who are more likely to relapse³⁰.

In a series of 92 radical prostatectomy patients carried out in the Netherlands, the expression of Ki-67 (MIB 1) correlated with recurrence but this did not reach significance in a multivariate analysis⁴⁰³.

In a series of 137 consecutive radical prostatectomies, with a mean follow up of 5.8 years available for 108 patients, a Ki-67 labelling index above 7.5% was predictive of recurrence in a multivariate analysis²⁰⁷. A similar study of 29 patients using a cut off of 10% also showed a statistically significant association between the proportion of Ki-67 cells labelled and biochemical

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

relapse¹⁵. A study of 104 patients focussing on the Ki-67 staining in the area of highest tumour grade with a median follow up of 56 months showed a median staining of 6.7% (range 1.2 – 42.6). Using a cut off of 6.7%, the Ki-67 staining was a predictor of relapse in a multivariate analysis¹⁵⁵.

A recent study of 122 patients who underwent radical prostatectomy selected 22 patients with “metastatic disease” (pT3b-4aN0M0 and pTanyN1M0) and 18 “nonmetastatic” (\leq pT3aN0M0). At a mean follow up of 33 (4 – 78) months, the proliferation index predicted relapse in both groups³⁷⁶.

A study of 20 radical prostatectomy specimens showed that the Ki-67 score is greater for peripheral zone tumours when compared to transition zone cancer¹¹⁵; this is in concordance with more aggressive behaviour of peripheral zone tumours.

3.82 Ki-67 expression and outcome in patients with radical radiotherapy.

Animal studies using radiated heterotransplanted DU-145 cell lines and comparing the expression of Ki-67 in the radiated and control groups have not shown a significant decrease in proliferation activity²⁸⁵.

Pre-radiotherapy material was stained from 106 patients who had definitive

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

prostate radiotherapy. The mean Ki-67 labelling index was 3.2%, and increased in high grade and high stage disease. A labelling index greater than 3.5% predicted early biochemical failure⁹⁰.

A further study of 42 patients with prostate cancer, both capsule confined and extra capsular, demonstrated that Ki-67 as a measure of cell proliferation could predict recurrence in both groups in a multivariate analysis³³⁸. A later study confirmed these findings¹⁹⁷.

In a series of 55 patients who had received radical radiotherapy and been followed up for a mean of 6.8 years, residual cancer was found in 67% (37/55) of patients, of these 64% (23/36) had active tumour cells as determined by Ki-67 staining. These authors concluded that the majority of patients with tumour on their post radiotherapy biopsies had viable tumour²²⁸.

A study comparing pre radiotherapy biopsies with post radiotherapy biopsies in 13 patients demonstrated an increase in Ki-67 score, between the pre and post biopsies. The score increasing from a mean of 5.6 to 10.5 in those patients that relapsed¹⁴².

3.83 Ki-67 expression and outcome in patients treated with androgen withdrawal.

A study of 153 unselected patients with prostatic carcinoma demonstrated that Ki-67 score was related to poorly differentiated tumour, and a high Ki-67 score was related to metastatic disease and poorer survival. In particular if patients with metastatic disease were stratified according to their Ki-67 scores after androgen withdrawal those with scores > 1% had a poorer outcome¹⁵⁹.

A study of 190 treated patients with prostate cancer with a mean follow up of 12.9 years, demonstrated that Ki-67 was significantly related to stage and grade. In a multivariate analysis Ki-67 expression had no independent predictive value in organ confined disease but was an independent prognostic factor over the entire cohort¹.

3.84 The effect of androgen ablation on Ki-67.

A study of 18 patients who had biopsies before and after androgen ablation showed a decrease in Ki-67 score after androgen ablation⁴¹⁴. A further study comparing 28 patients who had received neoadjuvant androgen ablation with 51 patients who had not, demonstrated that androgen ablation leads to a significant decrease in MIB-1 index i.e. 7.5% compared to 14.4% respectively³⁹⁴.

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

3.85 Ki-67 and PIN.

In a heterogeneous series of 35 patients the Ki-67 labelling index increased in a stepwise progression when comparing benign tissue, with PIN and cancer ¹⁸⁷.

This has been confirmed by others ^{160:275:379:393}.

Table20: The relationship between Ki-67 (MIB1) expression and outcome in patients with radical prostatectomy.

Ref.	Year of pub	Type of specimen	No of cases	No localised	No locally invasive	No mets	Method	Results/Conclusions
³⁰	1996	Radical Prostatectomy follow up 1-9 yr.	180	72	107	1	IHC	18 pts had low MIB1 expression. Significant correlation between MIB 1 score and nuclear grade. Gleason score, pathological stage. Low MIB1 score correlated with low chance of recurrence.
²⁰⁷	1996	Radical Prostatectomy follow up 1-15 yr.	137	44	93	34	IHC	Significant correlation between recurrence and Ki-67>7.5%.
¹⁵	1999	Radical Prostatectomy follow up 3yrs	29	39	?	8	IHC	High MIB1 expression associated with recurrence, and an independent prognostic factor in a multivariate analysis.
⁴⁰³	2000	Radical Prostatectomy follow up 0-17 yr.	92	65	27	6	IHC	High MIB1 expression associated with recurrence, but was not an independent prognostic factor in a multivariate analysis.
¹⁵⁵	2001	Radical Prostatectomy median follow up 56 months.	104	32	72	7	IHC	High MIB1 expression associated with recurrence, and an independent prognostic factor in a multivariate analysis.
⁸⁴	2002	Radical Prostatectomy	24	13	11	0	IHC	High Ki-67 expression in cancer.
¹⁷⁷	2005	Radical Prostatectomy	52	14	38	0	IHC	20/52 progressed. 16 Ki67+ relapsed and 4 Ki67-, Ki-67 predicted relapse.
²⁷⁶	2005	Radical Prostatectomy	40	6	12	22	IHC	High Ki-67. predicted progression, and metastatic disease.

Table21: The relationship between Ki-67 (MIB1) expression and outcome in patients with radical radiotherapy.

Ref.	Year of pub	Type of specimen	No of cases	No localised	No locally invasive	No mets	Method	Results/Conclusions
²²⁸	1996	Biopsies	55	?	?	?	IHC	Post DXT, there was still active proliferation.
³³⁸	1997	Biopsy/TURP	42	30	12	0	IHC	18/23 patients with recurrence had high Ki-67 scores, compared to 6/19 with no recurrence.
¹⁴²	1998	Biopsy	13	13	0	0	IHC	Comparing pre-treatment with post treatment biopsies in radiation failures showed an increase in Ki-67 from 5.6 to 10.5
¹⁹⁷	1999	TURP	42	35	7	0	IHC	The treatment failures had a significantly higher Ki-67 score compared to the responders.
⁹⁰¹	2002	Biopsy/TURP	106	74	32	0	IHC	Ki-67>3.5% predicts treatment failure.
²²⁵	2004	Biopsy	108	31	77	0	IHC	A staining index of greater than 3.5% was associated with a risk of developing local progression, and disease specific mortality.

Table 22: The relationship of Ki-67 to outcome in patients treated with androgen withdrawal.

Ref.	Year of pub	Type of specimen	No of cases	No localised	No locally invasive	No mets	Method	
⁴¹⁴	1995	Serial biopsy	18	3	6	9	IHC	Androgen ablation causes a decrease in Ki-67.
¹	1997	TURP/Biopsy	190	106	84	66	IHC	Ki-67 increased with grade and stage. In a multivariate analysis Ki-67 staining was an independent predictor of outcome.
³⁷⁰	1997	TURP mean follow up 71 months.	125	89	16	15	IHC	Ki-67 increased with grade and stage. In a multivariate analysis Ki-67>3% was an independent predictor of outcome.

Chapter 4:

Aims

- To determine the relationships between the volume of high grade prostatic intraepithelial neoplasia, cancer and benign tissue.
- To determine the relationship between serum PSA and the volume of high grade prostatic intraepithelial neoplasia, cancer and benign tissue.
- To determine using immunohistochemical techniques the levels of p53, bcl₂, bax, and Ki-67 in formalin fixed benign, malignant and tissue with PIN.
- To determine using immunohistochemical techniques the effect of androgen ablation on the levels of p53, bcl₂, bax, and Ki-67 in formalin fixed malignant tissue.

Chapter 5: Materials and Methods.

5.1 Methods

This study was divided into two parts; the first part was a prospective assessment of tumour volume, PIN volume, and total gland volume. This was carried out in 76 radical prostatectomy specimens. The second part was a retrospective assessment of the expression of p₅₃, bcl₂, bax, and Ki-67. This was carried out on paraffin fixed specimens identified from 97 radical prostatectomy specimens.

All radical prostate specimens were obtained from patients undergoing radical retro pubic prostatectomy. The patients preoperatively were staged with clinical examination, serum total PSA (Hybertech Assay), transrectal ultrasound guided sextant biopsy, CT scan and bone scan. A single surgeon (A V Kaisary) carried out the operations and a single pathologist carried out all the initial histopathological assessment (M Jarmulowicz).

The specimens were prepared as follows.

5.2 Fixation

Some authors advocate cutting the prostate into slices fresh and then placing in fixative³⁴⁴; we have not found this a satisfactory technique. It is difficult to cut the prostate fresh and get consistent slices. In addition the tissue is friable and distorted. This leads to curling of the edges. Others believe in inking the fresh prostate and then fixing it whole³⁴³. We feel the best results are achieved if the latter approach is adopted. After weighing, the whole specimen is left to fix in 10% buffered formaldehyde for at least 72 hours. The specimen is not incised to speed penetration of the fixative, as this will distort the subsequent slices.

After fixation the specimen is measured. The margins are painted with Indian ink. After painting the specimen is briefly immersed into Bouin's fixative, which instantly fixes and hardens the Indian ink, so minimising the spread of ink onto the cut surface on slicing. The specimen is then cut.

5.3 Cutting the specimen

The specimen is cut with a slicer. The machine setting is calibrated, using any available fixed tissue, to deliver slices which are 3.5 mm in thickness. Our experience of these machines is that towards the end of the cut, the specimen slice is pulled away producing a curved slice and distorted surgical margin at

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

that point. To prevent this, the specimen is almost cut through, and then withdrawn from the machine and the cut completed using a disposable microtome blade attached to a disposable handle.

There are various opinions on the best way to tackle the apical slice. In our laboratory the apical slice is further cut in the saggital plane and then sliced³⁴³.

5.4 Scanning the slices

Maintaining the correct orientation, both left / right and apex to base, the slices of prostate are laid onto an A4 sheet of transparent celluloid. The surface of the slices to be cut to produce the microscope section is laid face down onto the celluloid. The slices, together with a graduated rule, are then placed in a standard flatbed scanner. The specimen is covered with a white A4 sheet of paper, both to protect the lid of the scanner and to improve the background. As the lid will be three millimetres above the glass the background will have a green hue due to reflection from the lid.

Select the image size to be scanned as A4. This will make subsequent printing at real size easier. Scan the image at 180 dpi with printing at 720 dpi (Figure 14).

Prostatic Intraepithelial Neoplasia and Prostate Cancer.




Fig 14:After fixation the prostate is sliced and scanned.

5.5 Manipulating the image

Some centres use a 'whole mount' of the cut prostate slices, whereas others may not have the facilities and have to use the standard sized cassettes. Using the former method there should be some way of identifying left and right sides on the microscope slide. We use standard cassettes and a print out of the image forms a convenient template to record the numbering of the cassettes. The majority of slices need to be cut into four.

The specimen is then embedded in paraffin wax and processed as per standard protocol (see later). When examining the slides the areas of tumour and high grade PIN are marked on the glass slides using a glass marker. Transferring the tumour and PIN maps from the microscope slides to the computer bitmap image of the prostate slices is not difficult, nor is it a lengthy procedure. The quickest method is to first transfer the map from the microscope slides to a printed image. If the image has been printed at 100% size the tissue sections and printed image will match and slight variation to section shrinkage/stretching does not produce any significant difficulties. The printed image can be traced over the slide using a light box; the latter is the easier and more accurate (fig 15).

Prostatic Intraepithelial Neoplasia and Prostate Cancer.



Fig 15: Areas of PIN and cancer.

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

To achieve the final digitalised image the hand coloured printed hardcopy is re-scanned and overlaid onto the original image. This is straightforward as the latest version of software allows images to be manipulated as a series of separate image layers. The re-scanned image is slightly smaller than the original, because of the small margin introduced by the printer, but this is easily corrected using the resize command of the software, so that the two images are of the same size. The overlaid image, in a separate layer, is set at 50% transparency so that it can be moved to perfectly place it over the original image. A digitalised pad with a pen is used to facilitate this stage rather than a standard mouse. For clarity the marked tumour and PIN on the re-scanned layer is traced in the appropriate colour (Red = cancer, Blue = PIN), each in a separate image layer. The final image can then be printed (figure 16). The patient identification, slice labels and any comments can readily be inserted onto the image using the text facility of the image software. For convenience it is best to have separate image layers for prostate slices, patient/specimen identification, labels and comments, tumour and PIN. Using Paint Shop pro (version 6) the image saved as separate layers is approximately.

This technique provides a high quality image of the cut surface of the prostate, with an overlaid map of tumour and PIN.

Prostatic Intraepithelial Neoplasia and Prostate Cancer.




Fig 16: Composite image of prostate.

5.6 Calculation of total prostate, PIN and tumour volume

The acquired images of prostate slices were transferred to the image analysis program using proprietary software (Sigma Scan Pro, version 6.0 by Jandel Scientific, UK). In order to make accurate measurements of prostate gland areas; a scale bar was included in each image. The image analysis program would use the scale bar for calibration. Area measurements of the regions of interest were performed semi-automatically on Kontron KS400 image analysis software (Zeiss, UK). A macro was written to aid the use of this programme (Appendix 1). The macro was designed to interact with Paint-Shop-Pro images to convert the image from full colour to grey scale and then allow the images to be transported out to the KS400 environment. The regions of interest were then automatically segmented, giving the operator a choice to interact if necessary.

Regions measured included: 1) the entire prostate excluding the slices containing predominantly seminal vesicles, 2) PIN, and 3) cancer. The area of the apical cone was not included in the calculation, since it was difficult to be certain of its exact thickness and so the volume it represented. After the areas had been calculated they were multiplied by the shrinkage factor to correct for any changes in volume during processing of the specimen. All measurements were transferred to Microsoft Excel for statistical analysis.

5.7 Shrinkage factor

Shrinkage factors have been calculated by a number of authorities, and vary from 1 to 1.5^{301;344}. It is important to calculate it for your specific lab. In addition different labs use different techniques to fix the prostate with different fixatives and different protocols for the processing of the prostate³⁴³. Some labs cut the fresh prostate into slices and then fix the gland in slices³⁴³ where as others, ours included, fix the whole gland and then cut it into slices.

In our lab we do not have the resources to process the prostate slices as whole mount so as described previously we cut them into portions that will fit into our cassettes. Taking all this into account we decided to calculate the shrinkage fraction for our lab rather than accept one that may not be applicable to our institution.

The first problem is how to accurately the measure the volume of the prostate. Since prostates are an irregular shape and often compressible it is difficult to get reproducible measurements with a ruler. Therefore, the fresh prostates were taken from the operating theatre, loose tissue was removed and they were weighed and immersed in water and the volume of water displaced weighed. The measurement of the volume was repeated. The prostates are then placed in the fixative, 10% buffered formaldehyde for at least 72 hours.

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

Just prior to slicing the prostates, the measurement of volume and weight was repeated. This gave us the shrinkage factor for the first stage. This method of measuring changes in the volume takes into account any contraction that occurs in the urethral cavity whereas if simply the maximal linear measurement was taken before and after fixation this would not have been taken into account³⁴³. However the volume obtained also includes the seminal vesicles which may have different tissue characteristics with respect to fixation and processing. We did consider removing the vesicles prior to measuring the volumes, but this would have meant considerable dissection of the specimen prior to histopathological assessment of the specimen, and it was felt this would not be in the best interest of the patient. In addition other authorities have included the seminal vesicles when calculating the shrinkage fraction³⁰¹.

Since the slices are now scanned, this factor can be used for determining the total volume of the gland from the images. Then the slices are divided into convenient sections for placing in the cassettes; initially we attempted to use linear measurements with a vernier calliper. However this was not reproducible and there was considerable inter and intra user variation. Therefore this part of the procedure evolved, and now we scan the prostate slices whole after the fixation in formalin and then the processed slides stained with haematoxylin and eosin were scanned (fig 16).

Prostatic Intraepithelial Neoplasia and Prostate Cancer.




Fig 16: H&E stained slices for comparison with slices from fig 14.

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

Thus using the KS-400 software we could calculate the area of these slices and so the second correction factor. This can be multiplied with the previously calculated correction factor can provide an accurate result for prostate cancer and PIN. See appendix2.

5.8 Slice thickness.

The thickness of the prostate slices after fixation in formalin was measured with a pair of Venier callipers. Consecutive slices from six prostates were measured and the average thickness calculated. This was multiplied by the shrinkage fraction. See Appendix 3.

5.9 Preparation of specimens

The paraffin blocks were cooled in an ice bath before using a standard microtome to cut 3-4 micron thick serial sections of all specimens. The sections were floated on water heated to 45° C in a water bath. In some cases they needed to be straightened out with the aid of methyl alcohol covered slides before floating. Sections were examined with the naked eye to ensure that a cross section of the whole block was obtained. Each slide usually contained one piece of serial section from, the same block. The sections were then mounted on to APES coated microscopic glass slides (Appendix No. 4) to avoid tissue loss during micro waving. The slides were immediately labelled

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

using, a lead pencil, with their respective pathology number which consists of a 7 digit code (identical to number on block). The upper left corner of the frosted part of the slide was consistently used to ensure correct orientation when handling them. Slides were then transferred into racks and left for 24 - 48 hours in a warm room at 37⁰ C to dry.

5.10Dewaxing.

Prior to staining the slides had to dewaxed. This was carried out as per appendix 5. Then the specimens could be stained with haematoxin and eosin or with the appropriate antibodies.

5.11Staining methods

There are four main methods of immunoperoxidase staining that can be used to localise cellular antigens. The direct, indirect, PAP, and streptavidin-biotin methods. Each may have certain advantages and disadvantages which have been evaluated in our laboratory; we use the streptavidin-biotin method. The streptavidin-biotin technique lends itself to the localisation of numerous antigens in a variety of specimens including paraffin sections, cryostat sections, smears, imprints and cytopins. For a specific antigen it can be determined what type of cells produce this substance in normal and neoplastic tissue, levels of the substance produced, and the determination of tumour cell

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

differentiation.

Immunoperoxidase staining involves the use of antibodies and the enzyme peroxidase. Peroxidase is commonly used for several reasons:

1. Its small size will not hinder the binding of antibodies to adjacent sites.
2. It is easily obtainable in highly purified form so that the chance of contamination is minimised.
3. It is very stable, and therefore will remain unchanged during manufacture, storage and application.
4. Only small amounts are present in tissue specimens, and this endogenous peroxidase activity is easily quenched.
5. There is a wide availability of chromogens, which can be acted upon by peroxidase to form a coloured end product that will precipitate at the site of the antigen to be localised.

5.12 Streptavidin-biotin Method

This method is based on the ability of the egg white glycoprotein avidin to immunologically bind four molecules of the vitamin biotin. Three reagents are used. The first is a primary antibody specific for the antigen to be localised. The secondary antibody, capable of binding to that first, is conjugated to biotin. The third reagent is a complex of peroxidase conjugated biotin and

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

avidin. The free sites of the avidin molecule allow binding to the biotin on the secondary antibody. The peroxidase enzyme, and therefore the original antigen, is visualised with an appropriate chromagen.

Even though conjugated antibodies are used in this method, the strong affinity of avidin for biotin gives this method greater sensitivity than other conjugated antibody techniques such as the direct and indirect methods. Excellent results can be achieved on fixed, paraffin embedded specimens.

5.13 Staining Procedure

The details of the exact staining procedure are summarised in Appendix No.6.

5.14 Endogenous Peroxidase Activity (Appendix No. 7)

The substrate chromagen reaction used to visualise peroxidase cannot distinguish between the enzyme immunologically localising the cellular antigen, and similar enzymatic activity present in the specimen before staining. This endogenous peroxidase activity is confined mostly to red and white blood cells.

If it is not removed before adding the marking enzyme, positive staining will be observed that is due not to the specific antigen alone, but also due to peroxidase activity already present in the specimen. Removal of endogenous

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

peroxidase is essential to correct interpretation. This was achieved by quenching the mounted specimens in a solution of 0.5% hydrogen peroxide in methanol.

5.15 Antibodies

The Dako D07 was used for the detection of aberrant p₅₃ because this has been shown to be an effective antibody in comparative studies of six antibodies ¹³, and has been used in our laboratory routinely and by others with good results ¹⁰. Dako M887 was used for bcl₂, SantaCruz Bax N20 for bax alpha, Dako M722 for Ki-67, and Dako A562 for PSA.

5.16 Dilution of antibodies.

Optimally diluted antibodies must be used to achieve meaningful results. The dilution of any one antibody is dependent upon several factors:

- Concentration of specific antibody in the solution as provided by the manufacturer (antibody titre). The more specific antibody molecules per millilitre, the higher the titre and the lower the working concentration.
- A number of substances other than specific antibody are often present in the solution; high levels of contaminating proteins necessitate high dilutions to prevent non-specific background staining.

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

Generally the longer the incubation time, the more dilute the antibody can be used. In this study, antibody incubation times were held constant after experimental determination of optimal time.

Other considerations such as the choice of dilution buffer, use of a humidity chamber, the specimen fixation and processing protocol, will all affect the dilution used. Optimal dilutions for our laboratory conditions were ascertained for each individual antibody.

When interpreting dilution results, two criteria were evaluated: Specific staining and non-specific background staining. The goal was to achieve the greatest intensity of specific staining with the least amount of background interference. The higher the antibody dilution, the lower the background staining due to undesired protein binding.

5.17 Incubation times

Incubation times, antibody dilutions, non-specific background staining and specific staining intensity are all interrelated. The longer the incubation times the higher the antibody dilution. The more dilute the antibody, the lower the non-specific background stain. The lower the non-specific stain, the greater the contrast with the specific stain, making interpretation easier.

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

Incubations with antibody solutions were for 60 minutes to 16 hours. When processing several slides at once, there will be a one or two minute difference between the first slide and the last. To achieve consistency processing of slides was staggered so that all specimens undergo identical incubations. The temperature of incubation will also affect antibody dilutions. The optimal temperature is calculated for each antibody.

Haematoxylin and eosin stains were carried out on all the sections. This determined the presence of carcinoma, PIN, and benign prostate tissue. The areas of tumour and PIN were demarcated and then transferred to the computer-generated images.

5.19 Antigen retrieval (micro waving) (Appendix No.9)

The use of formalin fixed paraffin blocks was unreliable before the advent of antigen retrieval techniques such as microwave irradiation or a pressure cooking to reverse some of the effects of fixation (un-fixation). Studies indicate that antigen retrieval provides an extremely successful method for the demonstration of nuclear antigens with formalin fixed, paraffin embedded tissues^{351-355;383;384}, even after fixation in formalin for as long as 7 days³⁸³. The use of an antigen retrieval system provided excellent results with minimal background staining, and high contrast between positively stained and negative

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

cells. Studies for p₅₃ using single strand conformational polymorphism analysis of RNA/PCR products and sequencing have shown that using microwave irradiation as an antigen retrieval system detects 93% of mutations between exons 5-8⁴¹³.

For the immunohistochemical studies the slides were microwaved or placed in a pressure cooker as part of an antigen retrieval system. The microwave incubation times had been predetermined (PSA = 5mins, p₅₃ = 10 minutes, and bcl₂ = 20 minutes). For Ki-67 pressure-cooking has been used successfully in our laboratory as an antigen retrieval system, the time used was 90seconds. For bax no antigen retrieval is required.

5.19 Controls

Certain procedural and reagent controls are essential for the validation of the results of immunoperoxidase staining. Most of these controls centre on the primary antibody to ascertain that positive staining is the result of specific binding to the antigen. After determining the optimal dilution of a new primary antibody, several known positive and negative specimens were stained to certify the antibody's specificity.

Positive control: A specimen, processed identically to the unknown that contains the antigen in question.

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

Two negative controls: One specimen processed in an identical way except that non-immune serum is used instead of the primary antibody. This serves as a specimen blank since any staining observed is due not to antibody localisation of the antigen but to non-specific protein binding, endogenous peroxidase activity, or non-specific binding of the other antibody reagents.

The second specimen processed in an identical way except that non-immune serum is used in place of both primary and secondary antibodies. This serves to exclude staining that may be due to interaction between the chromogen used 3, 3 Diaminobenzidine Tetrachloride (DAB) and processed tissue (DAB negative control).

All controls underwent the same fixation and processing schedule as the unknown specimen and were processed along with the unknown to assure the accuracy of the results.

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

Antigen	Clone	Supplier	Antigen retrieval.	Ab Dilution	Incubation time.	Controls
P ₅₃	D07	Dako	10 mins microwave	1:50	60 mins @ room temp.	Vulval cancer
Bcl ₂	M887	Dako	20mins microwave	1:25	60 mins @ room temp.	Tonsil
Bax-alpha	Bax N20	SantaCruz	0	1:800	16hrs @4°C	Breast
Ki-67	M722	Dako	90s in 2l Citrate buffer in a pressure cooker.	1:500	60 mins @ room temp.	Tonsil
PSA	A562	Dako	5 mins microwave	1:800	60 mins @ room temp.	Benign prostate

Table23: A summary of the antibodies used the incubation times, and the controls.

5.20 Principles of Interpretation of immunostaining

The most difficult aspect of immuno-staining is in the correct evaluation of the finished product. The amount of chromagen precipitated, and therefore the intensity of the reaction has been found to be proportional to the amount of antigen present. Not all cells contain the same amount of antigen, and therefore stain with varying intensity. Non-antigen containing cells and cell components are contrasted by a counter stain (methylene blue) which makes distinguishing the two easier.

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

Specific staining is localised in cells, non-specific background staining is usually found in collagen and connective tissue. Interpretation is basically a comparison of the specific and non-specific staining pattern and intensity, if any, of the specimen with that of the control. The source of the tissue, antigen characteristics and the procedure used must be taken into account for proper evaluation.

Several artefacts may be found in specimens, which may produce staining and interfere with correct interpretation. They are mostly not confined within cells, but are spread randomly across the specimen. Any staining that occurred on the very edge or along knife marks is often non-specific and can be ignored. Positive staining of necrotic crushed or cells that have undergone autolysis and haemorrhagic tissue, is always regarded as non-specific. Only the staining pattern of viable cells is considered for interpretation.

5.21 Non-specific Background Staining

Positive staining of a specimen that is not a result of antigen antibody binding is termed non-specific background staining. The most common cause is attachment of protein to highly charged collagen and connective tissue elements of the specimen. Antibodies are proteins. If the first protein solution applied to the tissue is the primary antibody, it can be non-specifically adsorbed to these charged sites. The secondary antibody can still bind to the

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

primary and the peroxidase colour reaction will occur. Positive staining of these sites is due not to localisation of the tissue antigen, but to non-specific antibody attachment to collagen and connective tissue, and neurones and ganglion cells for bcl₂.

Other causes of non-specific background staining such as inappropriate antibody dilutions incomplete removal of paraffin, improper rinsing of slides and incorrect substrate incubation can be minimised by careful and accurate technique.

5.22 Scoring of immunostaining

All samples were scored firstly as positive or negative staining, secondly by assessing the intensity of staining, thirdly by assessing the proportion of cells stained. In carcinoma and PIN this grading was confined to only regions of interest within the specimen. The areas of PIN and cancer within each specimen were initially identified with H&E.

Cases were considered positive if any of the tumour cells manifested positivity. Intensity was scored on a four point scale: no staining = 0; weak staining = 1; moderate staining = 2 and intense staining = 3. Assessment of intensity of staining was always referenced to a standard positive control, stained at the time and under the same conditions as the test case.

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

The proportions of cells staining, for KI-67 in particular, were scored as a percentage of cells staining positive against total number of cells per high power microscopic field (x 400). All histopathological assessment was confirmed two histopathologists, Dr Michael Jarmulowicz, Consultant Histopathologist at the Royal Free Hospital, and Dr Tara Walker, Consultant Histopathologist at Queen Alexandra Hospital, Portsmouth. There was agreement in most (>90%) of cases with regard to the intensity and pattern of staining between the two pathologists. In the sections in which the scores differed, we agreed a common result. The percentage of positive cells within a section was estimated by 'eye-brain' co-ordination.

5.23 Background staining

Background staining was minimal or absent in most sections and did not therefore interfere with assessment. The p₅₃, bcl₂ and MIB-1 antibodies achieved particularly specific staining with an absence of background staining. The bax had some diffuse background staining.

5.24 Staining patterns

The criterion for a positive reaction was nuclear staining for p₅₃ and Ki-67 and cytoplasmic staining for bcl₂, bax alpha.

5.30 Statistics.

The data obtained from this study was tabulated in an excel spreadsheet. The raw data is presented in Appendix 10 and 11. The data was assumed not to follow a normal Gaussian pattern, and therefore treated as a non-parametric data set. Analysis designed for parametric data can not be applied to non-parametric data, but non-parametric analysis can be applied to parametric and non-parametric data to give valid results. Therefore the use of non-parametric tests would maintain the statistical stringency of the analysis.

The statistical package for social science (SPSS) 10.1 was used for data analysis. The analysis of variance (ANOVA) was used calculate the relationship between tumour, PIN, and benign tissue volume and serum PSA. The data obtained from the staining of the slides was also assumed to be non-parametric, so the Mann-Whitney U test was used to compare the relationship between the slides stained.

Assistance and guidance about statistical method was sought from Richard Morris (Department of Statistics, The Royal Free Hospital), David St George (Department of Statistics, The Royal Free Hospital) and Roger A'Hern (Department of Statistics, The Royal Marsden Hospital).

Chapter 6: Results

The results obtained during this piece of work have been divided into two sections. The first set of data was obtained from the morphometric analysis of 76 radical prostatectomy specimens. This was carried out prospectively. In summary the whole radical prostatectomy was fixed in 10% buffered formaldehyde for 72 hours. The specimen is then weighed and inked, slices are prepared at intervals of 3.46mm, the slices are scanned and the image obtained is used as a template. The slices are then cut into four and placed in a standard cassette. Slides are prepared from the cassettes and haemotoxylin and eosin stain is used to determine the areas of interest. This is transcribed onto the initial template. A shrinkage fraction for the template is 1.03, and for the areas of cancer, and PIN is 1.22.

The data is presented in graphical form wherever possible. The raw data is available in appendix 10.

6.1: Results 1: Morphometric Studies.




Fig 18: The age of the patients.

This graph represents the age distribution of the 76 patients whose prostatectomy specimens were used to investigate the relationship between serum PSA, cancer volume and PIN volume. The mean age of the patients at the time of surgery is 63.07 ± 7.4 (SD), with a range of 46 to 78 years.

Prostatic Intraepithelial Neoplasia and Prostate Cancer.




Fig 19: Stage of radical prostatectomy specimens.

This graph represents the stage distribution of the radical prostatectomy specimens obtained from the 76 patients whose prostatectomy specimens were used to investigate the relationship between serum PSA, cancer volume and PIN volume. The stage of specimens was assigned using the International Union against Cancer (UICC) TNM Classification of malignant tumours, 6th edition, published in 2002.

Prostatic Intraepithelial Neoplasia and Prostate Cancer.




Fig 20: Distribution of weight of gland.

The specimens were obtained fresh from the operating theatre. All loose tissue was removed. The specimens were then directly taken fresh to the histopathology department and weighed. The mean fresh weight is 61.3 g \pm 21.6 (SD) with a range of 26 to 173 g.

Prostatic Intraepithelial Neoplasia and Prostate Cancer.




Fig 21: Distribution of volume of glands.

Prostatectomy specimens are an irregular shape and therefore linear instruments can not be used to calculate their dimensions. The volume of the prostatectomy specimen was calculated by obtaining the fresh specimen, removing any loose tissue, and fixing it in formaldehyde. The slices were cut at 3.5mm intervals. They were then scanned using Paint Shop Pro (version 6.0). The scanned images were transferred using Sigma Scan Pro (version 6.0 by Jandel Scientific, UK). This made the images into a format accessible to image analysis software. The software used was Kontron KS400.

The regions of interest could then be identified and measured. In order to measure the volume of the gland, the area of all the slices except the apical slices were measured. This was multiplied by the thickness of the slices and the shrinkage fraction, to obtain the volume of the entire gland. There is an

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

assumption made that the prostate specimen is of uniform density. The mean calculated volume of the prostatectomy specimens is 50.9 mls \pm 20.6 (SD), with a range of 19.4 to 155.5 mls.

Prostatic Intraepithelial Neoplasia and Prostate Cancer.




Fig 22: Correlation between calculated volume and weight of gland ($R_{sq} = 0.985$).

The data obtained from the weight and the volume of the specimens was used to calculate a correlation coefficient. SPSS software was used and analysis of variance (ANOVA) was used to calculate the correlation coefficient.

There is good correlation between the volume of the specimens and the weight of the gland. This is reassuring and further supports using the scanned images to accurately calculate the volume of the gland. The majority of human tissue is composed of water so the density obtained from this calculation is close to 1g/ml.

The line does not cross the 0 on the x and y-axis simultaneously. This could be due to the volume of the apical slices not being included in the volume calculations.

Prostatic Intraepithelial Neoplasia and Prostate Cancer.




Fig 23: Distribution of Gleason score

The Gleason score is the sum of the two most common Gleason grades. The Gleason grade is determined by the architecture of the prostate cancer^{131;250}. The Gleason score will affect the management options available to the patient, for example patients with a Gleason score of seven or above are rarely offered surveillance. The Gleason score also has prognostic significance, in this series the mean Gleason score is 6.2 \pm 1.2 (SD) with a range of 2 to 9. This is in keeping with the results from other published series.

Prostatic Intraepithelial Neoplasia and Prostate Cancer.




Fig 24: The distribution of PSA (ng/ml) in the 76 patients studied for morphometric analysis.

The blood sample for the PSA was taken the night before the operation. The hybertech assay was used for PSA. The range of PSA was 3 to 45 ng/ml, with a mean of 9.1 ± 5.9 (SD), and a median of 5.5 ng/ml.

Prostatic Intraepithelial Neoplasia and Prostate Cancer.




Fig 25: The distribution of cancer volume.

The fresh prostate was fixed for 72 hours in formaldehyde. The specimen was then sliced at 3.5-mm intervals; the slices were then scanned using Scan Pro (version 6.0). Thus preparing a template. The prostate slices were placed in cassettes and embedded. Slides were prepared and stained with haematoxylin and eosin. The areas of cancer were identified. This was transcribed on to the template in red. The scanned images were transferred using Sigma Scan Pro (version 6.0 by Jandel Scientific, UK). This made the images into a format accessible to image analysis software. The software used was Kontron KS400. The regions of prostate cancer were identified and measured. In order to measure the volume of the prostate cancer, the area of all the areas of cancer except the apical slices were measured. This was multiplied by the thickness of the slices and the shrinkage fraction, to obtain the volume of the cancer. The range of the prostate cancer was .09 to 15 mls with a mean of 2.5 ± 2.7 (SD).




Fig 26: The distribution of PIN volume in the 76 prostates studied.

The fresh prostate was fixed for 72 hours in formaldehyde. The specimen was then sliced at 3.5-mm intervals; the slices were then scanned using Scan Pro (version 6.0). Thus preparing a template. The prostate slices were placed in cassettes and embedded. Slides were prepared and stained with haematoxylin and eosin. The areas of PIN were identified. This was transcribed on to the template in blue. The scanned images were transferred using Sigma Scan Pro (version 6.0 by Jandel Scientific, UK). This made the images into a format accessible to image analysis software. The software used was Kontron KS400. The regions of PIN were identified and measured. In order to measure the volume of PIN, the area of all the areas of PIN except the apical slices were measured. This was multiplied by the thickness of the slices and the shrinkage fraction, to obtain the total volume of PIN.

The range of PIN measured was 0 to 2.7 mls with a mean of 0.53 ± 0.6 (SD) mls. The area of the prostate taken up by benign tissue was calculated by

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

subtracting the volume occupied by PIN and prostate cancer from the calculated volume of the whole gland. The range of volume for benign gland was 47.5 mls with a range of 13.3 – 153.4 mls.

Prostatic Intraepithelial Neoplasia and Prostate Cancer.




Fig 27: Relationship between serum PSA and benign volume of prostate.

The volumes were calculated using the scanned images, multiplying by the slice thickness and making a correction for the shrinkage fraction. The benign volume of the gland was calculated by subtracting both the volume of PIN and cancer from the total volume of the gland i.e. total prostate volume – (volume of PIN + volume of prostate cancer). The volume of the benign tissue was plotted against the serum PSA (ng/ml) using SPSS10.1 software. The scatter chart generated correlation (spearman rho=0.24). The correlation between the serum PSA and the volume of benign prostate is significant ($p=0.05$).

Prostatic Intraepithelial Neoplasia and Prostate Cancer.




Fig 28: Relationship between serum PSA and volume of prostate cancer.

The volume of cancer was calculated from the scanned images; the area of the cancer was measured from the scanned images and multiplied by the thickness of the slices. The volume so calculated for each specimen was multiplied by a correction factor for shrinkage of 1.22. The volume of prostate cancer was plotted against the serum PSA. There is a correlation between the serum PSA and the volume of prostate cancer, spearman rho = 0.36, this is highly significant with a p value of 0.01).

Prostatic Intraepithelial Neoplasia and Prostate Cancer.




Fig 29: Relationship between serum PSA and volume of prostatic intraepithelial neoplasia.

The volume of PIN was calculated from the scanned images using Kontron KS400 software to determine the area of the regions of interest. This was multiplied by the slice thickness (3.46mm) and the shrinkage factor (1.22). The volume of PIN was plotted against serum PSA (in ng/ml). There was no correlation between the volume of PIN and serum PSA. Therefore, PIN using the model developed in this piece of work does not appear to contribute significantly to PSA.

Prostatic Intraepithelial Neoplasia and Prostate Cancer.




Fig 30: Relationship between PIN volume and cancer volume.

The volume of PIN was compared to the volume of cancer. Some authors have suggested there is a strong and significant inverse relationship between the volume of PIN and cancer. Hypothesising that this may be further evidence to support the view that PIN develops into prostate cancer⁹⁹. A weak inverse relationship was detected between the volume of PIN and prostate cancer (correlation coefficient = -0.155), but this does not reach significance ($p = 0.183$). Therefore this data does not support the work of de la Torre et al⁹⁹.

Prostatic Intraepithelial Neoplasia and Prostate Cancer.




Fig 31: Relationship between PIN volume versus total calculated volume.

The total volume of the gland was calculated using the prostate maps, and this was compared with the volume of PIN. There was a significant correlation with a coefficient of 0.4($p=0.03$). This may imply that both conditions respond to common stimuli, e.g. testosterone.

Prostatic Intraepithelial Neoplasia and Prostate Cancer.




Fig32: Cancer volume versus reciprocal of total volume.

The volume of cancer was compared to the volume of the gland measured from the templates. The volume of cancer is inversely related to the total volume of the gland with a correlation coefficient of 0.3, and a p value of 0.01.

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

This may be due to the fact that in the majority of patients in this data set, the trigger for a biopsy is PSA, and if the PSA reaches a critical threshold then the patient is offered a biopsy. Since both benign and malignant tissue can produce PSA, therefore a patient may be offered a biopsy if he has a large benign gland with minimal tumour - the majority of the PSA being generated by benign tissue; or a small gland with a high volume of tumour - the rise in serum PSA being due to prostate cancer.

Prostatic Intraepithelial Neoplasia and Prostate Cancer.




Fig 33: Cancer volume versus Gleason score.

The Gleason score is a predictor of outcome in patients with prostate cancer^{131:250}. Tumour volume correlates with stage and is also a prognostic factor. Therefore there may be a relationship between Gleason score and cancer volume. We investigated this. No relationship could be demonstrated between tumour volume and Gleason score.

Prostatic Intraepithelial Neoplasia and Prostate Cancer.




Fig 34: PIN volume versus Gleason score.

The volume of PIN was calculated using the templates, and this was compared to the Gleason score of the prostate specimens. There is no correlation between these two variables.

6.2: Results 2:

Immuno-histochemical determination of p₅₃, bcl₂, bax-alpha, and PSA in benign prostate tissue, PIN, and prostate cancer.

Areas of cancer, and PIN were identified on H&E sections of microscope slides obtained from radical prostatectomy specimens from 92 hormone naïve patients, and areas of cancer were identified from radical prostatectomy specimens obtained from 14 patients treated with three months of androgen ablation. It is difficult to identify areas of PIN after androgen ablation. Low power views were used to identify the areas of interest.

The mean age of the patients was 64.1 (range 46-71) years, mean PSA 9.6 (range 1.9-32.40) ng/ml, 77 patients had organ and capsule confined disease, and 29 patients had extracapsular disease. The mean Gleason score was 6.1 (range 2 to 9). All patients were fully staged and were NOM0.

Specimens obtained from 19 patients who had a TURP for benign disease were used as controls

For p₅₃ the D07 antibody was used, and nuclear positivity was sought, the presence of positive staining was regarded as positive. The presence of p₅₃ was sufficient for a specimen to be labelled as positive regardless of the percent of

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

cells being stained positive^{312;421}. In addition the intensity of the staining was compared (Table 25). There was no significant difference in the intensity of staining between the groups studied.

For Ki-67 staining, the MIB-1 antibody was used, in order to obtain a proliferation index. 1000 cells were examined at high power and the proliferation index was the percentage of cells staining tumours positively with MIB-1. Brown granular nuclear staining was considered positive for Ki-67.

For bcl₂ and bax cytoplasmic staining was used to determine a positive result. For bcl₂ the Dako M887 antibody was used, and for bax the Santa Cruz bax N20 antibody was used. The presence of positive staining cytoplasmic cells was regarded as positive; in addition for bcl₂ the proportion of positive staining cells in the region of interest was noted. The intensity was also measured for both of the antibodies.

The crude data is available in Appendix 11.

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

Table 24

Percentage of P53 positive cells per slide.

	N	Minimum	Maximum	Mean	Std. Deviation
Benign	19	0	0	.00	.000
PIN	91	0	100	3.46	15.859
Ca	92	0	100	6.96	23.334
LHRH Treated	12	0	100	25.83	39.648

Table 25

Intensity of Staining of P53 Staining.

	N	Minimum	Maximum	Mean	Std. Deviation
Benign	19	0	0	.00	.000
PIN	92	0	3	.10	.446
CA	92	0	2	.14	.459
LHRH	14	0	3	.43	.852

Table 26

Percentage of Bcl2 positive staining cells

	N	Minimum	Maximum	Mean	Std. Deviation
Benign	19	0	0	.00	.000
PIN	92	0	100	47.93	36.736
Ca	92	0	100	15.98	29.800
LHRH	14	0	100	55.71	46.029

Table 27

Intensity of Staining of BCL-2 Staining.

	N	Minimum	Maximum	Mean	Std. Deviation
Benign	19	0	0	.00	.000
PIN	92	0	5	1.39	.889
CA	92	0	2	.35	.563
LHRH	14	0	3	1.00	.961

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

Table 28

Percentage of Ki 67 positive staining cells

	N	Minimum	Maximum	Mean	Std. Deviation
Benign	19	0	0	.00	.000
PIN	92	0	40	4.50	6.010
CA	92	0	35	5.68	6.304
LHRH	14	1	25	2.79	6.399

Table 29

Intensity of Staining of Bax Staining.

	N	Minimum	Maximum	Mean	Std. Deviation
Benign	19	0	2	1.38	.650
PIN	84	0	3	1.80	.741
CA	84	0	3	1.88	.735
LHRH	14	.00	2.00	1.40	.986

Table 30

Intensity of Staining of PSA Staining.

	N	Minimum	Maximum	Mean	Std. Deviation
Benign	19	3	3	3.00	.000
PIN	92	1	3	2.59	.577
CA	92	0	2	.14	.459

		Total Number	No. +	Mean no of +ve cells +/- SD	Staining intensity +/- SD
P53	BPH	19	0	0	0
	PIN	92	5	3.46 +/- 15.86	0.1 +/- 0.45
	CaP	92	9	6.96 +/- 23.33	0.14 +/- 0.46
	LHRH	14	4	25.83 +/- 39.65	0.43 +/- 0.85
Bcl2	BPH	19	0	0	0
	PIN	92	75	47.93 +/- 36.74	1.39 +/- 0.89
	CaP	92	27	15.98 +/- 29.8	0.35 +/- 0.56
	LHRH	14	9	55.71 +/- 46.03	1 +/- 0.96
Baxalpha	BPH	13	13	100	1.38 +/- 0.65
	PIN	84	78	100	1.8 +/- 0.74
	CaP	84	80	100	1.88 +/- 0.74
	LHRH	10	10	100	1.4 +/- 0.96
KI-67	BPH	19	0	0	
	PIN	92	91	4.5 +/- 6.01	
	CaP	92	92	5.68 +/- 6.3	
	LHRH	14	14	2.79 +/- 6.4	
PSA	BPH	19	19	100	3 +/- 0
	PIN	92	92	100	2.6 +/- 0.56
	CaP	92	92	100	2.15 +/- 0.76

Table 31: Summary of results obtained by immunohistochemistry

Comparisons	Number of + specimens					Number of cells			Staining intensity			
	P ₅₃	Bcl 2	Bax	Ki-67	PSA	P ₅₃	Bcl2	Ki-67	P ₅₃	Bcl2	Bax alpha	PSA
PIN and BPH	NS	<.001	NS	<.001	NS	0.71	<0.001	<0.001	0.71	<0.001	0.06	0.0116
CaP and PIN	NS	<.001	NS	NS	NS	0.61	<0.001	0.16	0.62	<0.001	0.57	0.0002
CaP and BPH	NS	<.006	NS	<.001	NS	0.51	0.04	<0.001	0.51	0.04	0.03	<0.001
LHRH and CaP	.07	0.015	NS	NS	NS	0.19	0.006	0.006	0.27	0.016	0.04	

Table 32: Comparison of the benign, PIN, and cancer using the Mann Whitney U Test and the difference of the means for the number of cells and staining intensity. P-values in bold denote negative differences, i.e. the second factor is higher than the first. Otherwise the first factor is higher.

Chapter 7: Discussion

The volume of the prostate glands was calculated using digital images of the slices, which were corrected for the shrinkage factor. This calculated volume was compared to the weight of the gland. There was good correlation (correlation coefficient of 0.967). This would support this technique as a valid method of calculating prostate volumes.

7.1 Relationship between total volume of gland, volume of PIN, benign volume of gland, and volume of cancer.

Tumour volume as assessed in total prostatectomy specimen correlates with prognosis, especially tumour volumes greater than 4.0cm^3 even in patients who have relapsed^{93:367}. When comparing the tumour volume as a marker for recurrence, some have suggested that it is a better predictor of recurrence after radical prostatectomy than either Gleason score or preoperative PSA²⁹³.

Tumour differentiation decreases and DNA patterns deteriorate as the tumour volume increases. Some groups have shown that tumour volume correlates with histological grade^{301:365} and although it did predict progression, in a multivariate analysis it was not an independent predictor of recurrence³⁰¹. In

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

our series tumour volume did not correlate with Gleason grade.

In this study the volume of cancer was related to the stage ($p < 0.001$). The cancer volume also had an inverse correlation with the benign and the total volume of the gland ($p < .001$). This may be due to the fact that to trigger a biopsy and so the possible diagnosis of prostate cancer the patient needs to reach a threshold PSA. PSA can either be produced by the benign tissue, i.e. a patient will have a large gland with a small volume of cancer; or by malignant tissue and so a patient will have a small gland with a large volume of cancer sufficient to produce the PSA which leads to serum levels that trigger a biopsy.

Additionally this work has demonstrated that the size of the gland correlates with the volume of PIN. Both BPH and PIN increase with age and require androgens. Therefore, the stimulus for these lesions is probably identical, with PIN developing in the peripheral zone and BPH developing in the transition zone. If we accept that PIN is a premalignant lesion, then large glands have a corresponding increased volume of PIN, which would predispose to cancer. Therefore gland volume is a predictive factor for prostate cancer. The European Randomised Study of Screening for Prostate Cancer (ERSPC) screened a total of 8,621 men. Using a logistic regression model this group determined the most important predictors for prostate cancer was PSA, followed by prostate volume, DRE, and TRUS findings, in that order²⁰⁵. Thus

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

a large gland irrespective of the PSA may be indicative of prostate cancer, due to the increased volume of PIN.

No significant correlation was detected between the volume of PIN and the volume of cancer in this study. This is in contrast to other studies. De la Torre et al did demonstrate an inverse relationship between volume of PIN and the volume of cancer. Implying that as PIN transforms into prostate cancer, the volume of PIN decreases as the volume of cancer increases, however they did not differentiate between high and low grade PIN⁹⁹. Alexander et al demonstrated a significant correlation between PIN volume and the weight of the gland and the cancer volume⁶.

7.2 Relationship between volume of PIN, volume of benign tissue, volume of cancer and serum PSA.

In this study it has been shown using a regression model that the serum PSA significantly correlates with the volume of the gland and the volume of prostate cancer. In a multivariate analysis this correlation appears to be independent. Using SPSS ANOVA software, we have calculated one gm of benign prostate tissue correlates with 0.07ng/ml of serum PSA, and one gm of malignant tissue to 0.8 ng/ml of serum PSA.

Stamey et al in their seminal paper calculated that the serum PSA was related

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

to tumour volume, and that prostate cancer produced more PSA than benign tissue³⁶⁶. In the cohort of patients with prostate cancer no significant correlation was calculated between the volume of benign tissue and the serum PSA. In seven separate patients undergoing a Millen's retro pubic prostatectomy for benign disease they calculated that on average one gram of benign tissue corresponded to 0.29 ng/ml of PSA in the serum³⁶⁶. This group further studied the relationship between cancer and serum PSA and concluded that prostate cancer elevates serum PSA twelve fold compared to BPH and so one gram of cancer increases serum PSA by 3.5ng/ml³⁶⁵. However, of the 102 radical prostatectomy specimens used to calculate this figure only 48 were completely confined to the prostate, and therefore the total tumour volume was only measured with any degree of accuracy in less than half of this cohort. Stamey also detected a correlation between serum PSA and prostate weight, but not with the volume of benign tissue³⁶⁵. Partin et al studied the relationship between tumour volume, benign gland volume and serum PSA in 350 radical prostatectomy specimens and 72 men with benign prostatic hyperplasia. A correlation was found between the cancer volume and PSA, no correlation was detected between serum PSA and the volume of benign tissue³⁰⁰. Later, the same group in a series of 47 radical prostatectomy specimens detected a significant correlation between serum PSA and the weight of the gland and volume of malignant tissue⁴¹². This has been confirmed in this study and by others⁶. Ronnett et al demonstrated that in small volume tumours i.e. 0.5cc or

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

less, the weight of the gland correlates with the PSA, and the tumour volume does not³²⁶.

In a series of 159 cases with a clinically benign diagnosis and using TRUS to calculate the volume of the gland, and of the transition zone. Aarnich et al calculated that one-gram of the transition zone contributed to 0.150ng/ml of serum PSA and the remainder of the prostate contributed 0.077ng/ml per gram. The mean contribution for one gram of prostate was 0.092ng/ml². An earlier study of 472 men calculated a value of 0.72 ng/ml per gram of benign prostate tissue⁸⁶. The relationship between prostate volume and free PSA has also been studied. Using a multiple linear regression analysis in a model containing both free and total PSA values obtained from 681 men, free PSA was the only significant predictive variable of prostate volume²⁷¹. Studies have examined not only the effect of prostate volume on serum PSA but also age, it has been shown that serum PSA increases with age^{86:102:325}.

In this study we have demonstrated that there is no relationship between the volume of PIN and serum PSA.

Brawer postulated that for PSA to enter the serum it must traverse the basal cell layer, epithelial basement membrane, stroma, capillary basement membrane and the endothelial cells. In PIN there maybe disruption in the basal

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

cell layer and so it was postulated that serum PSA might be increased in patients with PIN⁵⁰. In a series of 81 patients, treated for bladder outlet obstruction by retro pubic Millen's prostatectomy and TURP, which included 25 men with PIN and 26 with BPH. The median serum PSA levels were significantly higher in the group with PIN (4.0 ng/ml) compared to the men with BPH (2.1ng/ml)⁵³. Using needle biopsies Morote et al identified 321 men with benign prostatic hyperplasia, 17 had PIN present and 304 had no PIN, there was a small but insignificantly higher total PSA in patients with benign tissue and associated PIN than those with no PIN (7.7ng/ml vs. 7.2ng/ml, $p>0.05$)²⁷⁰. Fowler et al carried out TRUS guided biopsies on 1961 consecutive men with suspected prostate cancer as part of a screening protocol. 93 patients had PIN diagnosed in the absence of cancer, and 957 patients had a benign diagnosis without PIN. The mean total PSA was 5.6 and 4.6 ng/ml, and the percentage free was 20.6 and 21.5 respectively. The difference in the percentage of free PSA was not significant, but the total PSA was significantly higher in the group with PIN ($p=0.02$)¹²⁰. Bostwick et al also detected an increased PSA in patients with PIN diagnosed on needle biopsy compared to a benign diagnosis⁴⁸

A study of 15 patients with BPH, 46 with PIN, and 16 with localised prostate cancer, the total mean PSA was significantly different between the three groups i.e. 2.78ng/ml for BPH, 4.89ng/ml for low grade PIN, 8.21ng/ml for

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

high grade PIN, and 12.70ng/ml for cancer. The mean free to total ratio was lower in patients with low grade PIN (25%) or high grade PIN (21%) compared to cancer (14%) but this did not reach significance¹⁹⁸. Minardi et al carried out a retrospective study of 87 patients, 30 patients with BPH, and 24 with high grade PIN, 8 with low grade PIN and 25 with carcinoma of the prostate²⁵⁸. The mean free to total ratio was 24.2% in those with BPH, 17.6% in the high grade PIN, and 13% in the prostate cancer group. These differences were significant. The mean values for the total PSA i.e. 6.17 ng/ml, 6.89 ng/ml, and 8.02 g/ml respectively, were not statistically significant²⁵⁸. Ramos et al similarly followed up 48 men who were diagnosed with PIN without prostate cancer. This cohort was compared to a group diagnosed with prostate cancer and a group with benign prostatic hyperplasia in the absence of PIN. The percentage free PSA was significantly lower in patients with prostate cancer, but there was no difference in the percentage free PSA in the men with PIN and those with BPH (20.8+/- 7.1% and 20.1+/-7.3%, respectively)³¹⁶. In a study of 81 consecutive patients undergoing radical prostatectomy, Morote et al found no significant difference in the total serum PSA between the specimens that contained PIN and cancer compared to those with cancer alone. But the free to total serum PSA was significantly lower in the group with PIN, but in a multivariate analysis PIN did not contribute to free serum PSA²⁶⁹. In the large Austrian screening study of 1474 patients, the mean total PSA of patients with BPH, PIN and cancer was 6.0, 5.9, and 8.7ng/ml, with the

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

percentage free ratio being 21.9, 15.0 and 12.1% respectively. With regard to total PSA there was no significant difference between BPH and PIN, but with regard to the percentage free PSA there was a significant difference between the BPH group and all the other groups, and the PIN and cancer groups¹⁷⁰. This study could not identify a clear cut off between the groups due to a significant overlap¹⁷⁰.

Alexander et al have shown that PIN volume does correlate with serum PSA concentration⁶. This group studied 194 whole mounted radical prostatectomy specimens and used a grid counting technique to determine the volume of PIN, and prostate cancer. They, however, concluded that the relationship between serum PSA and PIN was an artefact, in a multiple regression analysis adjusting for cancer volume, gland weight, Gleason score, and extraprostatic extension, log PIN volume did not contribute to log serum PSA⁶.

Ronnet et al using 65 whole mount radical prostatectomy specimens weighing less than 65 gm and with small tumour volumes of <0.5mls, found no correlation between PSA and tumour and PIN volume, but did find a correlation between serum PSA and total gland weight³²⁶.

Prostatic Intraepithelial Neoplasia and Prostate Cancer.




Fig35: PIN stained with PSA x 40




Fig36: Cancer stained with PSA x 40

7.3 P₅₃ in benign, PIN and malignant prostate tissue.

Expression of p₅₃ increased in a stepwise manner from benign to intra epithelial neoplasia to malignancy. 0/19 benign, 5/92 PIN, and 9/92 cancer specimens were positive for p₅₃. This increase was not significant. Loss of p₅₃ was thought to be a common event in localised prostate cancer²⁷⁴, however more recent studies have not supported this view²⁷³. In only two out of the five PIN specimens expressing p₅₃ was there also concurrent expression of p₅₃ in the cancer specimens implying aberrant expression of p₅₃ is not required for progression. Furthermore sequencing studies of aberrant p₅₃ in both PIN and cancer in the same specimen have revealed that the sites of the point mutations are not identical and therefore PIN and prostate cancer may arise from a different clone with in the same specimen^{377:426}. In addition we have shown that progression to cancer from PIN can occur in spite of the loss of aberrant p₅₃. Loss of p₅₃ is probably a late change in prostate cancer³⁶⁹, and not related to the early genesis of this disease.

Androgen ablation led to proportionally increased expression of p₅₃; this however did not reach significance (p=0.07). There may be a number of explanations for the increase in p₅₃. Androgen ablation leads to the withdrawal of a necessary stimulus for the prostate, the cells of the gland involute, and a p₅₃ dependant pathway may cause this. In addition increased bcl₂ inhibits the nuclear translocation of p₅₃²¹ and so may increase the amount

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

of free wild type p₅₃ antigenically available for the immunohistochemistry antibody to react with.

Increased p₅₃ after androgen ablation may also be a predictor of relapse. An analysis of 39 patients, who had received neoadjuvant LHRH treatment prior to definitive surgery, demonstrated 21 patients that were positive for p₅₃. 90% (19 patients) of these patients relapsed. Therefore the increased expression of p₅₃ associated with androgen ablation may predict relapse³¹².

Prostatic Intraepithelial Neoplasia and Prostate Cancer.




Fig37: PIN stained with P₅₃ x 40



Fig38: Cancer stained with P₅₃ x 40

7.4 Bcl₂ in benign, PIN and malignant prostate tissue.

Bcl₂ is always present in the stem cells of the prostate; therefore we only measured the expression of bcl₂ the luminal cells. The expression of bcl₂ was significantly different between the three groups studied. The expression of bcl₂ in the prostate cancer was comparable with the results of others^{20:74:238:239}. It was greatest in the PIN slides, and least in the benign tissue. In 22 specimens bcl₂ staining was detected concomitantly in PIN and cancer. Bcl₂ infers a degree of immortality and resistance to the cells from cytotoxic treatment, if PIN is a pre malignant lesion and not an epi phenomenon, this would imply that PIN luminal cells have a survival advantage. Others also share this view^{41:187}.

Expression of bcl₂ allows cells to remain viable despite androgen ablation²². In this study androgen ablation led to an up regulation of bcl₂ and therefore it may be postulated that androgen ablation selects out clones with a survival advantage²⁴⁶. Others have also supported this finding³⁹⁴. Up regulation of bcl₂ may also play a role in the development of radio resistant clones³²⁸ and increased expression of VEGF resulting in higher microvessel density¹¹⁶.

An interrelationship exists between p₅₃ and bcl₂. Up regulation of bcl₂ can inhibit the action of the pro apoptotic p₅₃²¹ and conversely p₅₃ can up regulate bax and down regulates bcl₂²⁶¹.

Prostatic Intraepithelial Neoplasia and Prostate Cancer.




Fig 39: Bcl₂ positive staining PIN x 40.




Fig 40: Bcl₂ positive staining cancer x 20.

7.5 Bax in benign, PIN and malignant prostate tissue.

A predetermined set point is thought to exist in cells for the ratio of bcl₂ and bax, and this is thought to play a critical role in the response of malignant cells to therapies, which lead to apoptosis²⁹². Bax inactivation abolishes induction of apoptosis by chemopreventive drugs¹⁷⁸. The expression of bax was almost uniform in all the groups. However, the staining intensity increased significantly from BPH to cancer and there was inactivation of bax with androgen ablation. This would imply up regulation of bax from benign to malignant tissue, so cancer cells are selectively more susceptible to chemotherapeutic/hormonal intervention. When comparing the specimens, which have been treated with androgen ablation there is a decrease in staining intensity, this may be because the susceptible cells have undergone apoptosis and only the resistant clones are left. The ratio of bax to bcl₂ is important and in the group treated with androgen ablation there is an increase in the expression of bcl₂, and so further adding weight to the argument that in this group androgen ablation is selecting resistant clones. Bax inactivation is important in human carcinogenesis, and contributes to tumour progression. Although the number of positive staining specimens does not show a significant change, but the staining intensity does, and therefore a quantitative method for determining the expression of bax is required to confirm its importance. If one measures expression of bax exclusively in terms of the number of specimens staining positive, then its loss does not appear to be

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

important in the initiation of prostate cancer. Others share this view^{187; 204}.



Fig41: Bax positive staining PIN x 40.




Fig 42: Bax positive staining cancer x 40.

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

7.6 Ki-67 in benign, PIN and malignant prostate tissue.

In this study there is a significant increase in the expression of Ki-67 between benign tissue, and PIN or cancer. However, the difference between PIN and cancer is not significant. Others have also detected low expression of Ki-67 in benign tissue²⁸⁵. Ki-67 positive cells shift from basal to luminal as one moves from benign, to PIN and cancer. This would imply, that not only is there a decrease in the level of apoptosis in cancer, but there is also increased cell production from benign to PIN or cancer and this is localised to the areas of malignant tissue.

The prostate is a hormone responsive organ and cell production is decreased by androgen ablation. Therefore this explains the decrease in Ki-67 expression associated with androgen ablation.

Prostatic Intraepithelial Neoplasia and Prostate Cancer.




Fig 43: Ki-67 positive staining PIN x 40.




Fig 44: Ki-67 positive staining cancer x 40.

Chapter 8: Conclusion.

In the third month of gestation the mesenchyme surrounding the urethra, in response to androgens, begins to differentiate into the capsule and the smooth muscle of the prostate. This along with the primitive lining of the urethra, which develops into ducts and the acini, forms the basic structures of the prostate. By the end of the 15th week of gestation, development of the basic prostate is complete. The dependence upon androgens that is so necessary early on will be hallmark of the development of this gland.

The prostate in response to androgens grows through out adult life. This results in a number of pathological conditions, both benign and malignant, which are major health issues. Yet, any intervention on the prostate gland is fraught, due to the close proximity of this gland to the urinary sphincter and the neurovascular bundles responsible for potency. Lowsley traditionally described the prostate as having five lobes, but a more useful zonal classification was proposed by McNeal²⁴⁸. The majority of prostate cancer develops in the peripheral zone.

Prostate cancer is the second commonest cause of cancer related death in males, yet the optimal management of localised carcinoma of the prostate represents a paradigm of uncertainty³³³. A number of risk factors have been implicated; these include age, race, family history, diet and androgens.

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

In this thesis, the various options have been outlined for the management of prostate cancer. The rational for these treatment modalities, whether it be watchful waiting, active surveillance, radiotherapy- external beam, conformal external beam, brachytherapy, or radiotherapy with adjuvant androgen ablation, surgery - open retro pubic, perineal, laparoscopic (extra or transperitoneal) has been discussed. Yet, despite the mushrooming options for the management of localised carcinoma only relatively recently has evidence been available that intervention does have any positive impact³³.

The increased interest in prostate cancer, has led to the discovery of a number of lesions which were thought to be premalignant, one of these is prostatic intraepithelial neoplasia⁴⁶, a lesion which exhibits the cytological changes of prostate cancer, but the basal cell layer is retained. Prostatic intra epithelial neoplasia has to be distinguished from atypia, transitional cell metaplasia, basal cell hyperplasia, clear cell cribriform hyperplasia, normal ejaculatory ducts, and seminal vesicle epithelium, high grade transitional cell carcinoma involving the ducts and acini, cribriform acinar and cribriform ductal prostatic carcinoma.

Initially prostatic intra epithelial neoplasia was thought to exist in three distinct forms, but this has been distilled to two, and then only one type, high-

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

grade prostatic intraepithelial neoplasia. Studies have shown that this lesion may precede prostate cancer, and can be detected in men as early as their third decade. The incidence of PIN in screening studies is between 0.7% and 16.5%, in TURP specimens it is between 2.8 and 3.2%. The discrepancy is probably due to the TRUS biopsies sampling the peripheral zone, and TURP chips being obtained from the transition zone.

PSA is a 240 amino acid glycoprotein serine protease with a molecular weight of 33,000. Initially used in forensic medicine it has evolved into one most important tumour markers used in medicine. The relationship between PIN and serum PSA had not been clearly defined, and it is hoped that this work has helped to elucidate this better.

To investigate the relationship between PIN, prostate cancer and serum PSA, this prospective study was undertaken. Prostate maps of fresh specimens obtained from radical prostatectomy specimens were constructed. The prostate was initially placed in 10% formaldehyde for 72 hours. The specimen was then sliced. The slices were scanned, and then fixed. Slides were prepared of the slices and areas of tumour and PIN identified and transcribed onto the prostate maps. The volume of cancer, PIN and of the total specimen was calculated from the scanned images, by measuring the area of interest from the scanned images and then multiplying it by the thickness of the slices. The volume so

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

calculated for each specimen was multiplied by a correction factor for shrinkage. The resulting volumes were compared to the serum PSA.

The volume of PIN is proportional to the total volume of the gland and the volume of benign tissue within the gland. The volume of cancer is inversely proportional to the benign and total volume of the gland, and directly related to the stage of the cancer.

PSA increases with the volume of benign and malignant tissue. One gram of benign tissue correlates with 0.07 ng/ml of serum PSA, and one gram of malignant tissue correlates with 0.8 ng/ml of serum PSA. Serum PSA does not correlate with the volume of PIN. The intensity of PSA staining decreases from benign, to PIN, to malignant tissue in a stepwise manner.

The importance of PIN is due to its predictive value for prostate cancer in up to 50% of patients with an initial diagnosis of PIN on biopsy. The relationship between PIN and prostate cancer was investigated at a molecular level using p₅₃, bcl-2, bax_{alpha} and Ki-67.

P₅₃ protein is the product of a gene located on the short arm of chromosome 17. The product of the p₅₃ gene has a molecular weight of 53 Kda, and the wild type p₅₃ by binding to DNA as a transcription factor inhibits the advancement

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

of cells from the G1 to the S phase, and so initiates apoptosis in defective cells - the Guardian of the genome. A mutant p₅₃ gene produces a defective protein, which can not regulate cell proliferation, and so cells may become immortal. Transfection of wild type p₅₃ into immortal cell lines can lead to growth arrest and apoptosis. Lack of a functioning p₅₃ confers an elevated risk of developing cancer – Li Fraumeni Syndrome⁴⁰⁰. The aberrant p₅₃ gene product has a longer half-life than the wild type and so is detected by immunohistochemical analysis.

Initially p₅₃ was thought to be an important prerequisite in localised carcinoma, and was used to predict outcome in patients at different stages of the disease with different Gleason scores. In patients who had undergone radical prostatectomy p₅₃ was thought to be able to predict outcome, but the weight of evidence at present points to its role being greater in metastatic disease, and causing extra prostatic spread.

In our experiments, p₅₃ expression was detected using immunohistochemical staining. The level of staining for p₅₃ did not increase significantly as one progressed from benign, to PIN, and to cancer. P₅₃ therefore does not appear to be important in the early genesis of prostate cancer. The presence of positive staining for p₅₃ appears to be a relatively infrequent event. There are some who agree with this⁴²¹ and others who do not³⁵⁶.

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

The discrepancy between our data and others may be due to the stage of the disease in patients. Even a recent study of 98 radical prostatectomy specimens, identified 57% of specimens were positive for p₅₃, but at a median follow up of 7.42 years (2.6 –12.9), 9 patients died from prostate cancer and 32 others progressed, implying relatively advanced disease at the time of surgery³⁵⁶.

There appears to be an increase in p₅₃ staining in the specimens obtained from patients who have received androgen ablation therapy prior to radical prostatectomy. However, this did not reach significance. The increase in expression may be due to androgen ablation causing apoptosis via a p₅₃-mediated pathway.

The bcl₂ gene was initially detected in B-cell follicular lymphomas. The bcl₂ proto-oncogene is located on the long arm of chromosome 18 and codes for a 26 kDa membrane protein. The bcl₂ protein inhibits apoptosis, possibly by limiting the release of pro-apoptotic enzymes from the mitochondria and so promotes survival.

Bcl₂ expression can allow prostate cancer cell lines to survive in the absence of androgens, and so can negate the therapeutic effect of androgen ablation in the management of prostate cancer³¹³. Further more in animal experiments

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

androgen ablation led to an increase in the mRNA for bcl₂, this may support the hypothesis that trigger for the development of androgen insensitive controls is the withdrawal of androgens²⁴⁶.

Bcl₂ expression in radical prostaectomy specimens is associated with higher stage and shorter progression free survival⁶¹. In patients being treated with radiotherapy, the expression of bcl₂ is associated with shorter time to relapse¹⁷³.

Bcl₂ expression increases from benign to PIN and then decreases in cancer. The high levels of expression of bcl₂ in PIN, would allow PIN to become relatively resistant to androgen ablation and chemotherapeutic intervention. The relative survival advantage conferred on PIN, could allow it to become the source for androgen insensitive clones. This is further supported by evidence that shows that bcl₂ expression appears to have an inverse relationship with androgen receptor expression⁴¹.

Androgen ablation led to an up regulation in the expression of bcl₂. This allows the cells to become androgen independent³¹³ and also inhibits p₅₃ mediated apoptosis by preventing the wild type p₅₃ from translocating to the nucleus and triggering apoptosis²¹. Increased bcl₂ also causes increased expression of VEGF, and so increased microvessel density, allowing cells to

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

survive in hostile environments¹¹⁶.

Bcl associated x protein or bax is a proapoptotic molecule. It is a cytosolic protein which forms a heterodimer with bcl₂. This causes the bax molecule to translocate to the mitochondrial membrane and bind to the membrane bound bcl₂ protein. This binding results in inactivation of bcl₂ and cell death.

The expression of bax was ubiquitous, in benign, PIN, and prostate cancer. The staining intensity was however greater in prostate cancer (p=0.03) and PIN (p=0.06) than in benign tissue, implying that there may be increased production of bax on a cellular basis leading to increased turnover of malignant and pre-malignant cells. Androgen ablation decreased the staining of cells expressing bax and so this may lead to decreased cell turnover via the bcl₂/bax axis, especially since there is increased expression of bcl₂.

The gene coding for the Ki-67 protein is on chromosome 10. Ki-67 is a nuclear protein expressed in the G1, S, G2, and M phases of the cell cycle, but not in G0 – the resting phase of the cell cycle. Therefore MIB-1 antibody staining can be used in formalin fixed tissue to determine the proliferation rate. The alternative is thymidine incorporation assays, but for this fresh tissue is required and this is a very demanding technique. The proliferation rate as measured by Ki-67 can be used to predict extra-capsular spread²²⁹,

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

recurrence¹⁴² and outcome after surgery or radiotherapy¹. The proliferation rate may also increase with Gleason grade

In our series the proliferation index increases from benign, to PIN, and cancer. The increase from benign to PIN and from benign to cancer is significant. If the proliferation rate as measured by Ki-67 is a predictor of the malignant potential of tissue, then PIN may be part of the way there.

Others smaller studies have supported this stepwise increase in MIB-1 expression when comparing benign tissue, with PIN and cancer
160:187:275:379:393

Prostate cancer is an androgen responsive tumour, and by removing the stimulus for proliferation by chemical androgen ablation, the proliferation rate declines to a level significantly less than cancer, but still higher than benign tissue. This would support the view that androgens are not the only regulator of proliferation, and that these cells are displaying some autonomy from their endocrine regulators even at this early stage.

Bcl₂ staining increases from benign to malignant tissue, and so allowing the cancer cells to be resistant to intervention with androgen ablation or chemotherapy. Ki-67 increases from BPH to PIN and from BPH to cancer.

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

Implying that there is increased cell production in malignant and pre-malignant tissue compared to benign tissue.

Androgen ablation led to a decrease in Ki-67 staining, an increase in the number of specimens expressing bcl_2 and a decrease in the staining intensity for bax, implying LHRH not only leads to a decrease in the production of new cells but also to the selection of androgen resistant clones.

In summary in this thesis, it has been shown that the volume of cancer is proportional to the local stage of the disease; it is inversely proportional to the volume of benign and PIN tissue. The total gland volume is proportional to the volume of PIN. One gram of benign tissue correlates with 0.07 ng/ml of serum PSA, and one gram malignant tissue correlates with 0.8 ng/ml of serum PSA, the volume of PIN does not correlate with serum PSA.

With respect to the immunohistochemical studies, p53 is a rare event in localized disease; bcl_2 is significantly over expressed in PIN compared to cancer or benign tissue, and the intensity of bax staining significantly increases when comparing benign tissue with PIN or malignant tissue, as does the expression of Ki-67.

Androgen ablation causes a significant increase in the expression of bcl_2 , and decrease in the expression of Ki-67 and staining intensity of bax.

Chapter 9: Publications and Presentations.

9.1 Presentations resulting from this work.

CO-EXISTENCE OF INTRA-EPITHELIAL NEOPLASIA AND PROSTATE CARCINOMA.

A V Kaisary, **S S Sandhu**, P J Richmond and M Jarmulowicz.

Presented at the Sixth International Prostate Cancer Update, Denver Colorado USA, January 1996.

THE CORRELATION OF SYSTEMATIC SEXTANT BIOPSY WITH PATHOLOGICAL EXAMINATION OF RADICAL PROSTATECTOMY SPECIMENS.

S S Sandhu, VB Matveev, M Jarmulowicz, and A V Kaisary.

Presented at the British Association of Urological Surgeons, Annual Meeting June 1997.

The British Journal of Urology 1997;79(Suppl 4);abs104, p26-27.

THE EFFECT OF LHRH AGONISTS ON PROSTATIC INTRAEPITHELIAL NEOPLASIA.

M Jarmulowicz, **S S Sandhu**, T Walker and A V Kaisary.

Presented at the Pathological Society of Great Britain and Ireland, Winter Meeting January 1998.

THE CORRELATION OF SYSTEMATIC SEXTANT BIOPSY WITH PATHOLOGY.

S S Sandhu, M Jarmulowicz, T Walker, N Simmonds, Wilson C and A V Kaisary.

Presented at the Sixth Mediterranean Congress of Urology, Cairo, Egypt. Sept 1999.

THE CORRELATION OF PREOPERATIVE TRUS GUIDED SEXTANT BIOPSIES WITH RADICAL PROSTATECTOMY SPECIMENS.

S S Sandhu, T Walker, D St George, M Jarmulowicz, and A V Kaisary.

Presented at the Congress of the European Association of Urology, Feb 2002. European Urology Supplement Vol 1, No 1; abs659, p167.

Selected as the Best Poster in the session.

HIGH GRADE PROSTATIC INTRAEPITHELIAL NEOPLASIA, SERUM PSA, AND INVASIVE CANCER.

S S Sandhu, T Walker, D St George, M Jarmulowicz, and A V Kaisary.

Presented at the British Prostate Group Spring Meeting, April 2002.

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

PROSTATE CANCER, PROSTATIC INTRAEPITHELIAL NEOPLASIA AND SERUM PSA.

S S Sandhu, T Walker, C Thrasyvoulou, M Jarmulowicz, and A V Kaisary.
Presented at Progress and Controversies in Oncological Urology (PACIO VII) and the seventh Congress of the Dutch Urological Association (DUA VII), October 2002.

TO INVESTIGATE USING MORPHOMETRIC CRITERIA THE RELATIONSHIP BETWEEN HIGH GRADE PROSTATIC INTRAEPITHELIAL NEOPLASIA PROSTATE CANCER, AND SERUM PSA.

S S Sandhu, T Walker, M Jarmulowicz and AV Kaisary.
Presented at the Société Internationale d'Urologie, Biannual Meeting October 2004. The British Journal of Urology International 2004: 94(Suppl 2);absUP-6.15, p194.

AN INVESTIGATION INTO THE RELATIONSHIP BETWEEN PROSTATIC INTRAEPITHELIAL NEOPLASIA AND PROSTATE CANCER.

S S Sandhu, T Walker, M Jarmulowicz and AV Kaisary.
Presented at the Société Internationale d'Urologie, Biannual Meeting October 2004. The British Journal of Urology International 2004: 94(Suppl 2);absUP-6.40, p200.

9.2 Publications resulting from this work.

TRIALS OF TREATMENT ARE NEEDED BEFORE TRIALS OF SCREENING.

S S Sandhu, R Morris, V Matveev, and A V Kaisary.
The British Medical Journal (1996): 312;709.

LOCALISED CARCINOMA OF THE PROSTATE: A PARADIGM OF UNCERTAINTY.

S S Sandhu and A V Kaisary.
Postgrad Med J 1997;**73**:691-696.

Appendix

Appendix 1: Area Calculation.

```
# Area measurements from micrographs for Sarb
# Chris Thras 4.01.2000
MSsetprop "FRAMEMODE",0
# showwindow "Editor",1

while 1
loops = 1 # reset loop number
    # imgsetpath "d:\sarb"
MSload "sarb"
# MSsetfeat "FIELDFEAT"

DBsetpath "c:\ks400\data\sarb"
dbnamef= "databases"
#specimen loop to allow data from several fields to be appended

while
read dbnamef, "Input the field data base name you want to use"
n = DBexist (dbnamef)
if n == 0 : break
MBok "Database name exists - try again"
endwhile

while 1
imgdelete "*"
Gclear 0
imgnew 1,768,1024,1,"Grey"
imgdisplay 1
MBok " Copy and past total image from PSP into image 1"
imgnew 2,768,1024,1,"Grey"
MBok " Copy and past PIN image from PSP into image 2"
imgnew 3,768,1024,1,"Grey"
MBok " Copy and past CANCER image from PSP into image 3"
imgdisplay 1
greytran 1,4,"invers.col"
disaut 4,7,0,1
binfill 7,10
imgdisplay 2
greytran 2,5,"invers.col"
```

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

```
disaut 5,8,0,1
imgdisplay 3
greytran 3,6,"invers.col"
disaut 6,9,0,1

if loops == 1
imgdisplay 1
MSsetgeom
! MSmeasmask 10,1,dbnamef,0,0,10
imgdisplay 8
! MSmeasmask 8,1,dbnamef,1,0,10
imgdisplay 9
! MSmeasmask 9,1,dbnamef,1,0,10
datalist dbnamef,0,0

else

imgdisplay 1
MSsetgeom
! MSmeasmask 10,1,dbnamef,0,0,10
imgdisplay 8
! MSmeasmask 8,1,dbnamef,1,0,10
imgdisplay 9
! MSmeasmask 9,1,dbnamef,1,0,10
datalist dbnamef,0,0

endif
loops = loops + 1
# MByesno "Do you have another section in this group?"
  if _STATUS == 0 :break

endwhile
MByesno "Another sample/specimen?"
  if _STATUS == 0 :break
endwhile
beep 100,880

stop
```


Appendix No. 2 : Shrinkage

Pt Id	Fresh Wt	Fresh Vol	Fresh Density	Fx Wt	Fx Vol	Fixed Density	% Change in Wt	% Change in Vol
D	71.57	67.75	1.06	71.61	66.90	1.07	-0.05	1.25
G	61.20	56.70	1.08	60.90	55.80	1.09	0.49	1.59
H	44.01	43.60	1.01	42.86	41.61	1.03	2.61	4.57
M	42.70	40.97	1.04	41.96	39.65	1.06	1.73	3.21
To	60.12	57.24	1.05	59.70	56.72	1.05	0.69	0.91
Tu	70.79	69.95	1.01	70.30	66.45	1.06	0.69	5.00
Average	58.40	56.03	1.04	57.89	54.52	1.06	1.03	2.76

Therefore if the true volume is x, the volume measured should be multiplied by $100/(100-\% \text{ Change in volume})$ ie.
1.03

Pt Id	Fx Slice Area	Mounted Slice Area H+E	Change in Area
D	1753.00	1560.62	10.97
G	1228.00	1027.30	16.34
H	1437.00	1165.50	18.89
M	1114.00	941.47	15.49
To	1842.50	1421.69	22.84
Tu	1535.60	1364.44	11.15

15.95

Therefore if the fixed slice area is y, the area measured should be multiplied by $100/(100-\% \text{ Change in area})$ ie.
1.19 and this will give the area of the H+E slides prior to being embedded in wax, and this should be multiplied by
1.03 and this will give the fresh area ie 1.22348

Appendix 3: Slice thickness

Pt Id	Slice 2	Slice 3	Slice 4	Slice 5	Slice 6	Slice 7	Slice 8	Total Slice Thickness	Av Slice thickness
D	3.17	3.38	3.39	3.31	3.52	3.3	3.77	23.84	3.41
G	2.77	2.92	3.08	3.01	3.06	2.7	3.62	21.16	3.02
H	3.02	3.36	3.32	3.42	3.75	4.28	3.28	24.43	3.49
M	3.19	3.26	3.41	3.79	3.61	4.21	4.33	25.80	3.69
To	2.42	2.63	3.04	3.38	3.58	3.23	3.29	21.57	3.08
Tu	3.88	3.62	3.19	3.24	3.36	3.41	3.63	24.33	3.48

To get the volume one should multiply the area by the thickness of the slices multiplied by the shrinkage fraction for the fixation in formalin i.e. 1.028335. i.e. 3.46. This has been corrected to one decimal place as 3.5mm.

Appendix 4:**Preparation of 3-Aminopropyltriethoxysilane (APES) coated slides.****Royal Free Hospital School of Medicine – Standard Operating Procedure
No.5**

METHOD (van Prooijen-Knegt ACet al. 1983; MaddoxPH and Jenkins, 1987).

1. Degrease slides in hot detergent solution, rinse in distilled water then alcohol (740 P) and air dry for 10 minutes.
2. Immerse slides in a freshly prepared 2 % solution of APES in acetone-
minimum 1 minute.
3. Drain slides briefly and wash in distilled water – 2 quick dips.
4. Drain slides on paper towels and dry at 40⁰C over night in oven on paper towels.
5. Store at room temperature in empty slide boxes.

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

Appendix No.5

Preparation Paraffin Sections for immunohistochemistry

Royal Free Hospital School of Medicine – Standard Operating Procedure No.2

1. Cool paraffin blocks.
2. Cut paraffin sections at 3 micro m.
3. Float sections on 20% alcohol then 45⁰C water bath.
4. Pick up sections on APES coated glass slides as appropriate.
5. Place slides in 37⁰C oven over night
6. De-parafinise sections and take them to alcohol.
 - a) Xylene 1 5 minutes
 - b) Xylene 2 5 minutes
 - c) Xylene 3 5 minutes
 - d) Absolute Alcohol 2 minutes
 - e) Absolute Alcohol 2 minutes
 - f) Absolute Alcohol 2 minutes
 - g) Double distilled water 2 minutes

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

Appendix No.6: Staining procedure.

Royal Free Hospital School of Medicine – Standard Operating Procedure No.13

1. Deparaffinise sections and take to alcohol.
2. Transfer sections to humidity chamber and block endogenous peroxidase activity.
3. Wash in 3 times changes of de-ionised or distilled water for 2 minutes each.
4. Use antigen retrieval system.
5. Transfer slides to a humidity chamber and cover sections with two changes of Tris buffered saline (TBS) for 5 minutes each.
6. Block non specific binding by covering with normal goat serum 1/10 in TBS for 15 minutes at room temperature.
7. Drain and cover sections with optimally diluted primary antibody diluted in TBS. Leave sections in humidity chamber for the appropriate time and temperature for the antibody.
8. Rinse in TBS twice for 3 minutes.
9. Drain and cover sections with biotinylated goat anti rabbit/mouse antisera 1/200 in 4% normal human sera for 30 minutes in a humidity chamber at room temperature.
10. Repeat step 8.
11. Drain and cover sections with STR-ABC 1/200 for 30 minutes in a humidity chamber.
12. Repeat step 8.
13. Develop sections with freshly prepared Diaminobenzidine (DAB) solution.
14. Wash in distilled water for 5-10 minutes.
15. Counter stain with Mayer's haemotoxylin for 2 minutes.
16. Rinse in distilled water. Differentiate in 0.5% acid alcohol and blue for 5 minutes.
17. Dehydrate, clear, and mount.
18. Sites of immunoperoxidase activity = Brown,
Nuclei = Blue.

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

Appendix No. 7 : Blocking endogenous peroxidase activity and alkaline phosphatase activity.

Royal Free Hospital School of Medicine – Standard Operating Procedure No.3

Hydrogen Peroxide/Water Method

1. De-paraffinise sections and take to water
2. Transfer slides to a humidity chamber at room temperature and cover the sections with freshly prepared 0.5 % H_2O_2 in distilled water for 10-15 minutes. This solution contains: 0.2 mls H_2O_2 (30% W/Vol.) and 11.8 mls water (enough for approx, 20 slides).
3. Wash sections in distilled water 3 times and proceed with immunostaining method.

Appendix No. 8
Buffers for Immuno-histochemistry

EQUIPMENT

Magnetic stirrer
pH meter
Balance
Conical flask
10 Litre plastic vat
Magnetic flea

A. Tris buffered saline (TBS) pH 7.6.

REAGENTS

Sodium chloride	80 gm
Tris (Tris hydroxymethylamine)	6.6 gm
1 M Hydrochloric Acid	44 mls
Double distilled water	10 litres

METHODS

1. Mix reagents with 2 litres of distilled water on a magnetic stirrer with a magnetic flea.
2. When components are dissolved transfer solution to a plastic 10-litre vat.
3. Add 8 litres of water and mix thoroughly.
4. Take an aliquot and pH to 7.6 (+/-0.05) using buffer components.
5. Buffers must be as fresh as possible in order to prevent adverse affects on staining. If storing, keep at 4 °C ready for use.

B. Phosphate.buffered saline (PBS) pH 7.4

REAGENTS

PBS tablets (Oxoid)	1
Double distilled water	100mls

METHODS

1. Mix reagents and check pH is 7.4 (+/- 0.05). If not correct pH check water pH and/or quality of tablets.

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

Appendix No. 9

Microwave technique for immuno-histochemistry ^{355 352 354 353 383 384}

EQUIPMENT

pH meter

Microwave (700W) oven with rotation plate.

Balance

Plastic coplin jars

1 L conical flask

Citric acid buffer

REAGENTS

Citric Acid Monohydrate

2.1g

Double distilled water.

10 litres

2M NAOH

13 mls

pH to 6.0 with buffer components

METHOD

1. De-paraffinise sections (appendix 2) sections must be mounted on APES slides (see appendix 1) or there will be section loss.
2. Block endogenous peroxidase activity (appendix 4)
3. Wash slides with PB S.
4. Fill plastic coplin jar 4/5 with citric acid buffer 0.01 m Citric acid buffer solution and submerge sections.

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

Appendix 10 Raw Morphimetric data

Pt Id	PSA	Total vol	Ca Vol	PIN Vol	Benign vol	Pros Weight
1	7	39605.6	2467.9	544.6	36593.1	42.03
2	5	37719.8	1922.2	250.9	35546.8	50.70
3	13.5	80401.6	8947.7	44.6	71409.3	93.00
4	14	64846.7	2656.1	307.5	61883.1	75.00
5	4.5	42495.4	2984.7	875.4	38635.3	52.30
6	9.6	59843.0	901.5	45.7	58895.7	67.80
7	9.6	42027.8	3774.7	836.9	37416.2	57.00
8	9.2	60964.2	612.5	243.0	60108.7	79.00
9	4.9	67633.6	8.9	12.5	67612.2	67.00
10	8.4	67063.1	22.2	1157.2	65883.7	77.80
11	5.6	80449.2	5089.8	234.5	75124.8	90.00
12	7.2	60497.7	2243.0	2300.2	55954.4	74.20
13	10	57438.6	957.8	117.9	56362.9	70.00
14	6.1	35346.4	1965.9	117.3	33263.2	47.60
15	9.7	89791.2	130.8	1132.1	88528.3	91.30
16	5.9	37108.8	3632.7	153.0	33323.1	44.00
17	5.5	70380.8	652.7	775.1	68952.9	75.50
18	6.2	39543.2	3265.3	487.2	35790.8	55.00
19	6.9	31638.0	1880.8	278.5	29478.7	46.00
20	5	40453.1	1543.9	510.0	38399.3	54.00
21	5.4	56137.8	430.2	462.9	55244.7	59.80
22	13.7	74299.1	2024.0	2728.5	69546.5	82.70
23	6.1	36654.9	1793.4	397.5	34464.0	42.00
24	13.2	39420.7	857.8	1514.2	37048.6	41.86
25	32.4	62789.4	15150.6	249.9	47388.9	88.40
26	5.3	41864.9	2519.3	180.8	39164.8	50.00
27	16.1	64089.7	293.3	479.5	63316.8	63.29
28	8.1	84674.2	4073.2	465.4	80135.7	99.00
29	10.3	55855.5	113.5	220.4	55521.6	71.00
30	9.6	83056.3	3704.6	400.0	78951.8	89.00
31	7.5	40394.0	1507.8	509.3	38376.9	43.01
32	4.1	46648.7	382.6	411.0	45855.1	57.80
33	5.2	29175.3	2767.0	64.9	26343.4	35.00
34	3.6	24503.2	1934.3	78.6	22490.3	32.20
35	5.7	69594.5	5324.7	343.3	63926.5	87.50
36	8.3	50470.0	3074.4	0.0	47395.6	59.00
37	6.2	28633.4	1564.1	283.5	26785.8	39.50
38	10.1	40458.6	1254.1	266.7	38937.8	48.00
39	11	39406.3	656.2	29.7	38720.4	61.60
40	7.3	43101.7	903.4	296.2	41902.1	52.30
41	5	62086.4	393.9	70.4	61622.1	81.00
42	15	23074.0	1427.0	132.2	21514.7	37.00
43	7.9	37088.2	1638.8	174.2	35275.2	42.40
44	12.4	34313.5	2153.4	207.1	31953.0	51.10
45	7.2	72421.6	694.0	1558.4	70169.2	67.80
46	8.2	70390.9	664.0	358.6	69368.3	71.00

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

47	4.2	19359.9	5839.4	222.6	13298.0	31.80
48	3	71721.5	139.1	985.9	70596.5	79.00
49	11.5	68491.4	3170.7	392.2	64928.5	73.80
50	7.4	41647.8	8847.8	328.5	32471.5	59.10
51	3.6	43831.8	946.1	2259.3	40626.3	54.00
52	10.7	51745.4	209.9	2406.5	49129.1	67.00
53	12.6	30462.3	2778.9	32.9	27650.5	42.00
54	7	34510.7	520.9	887.3	33102.4	52.00
55	5.3	32558.9	1874.2	605.9	30078.8	44.00
56	12.1	52898.0	12286.9	438.5	40172.5	57.35
57	5.3	35251.5	4213.2	156.8	30881.5	46.50
58	7.1	38953.1	3436.1	152.0	35365.0	42.00
59	8.1	67732.9	160.6	354.4	67217.9	49.75
60	8.3	45525.4	1550.7	1535.8	42438.9	65.00
61	45	59354.2	4282.1	635.8	54436.2	62.90
62	6.2	60233.1	854.2	255.9	59123.0	74.60
63	9.9	32028.5	5823.5	572.9	25632.1	34.67
64	14.4	31259.8	4905.4	449.5	25904.9	43.00
65	6.3	48891.0	57.2	23.5	48810.3	59.00
66	6	35443.7	2725.4	304.1	32414.2	41.00
67	11.5	56645.0	761.9	933.8	54949.3	70.00
68	14	51785.8	2094.0	21.6	49670.3	68.40
69	4.8	29300.4	1186.6	18.4	28095.4	39.00
70	9	24580.4	5062.5	26.5	19491.4	26.00
71	13	155519.0	142.2	1939.8	153437.0	173.00
72	6.2	47481.5	665.3	125.1	46691.1	55.70
73	6.7	34677.6	3283.6	378.7	31015.2	43.50
74	6.2	41129.4	2284.5	400.7	38444.3	48.22
75	7.5	48382.7	507.7	1151.6	46723.4	58.00
76	13.4	64003.2	466.7	2.9	63533.6	68.60

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

Appendix 11: Raw Immunohistochemical data.

Data obtained from regions of prostate cancer.

	Number of + cells P53 Ca	P53 Ca	PSA Ca	Number of + cells Bcl2 Ca	Bcl2 Ca	Ki67 Ca	Bax Ca
1	0	0	2	0	0	1	1
2	0	0	2	0	0	1	1
3	0	0	3	0	0	1	2
4	0	0	3	0	0	1	0
5	0	0	2	60	1	1	2
6	0	0	3	0	0	1	3
7	100	2	1	40	1	1	1
8	0	0	2	0	0	1	0
9	0	0	3	50	1	1	3
10	0	0	1	0	0	1	2
11	0	0	2	0	0	1	3
12	0	0	2	0	0	1	1
13	0	0	3	0	0	1	2
14	0	0	2	10	1	1	2
15	100	2	3	100	2	1	3
16	0	0	3	0	0	1	3
17	0	0	3	0	0	1	1
18	0	0	3	20	1	1	3
19	0	0	2	10	1	1	2
20	0	0	2	0	0	1	2
21	0	0	2	0	0	1	1
22	100	1	2	0	0	1	1
23	0	0	1	0	0	1	1
24	0	0	2	80	2	1	2
25	0	0	3	0	0	35	1
26	0	0	3	0	0	8	1
27	0	0	3	0	0	5	2
28	0	0	3	0	0	4	2
29	0	0	2	0	0	10	2
30	0	0	2	0	0	3	1
31	0	0	2	0	0	3	2
32	0	0	3	100	1	5	1
33	0	0	2	0	0	1	2
34	0	0	3	0	0	1	2
35	50	1	1	50	1	7	3
36	0	0	1	0	0	0	2
37	80	2	1	0	0	7	2
38	0	0	3	0	0	10	2
39	0	0	2	0	0	3	1
40	0	0	2	0	0	2	2
41	0	0	3	0	0	3	3
42	0	0	2	0	0	5	2
43	0	0	1	0	0	3	2
44	0	0	2	0	0	10	2

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

45	0	0	3	0	0	1	1
46	0	0	3	50	1	10	2
47	0	0	3	30	1	3	2
48	0	0	2	0	0	15	2
49	0	0	2	0	0	5	2
50	0	0	3	0	0	5	1
51	0	0	3	40	1	10	2
52	0	0	1	0	0	15	2
53	0	0	3	0	0	1	1
54	0	0	2	0	0	12	3
55	0	0	2	0	0	1	1
56	0	0	2	10	1	30	1
57	0	0	3	0	0	20	3
58	0	0	1	100	1	10	2
59	10	1	2	0	0	5	1
60	0	0	2	0	0	6	1
61	0	0	2	0	0	10	2
62	0	0	1	50	1	4	2
63	0	0	1	0	0	8	1
64	50	1	3	50	2	1	1
65	0	0	1	30	1	20	2
66	50	2	2	0	0	5	2
67	0	0	3	0	0	1	1
68	0	0	1	10	1	5	2
69	0	0	2	80	1	5	2
70	0	0	3	0	0	10	2
71	0	0	3	100	1	5	2
72	0	0	0	0	0	0	1
73	100	1	3	50	1	1	3
74	0	0	0	100	2	3	1
75	0	0	2	0	0	10	2
76	0	0	1	0	0	5	2
77	0	0	2	0	0	10	3
78	0	0	1	50	1	5	2
79	0	0	3	0	0	1	1
80	0	0	2	60	1	5	2
81	0	0	2	100	1	15	3
82	0	0	3	0	0	7	2
83	0	0	2	0	0	3	2
84	0	0	2	0	0	1	2
85	0	0	1	0	0	10	1
86	0	0	3	10	1	10	3
87	0	0	1	0	0	4	2
88	0	0	2	0	0	10	2
89	0	0	2	0	0	15	1
90	0	0	3	0	0	1	1
91	0	0	2	0	0	20	2
92	0	0	2	0	0	10	1

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

Data obtained from regions of PIN.

	Number of + cells P53 PIN	P53 PIN	PSA PIN	Number of + cells Bcl2 PIN	BCl2 PIN	Ki67 PIN	Bax PIN
1	0	0	3	60	2	1	1
2	0	0	2	0	0	1	0
3	0	0	3	80	1	1	1
4	0	0	3	0	0	1	2
5	0	0	3	0	0	1	2
6	0	0	3	60	2	1	2
7	0	0	3	20	1	1	1
8	0	0	3	100	2	1	1
9	0	0	2	80	2	1	1
10	0	0	3	100	2	1	1
11	0	0	3	50	2	1	1
12	0	0	3	30	2	1	2
13	0	0	3	25	2	1	1
14	0	0	3	40	1	1	2
15	0	0	3	100	2	1	1
16	0	0	3	100	2	1	1
17	100	3	3	10	2	1	1
18	0	0	3	80	2	1	2
19	0	0	3	60	2	1	3
20	0	0	3	5	1	1	2
21	0	0	3	25	2	1	2
22	0	0	2	0	0	1	2
23	0	0	3	0	0	0	2
24	0	0	2	60	1	5	2
25	0	0	2	100	1	1	2
26	25	2	3	80	2	5	1
27	0	0	2	60	2	2	3
28	0	0	3	0	0	5	2
29	0	0	2	20	1	5	2
30	0	0	2	100	1	3	2
31	0	0	2	0	3	5	2
32	0	0	3	10	2	5	2
33	0	0	3	10	2	1	2
34	0	0	3	60	1	2	1
35	50	2	3	100	1	5	3
36	0	0	2	100	5	4	2
37	50	1	2	0	0	7	2
38	0	0	3	30	1	5	2
39	0	0	2	70	3	5	1
40	0	0	3	60	2	4	2
41	0	0	3	50	2	3	3
42	0	0	1	80	2	1	1
43	0	0	1	0	0	3	2
44	90	1	3	50	1	3	2
45	0	0	3	100	2	2	2
46	0	0	3	30	2	3	2

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

47	0	0	3	80	2	20	2
48	0	0	2	0	0	15	2
49	0	0	2	70	2	2	2
50	0	0	3	70	2	1	2
51	0	0	3	0	0	5	2
52	0	0	2	30	2	3	2
53	0	0	3	10	1	1	1
54	0	0	2	30	2	5	3
55	0	0	3	100	2	1	1
56	0	0	2	5	1	40	1
57	0	0	3	30	1	3	3
58	0	0	2	80	2	5	2
59	0	0	2	0	0	5	0
60	0	0	0	0	0	0	0
61	0	0	3	50	1	15	2
62	0	0	3	50	2	1	2
63	0	0	2	40	2	5	1
64	0	0	0	0	0	0	0
65	0	0	1	10	2	15	2
66	0	0	3	100	2	10	2
67	0	0	3	0	0	1	1
68	0	0	2	10	2	2	2
69	0	0	3	100	1	8	2
70	0	0	3	0	0	10	2
71	0	0	3	70	1	5	0
72	0	0	3	10	1	5	2
73	0	0	2	100	2	1	0
74	0	0	0	0	0	0	1
75	0	0	3	0	0	10	2
76	0	0	2	80	1	1	2
77	0	0	3	30	1	25	3
78	0	0	3	30	1	3	2
79	0	0	1	100	3	1	1
80	0	0	3	50	2	5	2
81	0	0	2	100	1	15	3
82	0	0	3	0	0	5	2
83	0	0	3	60	2	3	2
84	0	0	3	10	1	1	2
85	0	0	2	50	1	2	0
86	0	0	3	60	2	3	3
87	0	0	3	100	1	2	2
88	0	0	2	100	2	5	2
89	0	0	2	40	1	5	1
90	0	0	2	80	1	1	1
91	0	0	2	30	1	20	2
92	0	0	3	100	1	10	1

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

Data obtained from specimens with 3 months of androgen ablation.

	Number of + cells	P53 LHRH	Number of + cells	BCI2 LHRH	Ki67 LHRH	Bax LHRH
1	0	0	100	3	1	1
2	0	0	70	1	1	1
3	0	0	0	0	1	1
4	0	0	50	1	1	1
5	0	0	100	2	1	2
6	0	0	100	2	1	2
7	0	0	0	0	25	1
8	0	0	60	1	1	2
9	50	3	0	0	1	2
10	80	1	100	2	1	1
11	0	0	0	0	2	2
12	100	1	0	0	1	1
13	0	0	100	1	1	2
14	80	1	100	1	1	1

Data obtained from benign (control specimens)

	P53 Benign	PSA Benign	Number of + cells	BCI2 Benign	Ki67	Bax
1	0	3	0	0	0	1
2	0	3	0	0	0	2
3	0	3	0	0	0	1
4	0	3	0	0	0	2
5	0	3	0	0	0	1
6	0	3	0	0	0	2
7	0	3	0	0	0	0
8	0	3	0	0	0	2
9	0	3	0	0	0	2
10	0	3	0	0	0	1
11	0	3	0	0	0	2
12	0	3	0	0	0	1
13	0	3	0	0	0	1
14	0	3	0	0	0	2
15	0	3	0	0	0	1
16	0	3	0	0	0	2
17	0	3	0	0	0	1
18	0	3	0	0	0	1
19	0	3	0	0	0	1

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

Reference List

1. Aaltomaa S, Lipponen P, Vesalainen S, Ala-Opas M, Eskelinen M, Syrjanen K. Value of Ki-67 immunolabelling as a prognostic factor in prostate cancer. *European Urology* 1997;**32**:410-5.
2. Aarnink RG, de la Rosette JJ, Huynen AL, Giesen RJ, Debruyne FM, Wijkstra H. Standardized assessment to enhance the diagnostic value of prostate volume; Part II: Correlation with prostate-specific antigen levels. *Prostate* 1996;**29**:327-33.
3. Abrahamson PA. Peptide hormone and serotonin immunoreactive cell in normal and hyperplastic prostate glands. *Path Res Pract* 1986:181, 675-683.
4. Adolfsson J, Carstensen J, Lowhagen T. Deferred treatment in clinically localised prostatic carcinoma. *British Journal of Urology* 1992;**69**:183-7.
5. Aihara M, Lebovitz RM, Wheeler TM, Scardino PT, Ohori M, Scardino PT. Prostate specific antigen and gleason grade: an immunohistochemical study of prostate cancer. *Journal of Urology* 1994; **151**:1558-64.
6. Alexander EE, Qian J, Wollan PC, Myers RP, Bostwick DG. Prostatic intraepithelial neoplasia does not appear to raise serum prostate-specific antigen concentration. *Urology* 1996;**47**:693-8.
7. Allen RT, Cluck MW, Agrawal DK. Mechanisms controlling cellular suicide: role of Bcl-2 and caspases. *Cellular & Molecular Life Sciences* 1998;**54**:427-45.
8. Alsikafi NF, Gerber GS, Yang XJ, Brendler CB. High grade prostatic intraepithelial neoplasia (PIN) with adjacent atypical glands is associated with a higher rate of cancer on subsequent needle biopsy than high grade PIN alone. *J Urol* 2000;**163**:90, Abs 394.

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

9. Andriani F, Nan B, Yu J, Li X, Weigel NL, McPhaul MJ *et al.* Use of the probasin promoter ARR2PB to express Bax in androgen receptor-positive prostate cancer cells. *Journal of the National Cancer Institute* 2001;**93**:1314-24.
10. Apakama I, Robinson MC, Walter NM, Charlton RG, Royds JA, Fuller CE *et al.* bcl-2 overexpression combined with p53 protein accumulation correlates with hormone-refractory prostate cancer. *British Journal of Cancer* 1996;**74**:1258-62.
11. Aprikian AG, Sarkis AS, Fair WR, Zhang ZF, Fuks Z, Cordon-Cardo C. Immunohistochemical determination of p53 protein nuclear accumulation in prostatic adenocarcinoma. *Journal of Urology* 1994;**151**:1276-80.
12. Aumuller, G. Prostate gland and seminal vesicles. 79. Berlin, Springer-verlag.
13. Baas IO, Mulder JW, Offerhaus GJ, Vogelstein B., Hamilton SR. An evaluation of six antibodies for immunohistochemistry of mutant p53 gene product in archival colorectal neoplasms. *Journal of Pathology* 1994;**172**:5-12.
14. Bagshaw MA, Kaplan I, Cox RC. Prostate cancer. Radiation therapy for localized disease. *Cancer* 1993; **71**:939-52.
15. Bai XZ, Masters JR, O'Donoghue N, Kirby R, Pan LX, Young M *et al.* Prognostic markers in clinically localised prostate cancer. *International Journal of Oncology* 1999;**14**:785-91.
16. Baretton GB, Klenk U, Diebold J, Schmeller N, Lohrs U. Proliferation- and apoptosis-associated factors in advanced prostatic carcinomas before and after androgen deprivation therapy: prognostic significance of p21/WAF1/CIP1 expression. *British Journal of Cancer* 1999;**80**:546-55.

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

17. Bartek J, Bartkova J, Vojtesek B, Staskova Z, Lukas J, Rejthar A *et al.* Aberrant expression of the p53 oncogene is a common feature of a wide spectrum of human malignancies. *Oncogene* 1991;**6**:1699-703.
18. Bartek J, Iggo R, Gannon J, Lane DP. Genetic and immunochemical analysis of mutant p53 in human breast cancer cell lines. *Oncogene* 1990;**5**:893-9.
19. Bauer JJ, Sesterhenn IA, Mostofi FK, McLeod DG, Srivastava S, Moul JW. p53 nuclear protein expression is an independent prognostic marker in clinically localized prostate cancer patients undergoing radical prostatectomy. *Clinical Cancer Research* 1995;**1**:1295-300.
20. Bauer JJ, Sesterhenn IA, Mostofi FK, McLeod DG, Srivastava S, Moul JW. Elevated levels of apoptosis regulator proteins p53 and bcl-2 are independent prognostic biomarkers in surgically treated clinically localized prostate cancer. *Journal of Urology* 1996;**156**:1511-6.
21. Beham A, Marin MC, Fernandez A, Herrmann J, Brisbay S, Tari AM *et al.* Bcl-2 inhibits p53 nuclear import following DNA damage. *Oncogene* 1997;**15**:2767-72.
22. Beham AW, Sarkiss M, Brisbay S, Tu SM, von Eschenbach AC, McDonnell TJ. Molecular correlates of bcl-2-enhanced growth following androgen-ablation in prostate carcinoma cells in vivo. *International Journal of Molecular Medicine* 1998;**1**:953-9.
23. Benchimol S, Pim D, Crawford L. Radioimmunoassay of the cellular protein p53 in mouse and human cell lines. *EMBO Journal* 1982;**1**:1055-62.
24. Bennett WP, Hollstein MC, He A, Zhu SM, Resau JH, Trump BF *et al.* Archival analysis of p53 genetic and protein alterations in Chinese esophageal cancer. *Oncogene* 1991;**6**:1779-84.
25. Bennett WP, Hollstein MC, Metcalf RA, Welsh JA, He A, Zhu SM *et al.* p53 mutation and protein accumulation during multistage human oesophageal carcinogenesis. *Cancer Research* 1992;**52**:6092-7.

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

26. Benson MC, Whang IS, Pantuck A, Ring K, Kaplan SA, Olsson CA *et al.* Prostate specific antigen density: a means of distinguishing benign prostatic hypertrophy and prostate cancer. *Journal of Urology* 1992;**147**:815-6.
27. Berchem GJ, Bosseler M, Sugars LY, Voeller HJ, Zeitlin S, Gelmann EP. Androgens induce resistance to bcl-2-mediated apoptosis in LNCaP prostate cancer cells. *Cancer Research* 1995;**55**:735-8.
28. Bermejo C, Pisters L. Cryotherapy for prostate cancer. *Expert Review of Anticancer therapy*. 2003;**3**:393-401.
29. Berner A, Geitvik G, Karlsen F, Fossa SD, Nesland JM, Borresen AL. TP53 mutations in prostatic cancer. Analysis of pre- and post-treatment archival formalin-fixed tumour tissue. *Journal of Pathology* 1995;**176**:299-308.
30. Bettencourt MC, Bauer JJ, Sesterhenn IA, Mostofi FK, McLeod DG, Moul JW. Ki-67 expression is a prognostic marker of prostate cancer recurrence after radical prostatectomy. *Journal of Urology* 1996;**156**:1064-8.
31. Bezzi M, Kressel HY, Allen KS, Schiebler ML, Altman HG, Wein AJ *et al.* Prostatic carcinoma: staging with MR imaging at 1.5 T. *Radiology* 1988;**169**:339-46.
32. Bhaskar V.S. KBV, Jennings TA, Ross JS, Breese K, Figge HL, Fisher HA *et al.* Alteration of the p53 locus in benign hyperplastic prostatic epithelium associated with high-grade prostatic adenocarcinoma. *Diagnostic Molecular Pathology* 1994;**3**:227-32.
33. Bill-Axelson A, Holmberg L, Ruutu M, Haggman M, Andersson SO, Spangberg A *et al.* Radical prostatectomy versus watchful waiting in early prostate cancer. *N Engl J Med* 2005;**352**:1977-84.
34. Blackwell KL, Bostwick DG, MR, Zincke H, Oesterling JE. Combining prostate specific antigen with cancer and gland volume to more reliably predict pathological stage: the influence of PSA-cancer

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

density. *Journal of Urology* 1994;**151**(6):565-70.

35. Blana A, Walter B, Rogenhoffer S, Wieland W. High-intensity focussed ultrasound for the treatment of localized prostate cancer: 5 year experience. *Urology* 2004;**63**:297-300.
36. Blasko JC. Brachytherapy. *Urology* 2000;**55**:306-8.
37. Bolla M, Collette L, Blank L, Warde P, Dubois J, Mirimanoff R-E *et al.* Long-term results with immediate androgen suppression and external irradiation in patients with locally advanced prostate cancer (an EORTC study): a phase III randomised trial. *Lancet* 2002;**360**:103-6.
38. Bolla M, Gonzalez D, Warde P, Dubois J, Mirimanoff R-E, Storme G *et al.* Improved survival in patients with locally advanced prostate cancer treated with radiotherapy and goserelin. *N Engl.J Med* 1997;**337**:295-300.
39. Bollens R, Vanden-Bossche M, Roumeguere T, Damoun A, Ekane S, Hoffman P *et al.* Extraperitoneal laparoscopic radical prostatectomy. Results after 50 cases. *European Urology* 2001;**40**:65-9.
40. Bonkhoff H. Role of the basal cells in premalignant changes of the human prostate: a stem cell concept for the development of prostate cancer. *European Urology* 1996;**30**:201-5.
41. Bonkhoff H, Fixemer T, Remberger K. Relation between Bcl-2, cell proliferation, and the androgen receptor status in prostate tissue and precursors of prostate cancer. *Prostate* 1998;**34**:251-8.
42. Bookstein R, MacGrogan D, Hilsenbeck SG, Sharkey F, Allred DC. p53 is mutated in a subset of advanced-stage prostate cancers. *Cancer Research* 1993;**53**:3369-73.
43. Borre M, Stausbol-Gron B, Overgaard J. p53 accumulation associated with bcl-2, the proliferation marker MIB-1 and survival in patients

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

with prostate cancer subjected to watchful waiting. *Journal of Urology* 2000;**164**:716-21.

44. Bostwick, D. G. Neoplasms of the Prostate. *Urologic Surgical Pathology* , 343-422. 1997.
45. Bostwick, D. G., Amin, M. B., and Dundore, P. Architectural patterns of high grade PIN. *Human Pathology* 24, 298-310. 1993.
46. Bostwick DG, Brawer MK, Anonymous, Richie JP, Kavoussi LR, Ho GT *et al.* Prostatic intra-epithelial neoplasia and early invasion in prostate cancer
Prostatic intraepithelial neoplasia: significance and correlation with prostate-specific antigen and transrectal ultrasound. Proceedings of a workshop of the National Prostate Cancer Detection Project. March 13, 1989, Bethesda, Maryland. *Cancer* 1987;**59**:788-94.
47. Bostwick DG, Dousa MK, Crawford BG, Wollan PC. Neuroendocrine differentiation in prostatic intraepithelial neoplasia and adenocarcinoma. *American Journal of Surgical Pathology* 1994;**18**:1240-6.
48. Bostwick DG, Qian J, Frankel K. The incidence of high grade prostatic intraepithelial neoplasia in needle biopsies. *Journal of Urology* 1995;**154**:1791-4.
49. Brasso K, Friis S, Juel K, Jorgensen T, Iversen P. Mortality of patients with clinically localized prostate cancer treated with observation for 10 years or longer: a population based registry study. *Journal of Urology* 1999;**161**:524-8.
50. Brawer MK. Prostatic intraepithelial neoplasia and prostate-specific antigen. *Urology* 1989;**34**:62-5.
51. Brawer MK, Aramburu EA, Chen GL, Preston SD, Ellis WJ. The inability of prostate specific antigen index to enhance the predictive the

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

value of prostate specific antigen in the diagnosis of prostatic carcinoma. *Journal of Urology* 1993;**150**:369-73.

52. Brawer MK, Lange PH. Prostate-specific antigen and premalignant change: implications for early detection. *Cancer* 1989;**39**:361-75.
53. Brawer MK, Rennels MA, Nagle RB, Schiffman R, Gaines JA. Serum prostate-specific antigen and prostate pathology in men having simple prostatectomy. *American Journal of Clinical Pathology* 1989;**92**:760-4.
54. Brenner PC, Rettig WJ, Sanz-Moncasi MP, Reuter V, Aprikian A, Old LJ *et al*. TAG-72 expression in primary, metastatic and hormonally treated prostate cancer as defined by monoclonal antibody CC49. *Journal of Urology* 1995;**153**:1575-9.
55. Brewster SF, Oxley JD, Trivella M, Abbott CD, Gillatt DA. Preoperative p53, bcl-2, CD44 and E-cadherin immunohistochemistry as predictors of biochemical relapse after radical prostatectomy. *Journal of Urology* 1999;**161**:1238-43.
56. Brooks JD, Bova GS, Ewing CM, Piantadosi S, Carter BS, Robinson JC *et al*. An uncertain role for p53 gene alterations in human prostate cancers. *Cancer Research* 1996;**56**:3814-22.
57. Brown DC, Gatter KC. Comparison of proliferation rates assessed using "multiblock" and conventional tissue blocks of lung carcinoma. *Journal of Clinical Pathology* 1992;**45**:579-82.
58. Bruckheimer EM, Brisbay S, Johnson DJ, Gingrich JR, Greenberg N, McDonnell TJ. Bcl-2 accelerates multistep prostate carcinogenesis in vivo. *Oncogene* 2000;**19**:5251-8.
59. Bryden AA, Freemont AJ, Clarke NW, George NJ. Ki-67 index in metastatic prostate cancer. *European Urology* 2001;**40**:673-6.
60. Bubendorf L, Sauter G, Moch H, Jordan P, Blochlinger A, Gasser TC

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

et al. Prognostic significance of Bcl-2 in clinically localized prostate cancer. *American Journal of Pathology* 1996;**148**:1557-65.

61. Bubendorf L, Sauter G, Moch H, Jordan P, Blochlinger A, Gasser TC *et al.* Prognostic significance of Bcl-2 in clinically localized prostate cancer. *American Journal of Pathology* 1996;**148**:1557-65.
62. Bubendorf L, Sauter G, Moch H, Schmid HP, Gasser TC, Jordan P *et al.* Ki67 labelling index: an independent predictor of progression in prostate cancer treated by radical prostatectomy. *Journal of Pathology* 1996;**178**:437-41.
63. Bubendorf L, Tapia C, Gasser TC, Casella R, Grunder B, Moch H *et al.* Ki67 labeling index in core needle biopsies independently predicts tumor-specific survival in prostate cancer. *Human Pathology* 1998;**29**:949-54.
64. Buckbinder L, Talbott R, Velasco-Miguel S, Takenaka I, Faha B, Seizinger BR *et al.* Induction of the growth inhibitor IGF-binding protein 3 by p53. *Nature* 1995;**377**:646-9.
65. Burton JL, Oakley N, Anderson JB. Recent advances in the histopathology and molecular biology of prostate cancer. *BJU Int.* 2000;**85**:87-94.
66. Byrne RL, Horne CH, Robinson MC, Autzen P, Apakama I, Bishop RI *et al.* The expression of waf-1, p53 and bcl-2 in prostatic adenocarcinoma. *British Journal of Urology* 1997;**79**:190-5.
67. Campo E, Miquel R, Jares P, Bosch F, Juan M, Leone A *et al.* Prognostic significance of the loss of heterozygosity of Nm23-H1 and p53 genes in human colorectal carcinomas. *Cancer* 1994;**73**:2913-21.
68. Caron de Fromentel C, May-Levin F, Mouriessse H, Lemerle J, Chandrasekaran K, May P. Presence of circulating antibodies against cellular protein p53 in a notable proportion of children with B-cell lymphoma. *International Journal of Cancer* 1987;**39**:185-9.

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

69. Carter HB, Pearson JD, Metter EJ, Brant LJ, Chan DW, Andres R *et al.* Longitudinal evaluation of prostate-specific antigen levels in men with and without prostate disease. *JAMA* 2002;**267**:2215-20.
70. Casson AG, Kerkvlelt N, O'Malley F. Prognostic value of p53 protein in esophageal adenocarcinoma. *Journal of Surgical Oncology* 1995;**60**:5-11.
71. Catalona WJ, Antenor J, Roehl K. Screening for prostate cancer in high risk populations. *J Urol* 2002;**168**:1980-4.
72. Catalona WJ, Smith DS, Ratliff TL. Measurement of prostate-specific antigen in serum as a screening test for prostate cancer. *New England Journal of Medicine* 2002;**324**:1156-61.
73. Cattoretti G, Becker MH, Key G, Duchrow M, Schluter C, Galle J *et al.* Monoclonal antibodies against recombinant parts of the Ki-67 antigen (MIB 1 and MIB 3) detect proliferating cells in microwave-processed formalin-fixed paraffin sections. *Journal of Pathology* 1992;**168**:357-63.
74. Cesinaro AM, Migaldi M, Ferrari G, Castagnetti G, Dotti A, De Gaetani C *et al.* Expression of p53 and bcl-2 in clinically localized prostate cancer before and after neo-adjuvant hormonal therapy. *Oncology Research* 2000;**12**:43-9.
75. Chang F, Syrjanen S, Kurvinen K, Syrjanen K. The p53 tumor suppressor gene as a common cellular target in human carcinogenesis. *American Journal of Gastroenterology* 1993;**88**:174-86.
76. Chaudhary KS, Abel PD, Lalani EN. Role of the Bcl-2 gene family in prostate cancer progression and its implications for therapeutic intervention. *Environmental Health Perspectives* 1999;**107** Suppl 1:49-57.
77. Cheng L, Cheville JC, Bostwick DG. Diagnosis of prostate cancer in needle biopsies after radiation therapy. *American Journal of Surgical Pathology* 1999;**23**:1173-83.

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

78. Cheng L, Leibovich BC, Bergstralh EJ, Scherer BG, Pacelli A, Ramnani DM *et al.* p53 alteration in regional lymph node metastases from prostate carcinoma: a marker for progression? *Cancer* 1999;**85**:2455-9.
79. Cheng L, Sebo TJ, Cheville JC, Pisansky TM, Slezak J, Bergstralh EJ *et al.* p53 protein overexpression is associated with increased cell proliferation in patients with locally recurrent prostate carcinoma after radiation therapy. *Cancer* 1999;**85**:1293-9.
80. Cheville JC, Reznicek MJ, Bostwick DG. The focus of "atypical glands, suspicious for malignancy" in prostatic needle biopsy specimens: incidence, histologic features, and clinical follow-up of cases diagnosed in a community practice. *American Journal of Clinical Pathology* 1997;**108**:633-40.
81. Chia SJ, Tang WY, Elnatan J, Yap WM, Goh HS, Smith DR. Prostate tumours from an Asian population: examination of bax, bcl-2, p53 and ras and identification of bax as a prognostic marker. *British Journal of Cancer* 2000;**83**:761-8.
82. Chodak GW, Thisted RA, Gerber GS, Johansson JE, Adolfsson J, Jones GW *et al.* Results of conservative management of clinically localized prostate cancer. *New England Journal of Medicine* 1994;**330**:242-8.
83. Chou JJ, Li H, Salvesen GS, Yuan J, Wagner G. Solution structure of BID, an intracellular amplifier of apoptotic signaling. *Cell* 1999;**96**:615-24.
84. Claudio PP, Zamparelli A, Garcia FU, Claudio L, Ammirati G, Farina A *et al.* Expression of cell-cycle-regulated proteins pRb2/p130, p107, p27(kip1), p53, mdm-2, and Ki-67 (MIB-1) in prostatic gland adenocarcinoma. *Clinical Cancer Research* 2002;**8**:1808-15.
85. Coffey, D. The molecular biology, endocrinology and physiology of the prostate and seminal vesicles. *Campbells Urology* Ed 6. 1992.

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

86. Collins GN, Lee RJ, McKelvie GB, Rogers AC, Hehir M. Relationship between prostate specific antigen, prostate volume and age in the benign prostate. *British Journal of Urology* 1993;**71**:445-50.
87. Collot T, Bass J, Denis F, Ranger R. Human p53 tumor suppressor p53 and DNA viruses. *Rev Med Virol* 2004;**14**:301-19.
88. Cooner WH, Mosley BR, Rutherford CLJr, Beard JH, Pond HS, Terry WJ *et al*. Prostate cancer detection in a clinical urological practice by ultrasonography, digital rectal examination and prostate specific antigen. *Journal of Urology* 2000; **143**:1146-54.
89. Cossman J, Schlegel R. p53 in the diagnosis of human neoplasia. *Journal of the National Cancer Institute* 1991;**83**:980-1.
90. Cowen D, Troncoso P, Khoo VS, Zagars GK, von Eschenbach AC, Meistrich ML *et al*. Ki-67 staining is an independent correlate of biochemical failure in prostate cancer treated with radiotherapy. *Clinical Cancer Research* 2002;**8**:1148-54.
91. Crawford LV, Pim DC, Bulbrook RD. Detection of antibodies against the cellular protein p53 in sera from patients with breast cancer. *International Journal of Cancer* 1982;**30**:403-8.
92. Crook T, Wrede D, Tidy JA, Mason WP, Evans DJ, Vousden KH. Clonal p53 mutation in primary cervical cancer: association with human papilloma virus negative tumours. *Lancet* 1992;**339**:1070-3.
93. D'Amico AV, Whittington R, Malkowicz SB, Schultz DS, Kaplan I, Beard CJ *et al*. *Journal of Clinical Oncology* 1998;**16**:3094-100.
94. D Kirk for the MRC Prostate Cancer Working Party and Investigators. Hormone therapy in advanced prostate cancer - report of the Medical Research Council "immediate" versus "deferred" treatment study. *British Journal of Urology* 1996;**77**:52.
95. Dahnert WF, Hamper UM, Eggleston JC, Walsh PC, Sanders RC. The

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

echogenic focus in prostatic sonograms, with xeroradiographic and histopathologic correlation. *Radiology* 1986;**159**:95-100.

96. Darson MF, Pacelli A, Roche P, Rittenhouse HG, Wolfert RL, Young CY *et al.* Human glandular kallikrein 2 (hK2) expression in prostatic intraepithelial neoplasia and adenocarcinoma: a novel prostate cancer marker. *Urology* 1997;**49**:857-62.
97. Davidoff A.M, Herndon JE, Glover NS, Kerns BJ, Pence JC, Iglehart JD *et al.* Relationship between p53 overexpression and established prognostic factors in breast cancer. *Surgery* 1991;**110**:259-64.
98. Davidoff A.M, Humphrey PA, Iglehart JD, Marks JR. Genetic basis for p53 overexpression in human breast cancer. *Proceedings of the National Academy of Sciences of the United States of America* 1991;**88**:5006-10.
99. de la Torre M, Haggman M, Brandstedt S, Busch C, Porter JR, Brawer MK *et al.* Prostatic intraepithelial neoplasia and invasive carcinoma in total prostatectomy specimens: distribution, volumes and DNA ploidy. *British Journal of Urology* 1993;**72**:207-13.
100. deVere White RW, Deitch AD, Jackson AG, Gandour-Edwards R, Marshalleck J, Soares SE *et al.* Racial differences in clinically localized prostate cancers of black and white men. *Journal of Urology* 1998;**159**:1979-82.
101. Dhanasekaran SM, Barrette TR, Ghosh D, Shah R, Varambally S, Kurachi K *et al.* Delineation of prognostic biomarkers in prostate cancer. *Nature* 2001;**412**:822-6.
102. Di Silverio F, Sciarra A, D'Eramo G, Casale P, Loreto A, Seccareccia F. Relationship among age, prostate-specific antigen, and prostate volume in men with lower urinary tract symptoms (LUTS) and in different groups of men with and without benign and malignant prostate diseases. *Prostate* 1998;**36**:1-7.
103. Dinjens WN, van der Weiden MM, Schroeder FH, Bosman FT,

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

Trapman J. Frequency and characterization of p53 mutations in primary and metastatic human prostate cancer. *International Journal of Cancer* 1994;**56**:630-3.

104. Djavan B, Remzi M, Seitz C, Anagnostou T, Dobrovits M, Harik M. A novel biopsy strategy defining the optimal number of cores based on PSA, age and prostate volume. *Journal of Urology* 2003;**169**:462 A.
105. Dolcetti R, Doglioni C, Maestro R, Gasparotto D, Barzan L, Pastore A *et al.* p53 over-expression is an early event in the development of human squamous-cell carcinoma of the larynx: genetic and prognostic implications. *International Journal of Cancer* 1992;**52**:178-82.
106. Donghi R, Longoni A, Pilotti S, Michieli P, Della Porta G, Pierotti MA. Gene p53 mutations are restricted to poorly differentiated and undifferentiated carcinomas of the thyroid gland. *Journal of Clinical Investigation*. 1993;**91**:1753-60.
107. Donovan J, Mills N, Smith M, Brindle L, Jacoby A, Peters T *et al.* Quality improvement report: Improving design and conduct of randomised trials by embedding them in qualitative research: ProtecT (prostate testing for cancer and treatment) study. Commentary: presenting unbiased information to patients can be difficult. *BMJ* 2002;**325**:766-70.
108. Drobnjak M, Latres E, Pollack D, Karpeh M, Dudas M, Woodruff JM *et al.* Prognostic implication of p53 nuclear over expression and high proliferation index of Ki-67 in adult soft tissue sarcomas. *Journal of the National Cancer Institute* 1994;**86**:549-54.
109. Dunsmuir WD, Gillett CE, Meyer LC, Young MP, Corbishley C, Eeles RA *et al.* Molecular markers for predicting prostate cancer stage and survival. *BJU Int.* 2000;**86**:869-78.
110. Eastham JA, Stapleton AM, Gousse AE, Timme TL, Yang G, Slawin KM *et al.* Association of p53 mutations with metastatic prostate cancer. *Clinical Cancer Research* 1995;**1**:1111-8.

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

111. Eeles RA, Dearnaley D, Arden-Jones A, Shearer R, Easton D, Ford D *et al.* Familial Prostate Cancer: The evidence and the Cancer Research Campaign/British Prostate Group (CRC/BPG) UK familial prostate cancer study. *BJU* 1997;**79** :8-14.
112. Effert PJ, Neubauer A, Walther PJ, Liu ET. Alterations of the P53 gene are associated with the progression of a human prostate carcinoma. *Journal of Urology* 1992;**147**:789-93.
113. Epstein JI., Carmichael MJ, Pivov G, Walsh PC. Influence of capsular penetration on progression following radical prostatectomy: a study of 196 cases with long-term followup. *Journal of Urology* 1993;**150**:135-41.
114. Fearon ER.,Vogelstein B. A genetic model for colorectal tumorigenesis. *Cell* 1990;**61**:759-67.
115. Feneley MR, Young MP, Chinyama C, Kirby RS, Parkinson MC. Ki-67 expression in early prostate cancer and associated pathological lesions. *Journal of Clinical Pathology* 1996;**49**:741-8.
116. Fernandez A, Udagawa T, Schwesinger C, Beecken W, Achilles-Gerte E, McDonnell T *et al.* Angiogenic potential of prostate carcinoma cells overexpressing bcl-2. *Journal of the National Cancer Institute* 2001;**93**:208-13.
117. Finnegan NM, Curtin JF, Prevost G, Morgan B, Cotter TG. Induction of apoptosis in prostate carcinoma cells by BH3 peptides which inhibit Bak/Bcl-2 interactions. *British Journal of Cancer* 2001;**85**:115-21.
118. Fonseca L, Yonemura Y, De Aretxabala X, Yamaguchi A, Miwa K, Miyazaki I. p53 detection as a prognostic factor in early gastric cancer. *Oncology* 1994;**51**:485-90.
119. Foti A, Ahuja HG, Allen SL, Koduru P, Schuster MW, Schulman P *et al.* Correlation between molecular and clinical events in the evolution of chronic myelocytic leukaemia to blast crisis. *Blood* 1991;**77**:2441-4.

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

120. Fowler JE, Jr., Bigler SA, Lynch C, Wilson SS, Farabaugh PB. Prospective study of correlations between biopsy-detected high grade prostatic intraepithelial neoplasia, serum prostate specific antigen concentration, and race. *Cancer* 2001;**91**:1291-6.
121. Fox SB, Persad RA, Royds J, Kore RN, Silcocks PB, Collins CC. p53 and c-myc expression in stage A1 prostatic adenocarcinoma: useful prognostic determinants? *Journal of Urology* 1993;**150**:490-4.
122. Furuya Y, Krajewski S, Epstein JI, Reed JC, Isaacs JT. Expression of bcl-2 and the progression of human and rodent prostatic cancers. *Clinical Cancer Research* 1996;**2**:389-98.
123. Gao X, Porter AT, Grignon DJ, Pontes JE, Honn KV. Diagnostic and prognostic markers for human prostate cancer. *Prostate* 1997;**31**:264-81.
124. Gaudin PB, Sesterhenn IA, Wojno KJ, Mostofi FK, Epstein JI. Incidence and clinical significance of high-grade prostatic intraepithelial neoplasia in TURP specimens. *Urology* 1997;**49**:558-63.
125. Gautschi O, Tschopp S, Olie RA, Leech SH, Simoes-Wust AP, Ziegler A *et al.* Activity of a novel bcl-2/bcl-xL-bispecific antisense oligonucleotide against tumors of diverse histologic origins. *Journal of the National Cancer Institute* 2001;**93**:463-71.
126. George NJ. Conservative management of prostate cancer. *Lancet* 1988;**1**:494-7.
127. Gerdes J. Ki-67 and other proliferation markers useful for immunohistological diagnostic and prognostic evaluations in human malignancies. *Seminars in Cancer Biology* 1990;**1**:199-206.
128. Gibbons RP, Correa TJ Jr, Brannen Ge, Weissman RM. Total prostatectomy for clinically localized prostatic cancer: long-term results. *Journal of Urology* 1989;**141**:564-6.

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

129. Gingrich JR, Barrios RJ, Kattan MW, Nahm HS, Finegold, MJ *et al.* Androgen-independent prostate cancer progression in the TRAMP model. *Cancer Research* 1997; **57**:4687-91.
130. Gingrich JR, Barrios RJ, Morton RA, Boyce BF, DeMayo FJ, Finegold MJ *et al.* Metastatic prostate cancer in a transgenic mouse. *Cancer Research* 1996; **56**:4096-102.
131. Gleason DF, Mellinger G. Prediction of prognosis for prostatic adenocarcinoma by combined histological and clinical staging. *Journal of Urology* 1974; **111**:58-64.
132. Gleave ME, Miyake H, Goldie J, Nelson C, Tolcher A. Targeting bcl-2 gene to delay androgen-independent progression and enhance chemosensitivity in prostate cancer using antisense bcl-2 oligodeoxynucleotides. *Urology* 1999; **54 (6A Suppl)**:36-46.
133. Glenister, TW. The development of the utricle and the so called "middle" or "median" lobe of the prostate. *Journal of Anatomy* 96, 443-455. 1962.
134. Gloushankova N, Ossovskaya V, Vasiliev J, Chumakov P, Kopnin B. Changes in p53 expression can modify cell shape of ras-transformed fibroblasts and epitheliocytes. *Oncogene* 1997; **15**:2985-9.
135. Gokden N, Roehl K, Catalona WJ, Humphrey PA. High-grade prostatic intraepithelial neoplasia in needle biopsy as a risk factor for the detection of adenocarcinoma: Current level of risk in screening population. *Urology* 2005; **65**:538-42.
136. Gould VE, Doljanskaia V, Gooch G, Bostwick DG. Immunolocalization of glycoprotein A-80 in prostatic carcinoma and prostatic intraepithelial neoplasia. *Human Pathology* 1996; **27**:547-52.
137. Graham SDjr and Bostwick, D. G. Report of the committee on staging and pathology. *Cancer* 70(suppl), 359-361. 1992.

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

138. Greenberg NM, DeMayo FJ, Finegold MJ, Medina D, Tilley WD, Aspinall JO *et al.* Prostate cancer in a transgenic mouse. *Proceedings of the National Academy of Sciences of the United States of America* 1995;**92**:3439-43.
139. Grignon DJ, Caplan R, Sarkar FH, Lawton CA, Hammond EH, Pilepich MV *et al.* p53 status and prognosis of locally advanced prostatic adenocarcinoma: a study based on RTOG 8610. *Journal of the National Cancer Institute* 1997;**89**:158-65.
140. Grimm P, Blasko JC, Sylvester J, Meier R, Cavanagh W. 10-year biochemical (prostate specific antigen) control of prostate cancer with (125) I brachytherapy. *Inte-J-Rad-Oncol-Biol-Phys* 2001;**51**:31-40.
141. Gross A, Yin XM, Wang K, Wei MC, Korsmeyer SJ. Enforced dimerization of BAX results in its translocation, mitochondrial dysfunction and apoptosis. *EMBO Journal* 1998;**17**:3878-85.
142. Grossfeld GD, Olumi AF, Connolly JA, Chew K, Gibney J, Bhargava V *et al.* Locally recurrent prostate tumors following either radiation therapy or radical prostatectomy have changes in Ki-67 labeling index, p53 and bcl-2 immunoreactivity. *Journal of Urology* 1998;**159**:1437-43.
143. Guillonnet B, Rozet F, Cathelineau X, Lay F, Barret E, Doublet JD *et al.* Perioperative complications of laparoscopic radical prostatectomy: the Montsouris 3-year experience. *J Urol* 2002;**167**:51-6.
144. Guillonnet B, el-Fettouh H, Baumert H, Cathelineau X, Doublet JD, Fromont G *et al.* Laparoscopic radical prostatectomy: oncological evaluation after 1,000 cases at Montsouris Institute. *J Urol* 2003;**169**:1261-6.
145. Guillonnet B, Vallencien G. Laparoscopic radical prostatectomy: the Montsouris experience. *J Urol* 2000;**163**:418-22.
146. Gusterson BA, Anbazhagan R, Warren W, Midgely C, Lane DP, O'Hare M *et al.* Expression of p53 in premalignant and malignant

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

squamous epithelium. *Oncogene* 1991;**6**:1785-9.

147. Haapasalo H, Isola J, Sallinen P, Kalimo H, Helin H, Rantala I. Aberrant p53 expression in astrocytic neoplasms of the brain: association with proliferation. *American Journal of Pathology* 1993;**142**:1347-51.
148. Haldar S, Basu A, Croce CM. Bcl2 is the guardian of microtubule integrity. *Cancer Research* 1997;**57**:229-33.
149. Haldar S, Chintapalli J, Croce CM. Taxol induces bcl-2 phosphorylation and death of prostate cancer cells. *Cancer Research* 1996;**56**:1253-5.
150. Haldar S, Jena N, Croce CM. Inactivation of Bcl-2 by phosphorylation. *Proceedings of the National Academy of Sciences of the United States of America* 1995;**92**:4507-11.
151. Hall MC, Navone NM, Troncoso P, Pollack A, Zagars GK, von Eschenbach AC *et al.* Frequency and characterization of p53 mutations in clinically localized prostate cancer. *Urology* 1995;**45**:470-5.
152. Hall PA. Cell proliferation. *Journal of Pathology* 1991;**165**:349-54.
153. Hall PA, Lane DP. p53 in tumour pathology: can we trust immunohistochemistry?--Revisited! *Journal of Pathology* 1994;**172**:1-4.
154. Hall PA, Levison DA, Woods AL, Yu CC, Kellock DB, Watkins JA *et al.* Proliferating cell nuclear antigen (PCNA) immunolocalization in paraffin sections: an index of cell proliferation with evidence of deregulated expression in some neoplasms. *Journal of Pathology* 1990;**162**:285-94.
155. Halvorsen OJ, Haukaas S, Hoisaeter PA, Akslen LA. Maximum Ki-67 staining in prostate cancer provides independent prognostic information after radical prostatectomy. *Anticancer Research*

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

2001;**21**:4071-6.

156. Han JP, Sabbatini P, Perez D, Rao L, Modha D, White E. The E1B 19K protein blocks apoptosis by interacting with and inhibiting p53-inducible and death promoting Bax protein. *Genes Dev* 1996;**10**:461-77.
157. Hardie C, Parker C, Norman A, Eeles RA, Horwich A, Huddart R *et al*. Early outcomes of active surveillance for localised prostate cancer. *BJU Int*. 2005;**95**:956-60.
158. Harper ME, Glynn-Jones E, Goddard L, Mathews P, Nicholson RI. Expression of androgen receptor and growth factors in premalignant lesions of the prostate. *Journal of Pathology* 1998;**186**:169-77.
159. Harper ME, Goddard L, Wilson DW, Matanhelia SS, Conn IG, Peeling WB *et al*. Pathological and clinical associations of Ki-67 defined growth fractions in human prostatic carcinoma. *Prostate* 1992;**21**:75-84.
160. Haussler O, Epstein JI, Amin MB, Heitz PU, Hailemariam S. Cell proliferation, apoptosis, oncogene, and tumor suppressor gene status in adenosis with comparison to benign prostatic hyperplasia, prostatic intraepithelial neoplasia, and cancer. *Human Pathology* 1999;**30**:1077-86.
161. Heidenberg HB, Sesterhenn IA, Gaddipati JP, Weghorst CM, Buzard GS, Moul *et al*. Alteration of the tumor suppressor gene p53 in a high fraction of hormone refractory prostate cancer. *Journal of Urology* 1995;**154**:414-21.
162. Henke RP, Kruger E, Ayhan N, Hubner D, Hammerer P, Huland H. Immunohistochemical detection of p53 protein in human prostatic cancer. *Journal of Urology* 1994;**152**:1297-301.
163. Hockenbery D, Nunez G, Millman C, Schreiber RD, Korsmeyer SJ. Bcl-2 is an inner mitochondrial membrane protein that blocks programmed cell death. *Nature* 1990;**348**:334-6.

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

164. Hodge KK, Mcneal JE, Stamey TA. Ultrasound guided transrectal core biopsies of the palpably abnormal prostate. *Journal of Urology* 1989;66-70.
165. Hodge KK, Mcneal JE, Terris MK, Stamey TA. Random systematic versus directed ultrasound guided transrectal core biopsies of the prostate. *Journal of Urology* 1989;142:71-5.
166. Hoedemaeker RF, Kranse R, Rietbergen JB, Kruger AE, Schroder FH, van der Kwast TH. Evaluation of prostate needle biopsies in a population-based screening study: the impact of borderline lesions. *Cancer* 1999;85:145-52.
167. Hoisaeter PA, Norlen BJ, Norming U, Ebert T, Matre R, Nilsson S *et al.* Imaging in the diagnosis and assessment of prognosis in localized prostate cancer. Consensus Conference on Diagnosis and Prognostic Parameters in Localized Prostate Cancer. Stockholm, Sweden, May 12-13, 1993. *Scandinavian Journal of Urology & Nephrology* 1993;Suppl:89-106.
168. Hollstein M, Sidransky D, Vogelstein B, Harris CC. p53 mutations in human cancers. *Science* 1991;253:49-53.
169. Holmberg L, Bill-Axelson A, Helgesen F, Salo JO, Folmerz P, Haggman *et al.* A randomized trial comparing radical prostatectomy with watchful waiting in early prostate cancer. *N Engl J Med* 2002;347:781-9.
170. Horninger W, Voigger H, Rogatsch H, Strohmeyer D, Steiner H, Hobisch A *et al.* Predictive value of total and percent free prostate specific antigen in high grade prostatic intraepithelial neoplasia lesions: results of the Tyrol Prostate Specific Antigen Screening Project. *Journal of Urology* 2001;165:1143-5.
171. Howley PM. Role of the human papillomaviruses in human cancer. *Cancer Research* 1991;51:5019s-22s.
172. Hu JC, Palapattu GS, Kattan MW, Scardino PT, Wheeler TM. The

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

association of selected pathological features with prostate cancer in a single-needle biopsy accession. *Human Pathology* 1998;**29**:1536-8.

173. Huang A, Gandour-Edwards R, Rosenthal SA, Siders DB, Deitch AD, White RW. p53 and bcl-2 immunohistochemical alterations in prostate cancer treated with radiation therapy. *Urology* 1998;**51**:346-51.
174. Hughes JH, Cohen MB, Robinson RA. p53 immunoreactivity in primary and metastatic prostatic adenocarcinoma. *Modern Pathology* 1995;**8**:462-6.
175. Humphrey PA, Swanson PE. Immunoreactive p53 protein in high-grade prostatic intraepithelial neoplasia. *Pathology, Research & Practice* 1995;**191**:881-7.
176. Incognito LS, Cazares LH, Schellhammer PF, Kuban DA, Van Dyk EO, Moriarty RP *et al*. Overexpression of p53 in prostate carcinoma is associated with improved overall survival but not predictive of response to radiotherapy. *International Journal of Oncology* 2000;**17**:761-9.
177. Inoue T., Segawa T, Shiraishi T, Yoshida Y, Toda Y, Yamada T *et al*. Androgen receptor, Ki67, and p53 expression in radical prostatectomy specimens predicts treatment failure in Japanese population. *Urology* 2005;**66**:332-7.
178. Ionov Y, Yamamoto H, Krajewski S, Reed JC, Perucho M. Mutational inactivation of the proapoptotic gene BAX confers selective advantage during tumor clonal evolution. *Proceedings of the National Academy of Sciences of the United States of America* 2000;**97**:10872-7.
179. Isaacs WB. Molecular genetics of prostate cancer. *Cancer Surveys* 1995;**25**:357-79.
180. Isaacs WB, Carter BS, Ewing CM. Wild-type p53 suppresses growth of human prostate cancer cells containing mutant p53 alleles. *Cancer Research* 1991;**51**:4716-20.

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

181. Isobe T, Hiyama K, Yoshida Y, Fujiwara Y, Yamakido M. Prognostic significance of p53 and ras gene abnormalities in lung adenocarcinoma patients with stage 1 disease after curative resection. *Japanese Journal of Cancer Research*. 1994;**85**:1240-6.
182. Isola J, Visakorpi T, Holli K, Kallioniemi OP. Association of over expression of tumor suppressor protein p53 with rapid cell proliferation and poor prognosis in node-negative breast cancer patients. *Journal of the National Cancer Institute* 1992;**84** :1109-14.
183. Ito T, Seyama T, Mizuno T, Tsuyama N, Hayashi T, Hayashi Y *et al*. Unique association of p53 mutations with undifferentiated but not differentiated carcinomas of the thyroid gland. *Cancer Research* 1992;**52**:1369-71.
184. Johansson JE, Adami HO, Andersson SO, Bergstrom R, Krusemo UB, Kraaz W. Natural history of localised prostatic cancer. A population-based study in 223 untreated patients. *Lancet* 1989;**1**:799-803.
185. Johnson MI, Hamdy FC. Apoptosis regulating genes in prostate cancer *Oncology Reports* 1998;**5**:553-7.
186. Johnson MI, Hamdy FC. Apoptosis regulating genes in prostate cancer. *Oncology Reports* 1998;**5**:553-7.
187. Johnson MI, Robinson MC, Marsh C, Robson CN, Neal DE, Hamdy FC. Expression of Bcl-2, Bax, and p53 in high-grade prostatic intraepithelial neoplasia and localized prostate cancer: relationship with apoptosis and proliferation. *Prostate* 1998;**37**:223-9.
188. Kabalin JN, Mcneal JE, Johnstone IM, Stamey TA. Serum prostate specific antigen and the biologic progression of prostate cancer. *Urology* 1995;**46**:65-70.
189. Kakeji Y, Korenaga D, Tsujitani S, Baba H, Anai H, Maehara Y *et al*. Gastric cancer with p53 over expression has a high potential for metastasising to lymph nodes. *British Journal of Cancer* 1993;**67**:589-93.

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

190. Kaklamanis L, Gatter KC, Mortensen N, Baigrie RJ, Heryet A, Lane DP *et al.* p53 expression in colorectal adenomas. *American Journal of Pathology* 1993;**142**:87-93.
191. Kallakury BV, Figge J, Leibovich B, Hwang J, Rifkin M, Kaufman R *et al.* Increased bcl-2 protein levels in prostatic adenocarcinomas are not associated with rearrangements in the 2.8 kb major breakpoint region or with p53 protein accumulation. *Modern Pathology* 1996;**9**:41-7.
192. Keetch DW, Humphrey P, Stahl D, Smith DS, Catalona WJ. Morphometric analysis and clinical followup of isolated prostatic intraepithelial neoplasia in needle biopsy of the prostate. *Journal of Urology* 1995;**154**:347-51.
193. Kelavkar P, Cohen C, Kamitani H, Eling E, Badr F. Concordant induction of 15-lipoxygenase-1 and mutant p53 expression in human prostate adenocarcinoma: correlation with Gleason staging. *Carcinogenesis* 2000;**21**:1777-87.
194. Keshgegian AA, Johnston E, Cnaan A. Bcl-2 oncoprotein positivity and high MIB-1 (Ki-67) proliferative rate are independent predictive markers for recurrence in prostate carcinoma. *American Journal of Clinical Pathology* 1998;**110**:443-9.
195. Key G, Becker MH, Baron B, Duchrow M, Schluter C, Flad HD *et al.* New Ki-67-equivalent murine monoclonal antibodies (MIB 1-3) generated against bacterially expressed parts of the Ki-67 cDNA containing three 62 base pair repetitive elements encoding for the Ki-67 epitope. *Laboratory Investigation* 1993;**68**:629-36.
196. Key G, Petersen JL, Becker MH, Duchrow M, Schluter C, Askaa J *et al.* New antiserum against Ki-67 antigen suitable for double immunostaining of paraffin wax sections. *Journal of Clinical Pathology* 1993;**46**:1080-4.
197. Khoo VS, Pollack A, Cowen D, Joon DL, Patel N, Terry NH *et al.* Relationship of Ki-67 labeling index to DNA-ploidy, S-phase fraction, and outcome in prostate cancer treated with radiotherapy. *Prostate*

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

1999;**41**:166-72.

198. Kilic S, Kukul E, Danisman A, Guntekin E, Sevuk M. Ratio of free to total prostate-specific antigen in patients with prostatic intraepithelial neoplasia. *European Urology* 1998;**34**:176-80.
199. Kimura K, Markowski M, Bowen C, Gelmann EP. Androgen blocks apoptosis of hormone-dependent prostate cancer cells. *Cancer Research* 2001;**61**:5611-8.
200. Koch M, de Miguel M, Hofler H, Diaz-Cano SJ. Kinetic profiles of intraepithelial and invasive prostatic neoplasias: the key role of down-regulated apoptosis in tumor progression. *Virchows Archiv* 2000;**436**:413-20.
201. Koivisto PA, Rantala I. Amplification of the androgen receptor gene is associated with P53 mutation in hormone-refractory recurrent prostate cancer. *Journal of Pathology* 1999;**187**:237-41.
202. Korsmeyer SJ. Bcl-2 initiates a new category of oncogenes: regulators of cell death. *Blood* 1992;**80**:879-86.
203. Koshland DE, Jr. Molecule of the year. *Science* 1993;**262**:1953.
204. Krajewska M, Krajewski S, Epstein JI, Shabaik A, Sauvageot J, Song K *et al*. Immunohistochemical analysis of bcl-2, bax, bcl-X, and mcl-1 expression in prostate cancers. *American Journal of Pathology* 1996;**148**:1567-76.
205. Kranse R, Beemsterboer P, Rietbergen J, Habbema D, Hugosson J, Schroder FH. Predictors for biopsy outcome in the European Randomized Study of Screening for Prostate Cancer (Rotterdam region). *Prostate* 1999;**39**:316-22.
206. Krongrad A, Lai H, Lamm SH, Lai S. Mortality in prostate cancer. *Journal of Urology* 1996;**156**:1084-91.

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

207. Kronz JD, Allan CH, Shaikh AA, Epstein JI. Predicting cancer following a diagnosis of high-grade prostatic intraepithelial neoplasia on needle biopsy: data on men with more than one follow-up biopsy. *American Journal of Surgical Pathology* 2001;**25**:1079-85.
208. Kubbutat MH, Key G, Duchrow M, Schluter C, Flad HD, Gerdes J. Epitope analysis of antibodies recognising the cell proliferation associated nuclear antigen previously defined by the antibody Ki-67 (Ki-67 protein). *Journal of Clinical Pathology* 1994;**47**:524-8.
209. Kuczyk MA, Serth J, Bokemeyer C, Machtens S, Minssen A, Bathke W *et al.* The prognostic value of p53 for long-term and recurrence-free survival following radical prostatectomy. *European Journal of Cancer* 1998;**34**:679-86.
210. Kunimi K, Amano T, Uchibayashi T. Point mutation of the p53 gene is an infrequent event in untreated prostate cancer. *Cancer Detection & Prevention* 1996;**20**:218-22.
211. Kyprianou N, King ED, Bradbury D, Rhee JG. bcl-2 over-expression delays radiation-induced apoptosis without affecting the clonogenic survival of human prostate cancer cells. *International Journal of Cancer* 1997;**70**:341-8.
212. Labrecque S, Naor N, Thomson D, Matlashewski G. Analysis of anti-p53 antibody response in cancer patients. *Cancer Research* 1993;**53**:3468-71.
213. Lane DP. Cancer. p53, guardian of the genome. [letter; comment]. [see comments]. *Nature* 1992;**358**:15-6.
214. Lane DP, Crawford LV. T antigen is bound to a host protein in SV40-transformed cells. *Nature* 1979;**278**:261-3.
215. Lang C, Unteregger G, Kartarius S, Gunther J, Bonkhoff H, Montenarh M *et al.* p53 autoantibodies in patients with urological tumours. *British Journal of Urology* 1998;**82**:721-6.

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

216. Langer JE, Rovner ES, Coleman BG, Yin D, Arger PH, Malkowicz SB *et al.* Strategy for repeat biopsy of patients with prostatic intraepithelial neoplasia detected by prostate needle biopsy. *Journal of Urology* 1996;**155**:228-31.
217. Lebedeva I, Rando R, Ojwang J, Cossum P, Stein CA. Bcl-xL in prostate cancer cells: effects of overexpression and down-regulation on chemosensitivity. *Cancer Research* 2000;**60**:6052-60.
218. Lee F, Torp-Pedersen ST, Carroll JT, Siders DB, Christensen-Day C, Mitchell AE. Use of transrectal ultrasound and prostate-specific antigen in diagnosis of prostatic intraepithelial neoplasia. *Urology* 1989;**34**:4-8.
219. Legros Y, Lacabanne V, d'Agay MF, Larsen CJ, Pla M, Soussi T. Production of human p53 specific monoclonal antibodies and their use in immunohistochemical studies of tumor cells. *Bulletin du Cancer*. 1993;**80**:102-10.
220. Leibovich BC, Cheng L, Weaver AL, Myers RP, Bostwick DG. Outcome prediction with p53 immunostaining after radical prostatectomy in patients with locally advanced prostate cancer. *Journal of Urology* 2000;**163**:1756-60.
221. Lentle BC, McGowan DG, Dierich H. Technetium-99M polyphosphate bone scanning in carcinoma of the prostate. *BJU* 1994;**46**:543-8.
222. Leung S, Miyake H, Zellweger T, Tolcher A, Gleave ME. Synergistic chemosensitization and inhibition of progression to androgen independence by antisense Bcl-2 oligodeoxynucleotide and paclitaxel in the LNCaP prostate tumor model. *International Journal of Cancer* 2001;**91**:846-50.
223. Levesque MA, Katsaros D, Yu H, Zola P, Sismondi P, Giardina G *et al.* Mutant p53 over expression is associated with poor outcome in patients with well or moderately differentiated ovarian carcinoma. *Cancer* 1995;**75**:1327-38.

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

224. Levine AJ, Momand J, Finlay CA. The p53 tumour suppressor gene. *Nature* 1991;**351**:453-6.
225. Li R, Heydon K, Hammond EH, Grignon DJ, Roach M, Wolkov H *et al.* Ki-67 stainin index predicts metastasis and survival in locally advanced prostate cancer treated with radiotherapy: An analysis of patients in radiation therapy oncology group protocol 86-10. *Clinical Cancer Research* 2004;**10**:4118-24.
226. Li X, Marani M, Yu J, Nan B, Roth JA, Kagawa S *et al.* Adenovirus-mediated Bax overexpression for the induction of therapeutic apoptosis in prostate cancer. *Cancer Research* 2001;**61**:186-91.
227. Lipponen P, Vesalainen S. Expression of the apoptosis suppressing protein bcl-2 in prostatic adenocarcinoma is related to tumor malignancy. *Prostate* 1997;**32**:9-15.
228. Ljung G, Egevad L, Norberg M, de la TM, Holmberg L, Busch C *et al.* Assessment of proliferation indicators in residual prostatic adenocarcinoma cells after radical external beam radiotherapy. *Prostate* 1996;**29**:303-10.
229. Lloyd SN, Brown IL, Leake RE. Ki-67 antibody immunostaining in benign and malignant human prostatic disease. *International Journal of Biological Markers* 1992;**7**:256-9.
230. Lowe SL, Rubinchik S, Honda T, McDonnell TJ, Dong JY, Norris JS. Prostate-specific expression of Bax delivered by an adenoviral vector induces apoptosis in LNCaP prostate cancer cells. *Gene Therapy* 2001;**8**:1363-71.
231. Lowsley O. *Am J Surg* 1930;**8**:526-41.
232. Lu-Yao GL, Albertsen PC, tanford JL, tukel TA, alker-Corkery ES, arry MJ. Natural experiment examining impact of aggressive screening and treatment on prostate cancer mortality in two fixed cohorts from Seattle area and Connecticut. *BMJ* 2002; **325**:740.

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

233. Lu-Yao GL, Yao SL. Population-based study of long-term survival in patients with clinically localised prostate cancer. [see comments.]. *Lancet* 1997;**349**:906-10.
234. Lunaceck A, Schwentner C, Fritsch H, Bartsch G, Strasser H. Anatomical radical retropubic prostatectomy: 'curtain dissection' of the neurovascular bundles. *BJU Int.* 2005;**95**:1226-31.
235. Mackey TJ, Borkowski A, Amin P, Jacobs SC, Kyprianou N. bcl-2/bax ratio as a predictive marker for therapeutic response to radiotherapy in patients with prostate cancer. *Urology* 1998;**52**:1085-90.
236. Macoska JA, Powell IJ, Sakr W, Lane MA. Loss of the 17p chromosomal region in a metastatic carcinoma of the prostate. *Journal of Urology* 1992;**147**:1142-6.
237. Mariappan P, Chong W, Sundaram M, Mohamed S. Increasing prostate biopsy cores based on volume vs the sextant biopsy: a prospective randomized controlled clinical study on cancer detection rates and morbidity. *BJU Int.* 2004; **94**.(3), P.: 307.-10.
238. Masuda M, Takano Y, Iki M, Asakura T, Hashiba T, Noguchi S *et al.* Prognostic significance of Ki-67, p53, and Bcl-2 expression in prostate cancer patients with lymph node metastases: a retrospective immunohistochemical analysis. *Pathology International* 1998;**48**:41-6.
239. Matsushima H, Kitamura T, Goto T, Hosaka Y, Homma Y, Kawabe K. Combined analysis with Bcl-2 and P53 immunostaining predicts poorer prognosis in prostatic carcinoma. *Journal of Urology* 1997;**158**:2278-83.
240. Matsushima H, Sasaki T, Goto T, Hosaka Y, Homma Y, Kitamura T *et al.* Immunohistochemical study of p21WAF1 and p53 proteins in prostatic cancer and their prognostic significance. *Human Pathology* 1998;**29**:778-83.
241. Matsuura H, Hayashi N, Kawamura J, Shiraishi T, Yatani R. Prognostic significance of Ki-67 expression in advanced prostate

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

cancers in relation to disease progression after androgen ablation. *European Urology* 2000;**37**:212-7.

242. McCarthy P, Pollack H. Imaging of patients with stage D prostatic carcinoma. *Urol Clin N Am* 1991;**18**:35-53.
243. McClennan BL. Transrectal US of the prostate: is the technology leading the science? *Radiology* 1988;**168**:571-5.
244. McCormick D, Chong H, Hobbs C, Datta C, Hall PA. Detection of the Ki-67 antigen in fixed and wax-embedded sections with the monoclonal antibody MIB1. *Histopathology* 1993;**22**:355-60.
245. McDonnell TJ, Navone NM, Troncoso P, Pisters LL, Conti C, von Eschenbach AC *et al*. Expression of bcl-2 oncoprotein and p53 protein accumulation in bone marrow metastases of androgen independent prostate cancer. *Journal of Urology* 1997;**157**:569-74.
246. McDonnell TJ, Troncoso P, Brisbay SM, Logothetis C, Chung LW, Hsieh JT *et al*. Expression of the protooncogene bcl-2 in the prostate and its association with emergence of androgen-independent prostate cancer. *Cancer Research* 1992;**52**:6940-4.
247. McLoughlin J, Foster CS, Price P, Williams G, Abel PD. Evaluation of Ki-67 monoclonal antibody as prognostic indicator for prostatic carcinoma. *British Journal of Urology* 1993;**72**:92-7.
248. McNeal, J. E. Normal histology of the prostate. *American journal of surgical pathology* 1988;**12**, 619- 633.
249. McNeal JE, Alroy J, Leav I, Redwine EA, Freiha FS, Stamey T. Immunohistochemical evidence for impaired cell differentiation in the premalignant phase of prostate carcinogenesis. *American Journal of Clinical Pathology* 1988;**90**:23-32.
250. Mellinger, GT, Gleason, D. F., and Bailar, J. The histology and

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

prognosis of prostate cancer. *Journal of Urology* 1967;97(2), 331-337.

251. Mellon K, Thompson S, Charlton RG, Marsh C, Robinson M, Lane DP *et al.* p53, c-erbB-2 and the epidermal growth factor receptor in the benign and malignant prostate. *Journal of Urology* 1992;**147**:496-9.
252. Merrill R, Hilton S, Wiggins C, Sturgeon J. Towards a better understanding of the comparatively high prostate cancer incidence rates in Utah. *BMC Cancer* 2003;**3**:14.
253. Mettlin C, Lee F, Drago J, Murphy GP, Skjorten FJ, Berner A *et al.* The American Cancer Society National Prostate Cancer Detection Project. Findings on the detection of early prostate cancer in 2425 men Prostatic intraepithelial neoplasia in surgical resections: relationship to coexistent adenocarcinoma and atypical adenomatous hyperplasia of the prostate. *Cancer* 1991;**67**:2949-58.
254. Meyers FJ, Gumerlock PH, Chi SG, Borchers H, Deitch AD, deVere White RW. Very frequent p53 mutations in metastatic prostate carcinoma and in matched primary tumors. *Cancer* 1998;**83**:2534-9.
255. Miyake H, Tolcher A, Gleave ME. Antisense Bcl-2 oligodeoxynucleotides inhibit progression to androgen-independence after castration in the Shionogi tumor model. *Cancer Research* 1999;**59**:4030-4.
256. Miyake H, Tolcher A, Gleave ME. Chemosensitization and delayed androgen-independent recurrence of prostate cancer with the use of antisense Bcl-2 oligodeoxynucleotides. *Journal of the National Cancer Institute* 2000;**92**:34-41.
257. Middleton RG, Thompson IM, Astenfeld M, Cooner W, Correa RJ, Gibbons RP *et al.* Prostate Cancer Clinical Guidelines Panel Summary report on the management of clinically localized prostate cancer. The American Urological Association. *Journal of Urology* 1995;**154**:2144-8.

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

258. Minardi D, Galosi AB, Dell'Atti L, Hanitzsch H, Mario P, Muzzonigro G. Production of serum-free and total prostate-specific antigen due to prostatic intraepithelial neoplasia. *Scandinavian Journal of Urology & Nephrology* 2002;**36**:323-9.
259. Mirchandani D, Zheng J, Miller GJ, Ghosh AK, Shibata DK, Cote RJ *et al.* Heterogeneity in intratumor distribution of p53 mutations in human prostate cancer. *American Journal of Pathology* 1995;**147**:92-101.
260. Mirtti T, Kallajoki M, Aaltonen M, Alanen K. Cyclin A and Ki-67 with DNA content in benign and malignant prostatic epithelial lesions. *Analytical & Quantitative Cytology & Histology* 2001;**23**:229-37.
261. Miyashita T, Reed JC. Tumor suppressor p53 is a direct transcriptional activator of the human bax gene. *Cell* 1995;**80**:293-9.
262. Momand J, Zambetti GP, Olson DC, George D, Levine AJ. The mdm-2 oncogene product forms a complex with the p53 protein and inhibits p53-mediated transactivation. *Cell* 1992;**69**:1237-45.
263. Montironi R, Bostwick DG, Bonkhoff H, Cockett AT, Helpap B, Troncoso *et al.* Origins of prostate cancer. *Cancer* 1996;**78**:362-5.
264. Montironi R, Diamanti L, Thompson D, Bartels HG, Bartels PH. Analysis of the capillary architecture in the precursors of prostate cancer: recent findings and new concepts. *European Urology* 1996;**30**:191-200.
265. Montironi R, Galluzzi CM, Diamanti L, Giannulis I, Pisani E, Scarpelli M. Prostatic intra-epithelial neoplasia: expression and location of proliferating cell nuclear antigen in epithelial, endothelial and stromal nuclei. *Virchows Archiv - A, Pathological Anatomy & Histopathology* 1992;**422**:185-92.
266. Montironi R, Magi-Galluzzi C, Scarpelli M, Giannulis I, Diamanti L. Occurrence of cell death (apoptosis) in prostatic intra-epithelial neoplasia. *Virchows Archiv - A, Pathological Anatomy &*

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

Histopathology 1993;**423**:351-7.

267. Moore CK, Karikehalli S, Nazeer T, Fisher HA, Kaufman R, Mian B. Prognostic significance of high grade prostatic intraepithelial neoplasia and atypical small acinar proliferation in the contemporary era. *Journal of Urology*: 2005;**173**:70-2.
268. Morkove O, Halvorsen OJ, Skjaerven R, Stangeland L, Gulsvik A, Laerum OD. Prognostic significance of p53 protein expression and DNA ploidy in surgically treated non-small cell lung carcinomas. *Anticancer Research* 1993;**13**:571-8.
269. Morote J, Raventos CX, Encabo G, Lopez M, e Torres IM. Effect of high-grade prostatic intraepithelial neoplasia on total and percent free serum prostatic-specific antigen. *European Urology*. 2000;**37**:456-9.
270. Morote J, Encabo G, Lopez M, de Torres IM. Influence of high-grade prostatic intra-epithelial neoplasia on total and percentage free serum prostatic specific antigen. *BJU Int*. 1999;**84**:657-60.
271. Morote J, Encabo G, Lopez M, de Torres IM. Prediction of prostate volume based on total and free serum prostate-specific antigen: is it reliable? *European Urology* 2000;**38**:91-5.
272. Morrin M, Kelly M, Barrett N, Delaney P. Mutations of Ki-ras and p53 genes in colorectal cancer and their prognostic significance. *Gut* 1994;**35**:1627-31.
273. Mottaz AE, Markwalder R, Fey MF, Klima I, Merz VW, Thalmann GN *et al*. Abnormal p53 expression is rare in clinically localized human prostate cancer: comparison between immunohistochemical and molecular detection of p53 mutations. *Prostate* 1997;**31**:209-15.
274. Moul JW, Bettencourt MC, Sesterhenn IA, Mostofi FK, McLeod DG, Srivastava S *et al*. Protein expression of p53, bcl-2, and KI-67 (MIB-1) as prognostic biomarkers in patients with surgically treated, clinically localized prostate cancer. *Surgery* 1996;**120**:159-66.

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

275. Mucci NR, Rubin MA, Strawderman MS, Montie JE, Smith DC, Pienta KJ. Expression of nuclear antigen Ki-67 in prostate cancer needle biopsy and radical prostatectomy specimens. *Journal of the National Cancer Institute* 2000;**92**:1941-2.
276. Myers RB, Grizzle WE. Biomarker expression in prostatic intraepithelial neoplasia. *European Urology* 1996;**30**:153-66.
277. Myers RB, Grizzle WE. Changes in biomarker expression in the development of prostatic adenocarcinoma. [Review] [73 refs]. *Biotechnic & Histochemistry* 1997;**72**:86-95.
278. Myers RB, Oelschlager D, Srivastava S, Grizzle WE. Accumulation of the p53 protein occurs more frequently in metastatic than in localized prostatic adenocarcinomas. *Prostate* 1994; **25**:243-8.
279. Nagata S. Apoptosis by death factor. *Cell* 1997;**88**:355-65.
280. Nakase M, Inui M, Okumura K, Kamei T, Nakamura S, Tagawa T. p53 gene therapy of human osteosarcoma using a transferrin modified cationic liposome. *Mol-Cancer-Ther* 2005;**4**:625-31.
281. Narayan P, Fournier G, Gajendran V, Leidich R, Lo R, Jacob G *et al*. Utility of preoperative serum prostate-specific antigen concentration and biopsy Gleason score in predicting risk of pelvic lymph node metastases in prostate cancer. *Urology* 1994;**44**:519-24.
282. National Institute for Clinical Excellence. Guidance on Cancer Services: Improving Outcomes in Urological Cancers. The Manual. www.nice.org.uk 2002.
283. Navone NM, Troncoso P, Pisters LL, Goodrow TL, Palmer JL, Nichols WW *et al*. p53 protein accumulation and gene mutation in the progression of human prostate carcinoma. *Journal of the National Cancer Institute* 1993;**85**:1657-69.
284. Naya Y, Ayala AG, Tamboli P, Babaian RJ. Can the number of cores

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

with high grade prostatic intraepithelial neoplasia predict cancer in men who undergo repeat biopsy? *Urology* 2004; **3**:503-8.

285. Nilsson S, Nordgren H, Eklov S, Logdahl M. Expression of Ki-67--a proliferation-associated antigen--in prostatic cancer. *Acta Oncologica*. 1991;**30**:177-9.
286. Noordzij MA, van der Kwast TH, van Steenbrugge GJ, van Weerden WM, Oomen MH, Schroder FH. Determination of Ki-67 defined growth fraction by monoclonal antibody MIB-1 in formalin-fixed, paraffin-embedded prostatic cancer tissues. *Prostate* 1995;**27**:154-9.
287. Novis DA, Zarbo RJ, Valenstein PA. Diagnostic uncertainty expressed in prostate needle biopsies. A College of American Pathologists Q-probes Study of 15,753 prostate needle biopsies in 332 institutions. *Archives of Pathology & Laboratory Medicine* 1999;**123**:687-92.
288. Nuorva K, Soini Y, Kamel D, Autio-Harminen H, Risteli L, Risteli J *et al*. Concurrent p53 expression in bronchial dysplasias and squamous cell lung carcinomas. *American Journal of Pathology* 1993;**142**:725-32.
289. O'Reilly MS, Holmgren L, Shing Y, Chen C, Rosenthal RA, Moses M *et al*. Angiostatin: a novel angiogenesis inhibitor that mediates the suppression of metastases by a Lewis lung carcinoma. *Cell* 1994;**79**:315-28.
290. Oesterling JE, CW, Jacobsen. Influence of patient age on the serum PSA concentration . An important clinical observation. *Urol Clin N Am* 1993;**20**:607-20.
291. Oliner JD, Kinzler KW, Meltzer PS, George DL, Vogelstein, B. Amplification of a gene encoding a p53-associated protein in human sarcomas. *Nature* 1992;**358**:80-3.
292. Oltvai ZN, Milliman CL, Korsmeyer SJ. Bcl-2 heterodimerizes in vivo with a conserved homolog, Bax, that accelerates programmed cell death. *Cell* 1993;**74**:609-19.

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

293. Olumi AF, Richie JP, Schultz DJ, D'Amico AV. Calculated volume of prostate cancer identifies patients with clinical stage T1C disease at high risk of biochemical recurrence after radical prostatectomy: a preliminary study. *Urology*;2000;**56**:273-7.
294. Orozco R, O'Dowd G, Kunnel B, Miller MC, Veltri RW. Observations on pathology trends in 62,537 prostate biopsies obtained from urology private practices in the United States. *Urology* 1998;**51**:186-95.
295. Osman I, Drobnjak M, Fazzari M, Ferrara J, Scher HI, Cordon-Cardo C. Inactivation of the p53 pathway in prostate cancer: impact on tumor progression. *Clinical Cancer Research* 1999;**5**:2082-8.
296. Pacelli A, Bostwick DG. Clinical significance of high-grade prostatic intraepithelial neoplasia in transurethral resection specimens. *Urology* 1997;**50**:355-9.
297. Parker C. Active surveillance: An individualised approach to early prostate cancer. *BJU Int.* 2003;**92**:2-3.
298. Parkes C, Wald NJ, Murphy P, George L, Watt HC, Kirby R *et al.* Prospective observational study to assess value of prostate specific antigen as screening test for prostate cancer. *British Medical Journal* 1995;**311**:1340-3.
299. Parkinson MC. Pre-neoplastic lesions of the prostate. *Histopathology* 1995;**27**:301-11.
300. Partin AW, Carter HB, Chan DW, Epstein JI, Oesterling JE, Rock RC *et al.* Prostate specific antigen in the staging of localized prostate cancer: influence of tumor differentiation, tumor volume and benign hyperplasia. *Journal of Urology* 1990;**143**:747-52.
301. Partin AW, Epstein JI, Cho KR, Gittelsohn AM, Walsh PC. Morphometric measurement of tumor volume and per cent of gland involvement as predictors of pathological stage in clinical stage B prostate cancer. *Journal of Urology* 2001;**141**:341-5.

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

302. Partin AW, Mangold L, Lamm D, Walsh PC, Epstein JI., Pearson J. Contemporary update of prostate cancer staging nomograms (Partin Tables) for the new millennium. *Urology* 2001;**58**:843-8.
303. Paulson DF. The impact of current staging procedures in assessing disease extent of prostatic adenocarcinoma. *Journal of Urology* 1979;**121**:300-2.
304. Paulson DF, Moul JW, Walther PJ. Radical prostatectomy for clinical stage T1-2N0M0 prostatic adenocarcinoma: long-term results. *Journal of Urology* 1990;**144**:1180-4.
305. Perachino M, di Ciolo L, Barbetti V, Ardoino S, Vitali A, Introini C *et al.* Results of rebiopsy for suspected prostate cancer in symptomatic men with elevated PSA levels. *European Urology* 1997;**32**:155-9.
306. Phillips ME, Kressel HY, Spritzer CE, Arger PH, Wein AJ, Axel L *et al.* Prostatic disorders: MR imaging at 1.5 T. *Radiology* 1987;**164**:386-92.
307. Pignatelli M, Stamp GW, Kafiri G, Lane D, Bodmer WF. Overexpression of p53nuclear oncoprotein in colorectal adenomas. *International Journal of Cancer* 1992;**50**:683-8.
308. Pirtskhalaishvili G, Shurin GV, Esche C, Cai Q, Salup RR, Bykovskaia SN *et al.* Cytokine-mediated protection of human dendritic cells from prostate cancer-induced apoptosis is regulated by the Bcl-2 family of proteins. *British Journal of Cancer* 2000;**83**:506-13.
309. Platt JF, Bree RL, Schwab RE. Accuracy of CT in the staging of carcinoma of the prostate. *Am J Roentgenol* 1987;**149**:315-8.
310. Prendergast NJ, Atkins MR, Schatte EC, Paulson DF, Walther PJ. p53 immunohistochemical and genetic alterations are associated at high incidence with post-irradiated locally persistent prostate carcinoma. *Journal of Urology* 1996;**155**:1685-92.

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

311. Qian J, Wollan P, Bostwick DG. The extent and multicentricity of high-grade prostatic intraepithelial neoplasia in clinically localized prostatic adenocarcinoma. *Human Pathology* 1997;**28**:143-8.
312. Quinn DI, Henshall SM, Head DR, Golovsky D, Wilson JD, Brenner PC *et al*. Prognostic significance of p53 nuclear accumulation in localized prostate cancer treated with radical prostatectomy. *Cancer Research* 2000;**60**:1585-94.
313. Raffo AJ, Perlman H, Chen MW, Day ML, Streitman JS, Buttyan R. Overexpression of bcl-2 protects prostate cancer cells from apoptosis in vitro and confers resistance to androgen depletion in vivo. *Cancer Research* 1995;**55**:4438-45.
314. Rakozy C, Grignon DJ, Li Y, Gheiler E, Gururajanna B, Pontes JE *et al*. p53 gene alterations in prostate cancer after radiation failure and their association with clinical outcome: a molecular and immunohistochemical analysis. *Pathology, Research & Practice* 1999;**195**:129-35.
315. Rakozy C, Grignon DJ, Sarkar FH, Sakr WA, Littrup P, Forman J. Expression of bcl-2, p53, and p21 in benign and malignant prostatic tissue before and after radiation therapy. *Modern Pathology* 1998;**11**:892-9.
316. Ramos CG, Carvahal GF, Mager DE, Haberer B, Catalona WJ. The effect of high grade prostatic intraepithelial neoplasia on serum total and percentage of free prostate specific antigen levels. *Journal of Urology* 1999;**162**:1587-90.
317. Rampino N, Yamamoto H, Ionov Y, Li Y, Sawai H, Reed JC *et al*. Somatic frameshift mutations in the BAX gene in colon cancers of the microsatellite mutator phenotype. *Science* 1997;**275**:967-9.
318. Renshaw AA, Santis WF, Richie JP. Clinicopathological characteristics of prostatic adenocarcinoma in men with atypical prostate needle biopsies. *Journal of Urology* 1998;**159**:2018-21.

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

319. Revelos K, Petraki C, Gregorakis A, Scorilas A, Papanastasiou P, Koutsilieris M. Immunohistochemical expression of Bcl-2 is an independant predictor of time-to-biochemical failure in patients with clinically localized prostate cancer following radical prostatectomy. *Anticancer Research* 2005;**25**:3123-33.
320. Reyes AO, Humphrey PA. Diagnostic effect of complete histologic sampling of prostate needle biopsy specimens. *American Journal of Clinical Pathology* 1998;**109**:416-22.
321. Richie JP, Kavoussi LR, Ho GT, Vickers MA, O'Donnell MA, St Laurent D *et al.* Prostate cancer screening: role of the digital rectal examination and prostate-specific antigen. *Annals of Surgical Oncology* 1994;**1**:117-20.
322. Rifkin M, Dahnert WF, Kurtz AB. State of the art: endorectal sonography of the prostate gland. *Am J Roentgenol* 1990;**154**:691-700.
323. Rockall A, Sohaib S, Harisinghani M, Babar S, Singh N, Jeyarajah A *et al.* Diagnostic performance of nanoparticle-enhanced MRI in the diagnosis of lymph node metastases in patients with endometrial and cervical cancer. *Journal of Clinical Oncology* 2005;**23**:2813-21.
324. Roehl K, Antenor J, Catalona WJ. Serial biopsy results in prostate cancer screening study. *Journal of Urology* 2002;**167**:2435-9.
325. Roehrborn CG, Boyle P, Gould AL, Waldstreicher J. Serum prostate-specific antigen as a predictor of prostate volume in men with benign prostatic hyperplasia. *Urology* 1999;**53**:581-9.
326. Ronnett BM, Carmichael MJ, Carter HB, Epstein JI. Does high grade prostatic intraepithelial neoplasia result in elevated serum prostate specific antigen levels? *Journal of Urology* 1993;**150**:386-9.
327. Rosanelli GP, Steindorfer P, Wirsberger GH, Klimpfinger M, Ratschek M, Puerstner P *et al.* Mutant p53 expression and DNA analysis in human breasy cancer comparison with conventional clinicopathologic parameters. *Anticancer Research* 1995;**15**:581-6.

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

328. Rosser C, Reyes AO, Vakar L, Levy L, Kuban DA, Hoover D *et al.* Bcl-2 is significantly overexpressed in localized radio-recurrent prostate carcinoma, compared to localized radio-naïve prostate carcinoma. *Inte-J-Rad-Oncol-Biol-Phys* 2003; **56**:1-6.
329. Rowinsky EK, Onetto N, Canetta RM, Arbuck SG. Taxol: the first of the taxanes, an important new class of antitumor agents. [Review] [56 refs]. *Seminars in Oncology* 1992;**19**:646-62.
330. Royuela M, De Miguel MP, Bethencourt FR, Fraile B, Arenas MI, Paniagua R. IL-2, its receptors, and bcl-2 and bax genes in normal, hyperplastic and carcinomatous human prostates: immunohistochemical comparative analysis. *Growth Factors* 2000;**18**:135-46.
331. Sakr, W. A. The frequency of carcinoma and intra-epithelial neoplasia of the prostate in the young male patients. *Journal of Urology* 150, 379-385. 1993.
332. Sakr WA, Angelakis K, deGuia K, Cher ML, Wood DP, Jr., Grignon DJ *et al.* The prevalence and evolution of isolated high grade prostatic intraepithelial neoplasia (HGPIN) in african american (AA) and caucasian men- A study of needle biopsies. *Journal of Urology* 2002;**163**:56, Abs 245.
333. Sandhu SS, Kaisary AV. Localised carcinoma of the prostate: a paradigm of uncertainty. *Postgrad.Med.J.* 1997;**73**:691-6.
334. Sasor A, Wagrowska-Danilewicz M, Danilewicz M. Ki-67 antigen and P53 protein expression in benign and malignant prostatic lesions. Immunohistochemical quantitative study. *Polish Journal of Pathology* 2000;**51**:31-6.
335. Sato T, Hanada M, Bodrug S, Irie S, Iwama N, Boise LH *et al.* Interactions among members of the Bcl-2 protein family analyzed with a yeast two-hybrid system. [erratum appears in Proc Natl Acad Sci U S A 1995 Feb 28;92(5):2016]. *Proceedings of the National Academy of Sciences of the United States of America* 1994;**91**:9238-42.

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

336. Sawan A, Randall B, Angus B, Wright C, Henry JA, Ostrowski J *et al*. Retinoblastoma gene and p53 gene expression related to relapse and survival in human breast cancer: an immunohistochemical study. *Journal of Pathology* 1992;**168**:23-8.
337. Sawhney N, Hall PA. Ki67--structure, function, and new antibodies. *Journal of Pathology* 1992;**168**:161-2.
338. Scalzo DA, Kallakury BV, Gaddipati RV, Sheehan CE, Keys HM, Savage D *et al*. Cell proliferation rate by MIB-1 immunohistochemistry predicts postradiation recurrence in prostatic adenocarcinomas. *American Journal of Clinical Pathology* 1998;**109**:163-8.
339. Scardino PT, Shinohara K, Wheeler TM, Carter SS. Staging of prostate cancer. Value of ultrasonography. *Urol Clin N Am* 1989;**16**:713-34.
340. Scardino PT, Weaver R, Hudson MA. Early detection of prostate cancer. *Human Pathology* 1992;**23**:211-22.
341. Scherr DS, Vaughan ED, Jr., Wei J, Chung M, Felsen D, Allbright R *et al*. BCL-2 and p53 expression in clinically localized prostate cancer predicts response to external beam radiotherapy. *Journal of Urology* 1999;**162**:12-6.
342. Schlichtholz B, Legros Y, Gillet D, Gaillard, Marty M, Lane D *et al*. The immune response to p53 in breast cancer patients is directed against immunodominant epitopes unrelated to the mutational hot spot. *Cancer Research* 1992;**52**:6380-4.
343. Schmid H-P, McNeal JE. An abbreviated standard procedure for accurate tumor volume estimation in prostate cancer. *American Journal of Surgical Pathology* 1992;**16**:184-91.
344. Schned AR, Wheeler KJ, Hodorowski CA, Heaney JA, Ernstoff MS, Amdur *et al*. Tissue-shrinkage correction factor in the calculation of prostate cancer volume. *American Journal of Surgical Pathology* 1996;**20**:1501-6.

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

- 345. Scholzen T, Gerdes J. The Ki-67 protein: from the known and the unknown. *Journal of Cellular Physiology*. 2000;**182**:311-22.
- 346. Scott RJ, Hall PA, Haldane JS, van Noorden S, Price Y, Lane DP *et al*. A comparison of immunohistochemical markers of cell proliferation with experimentally determined growth fraction. *Journal of Pathology* 1991;**165**:173-8.
- 347. Seaman EK, Whang IS, Cooner W, Olsson CA, Benson MC. Predictive value of prostate-specific antigen density for the presence of micrometastatic carcinoma of the prostate. *Urology* 2002;**43**:645-8.
- 348. Segal NH, Cohen RJ, Haffejee Z, Savage N. BCL-2 proto-oncogene expression in prostate cancer and its relationship to the prostatic neuroendocrine cell. *Archives of Pathology & Laboratory Medicine* 1994;**118**:616-8.
- 349. Serth J, Kuczyk MA, Bokemeyer C, Hervatin C, Nafe R, Tan HK *et al*. p53 immunohistochemistry as an independent prognostic factor for superficial transitional cell carcinoma of the bladder. *British Journal of Cancer* 1995;**71**:201-5.
- 350. Shepherd D, Keetch DW, Humphrey PA, Smith DS, Stahl D. Repeat biopsy strategy in men with isolated prostatic intraepithelial neoplasia on prostate needle biopsy. *Journal of Urology* 1996;**156**:460-2.
- 351. Shi SR, Chaiwun B, Young L, Imam A, Cote RJ, Taylor CR. Antigen retrieval using pH 3.5 glycine-HCl buffer or urea solution for immunohistochemical localization of Ki-67. *Biotechnic & Histochemistry* 1994;**69**:213-5.
- 352. Shi SR, Cote RJ, Taylor CR. Antigen retrieval immunohistochemistry: past, present, and future. *Journal of Histochemistry & Cytochemistry* 1997;**45**:327-43.
- 353. Shi SR, Cote RJ, Taylor CR. Antigen retrieval immunohistochemistry and molecular morphology in the year 2001. *Applied Immunohistochemistry & Molecular Morphology* 2001;**9**:107-16.

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

354. Shi SR, Cote RJ, Taylor CR. Antigen retrieval techniques: current perspectives. *Journal of Histochemistry & Cytochemistry* 2001;**49**:931-7.
355. Shi SR, Imam SA, Young L, Cote RJ, Taylor CR. Antigen retrieval immunohistochemistry under the influence of pH using monoclonal antibodies. *Journal of Histochemistry & Cytochemistry* 1995;**43**:193-201.
356. Shi X, Gandour-Edwards R, Beckett L, Deitch AD, de Vere WR. A modified yeast assay used on archival samples of localised prostate cancer tissue improves the detection of p53 abnormalities and increases their predictive value. *BJU Int.* 2004;996-1002.
357. Shiao YH, Rugge M, Correa P, Lehman HP, Scheer WD. p53 alteration in precancerous gastric lesions. *American Journal of Pathology* 1994;**144**:511-7.
358. Sidransky D, Mikkelsen T, Schwechheimer K, Rosenbaum ML, Cavanee W, Vogelstein B. Clonal expansion of p53 mutant cells is associated with brain tumour progression. *Nature* 1992;**355**:846-7.
359. Silvestrini R, Benini E, Daidone MG, Veneroni S, Boracchi P, Cappelletti V *et al.* p53 as an independant prognostic marker in lymph node-negative breast cancer patients. *Journal of the National Cancer Institute* 1993;**85**:965-70.
360. Skjorten FJ, Berner A, Harvei S, Røksahm TE, Tretli S. Prostatic intraepithelial neoplasia in surgical resections: relationship to coexistent adenocarcinoma and atypical adenomatous hyperplasia of the prostate. *Cancer* 1997;**79**:1172-9.
361. Soussi T, Legros Y, Lubin R, Ory K, Schlichtholz B. Multifactorial analysis of p53 alteration in human cancer: a review. *International Journal of Cancer* 1994; **57**:1-9.
362. Sozzi G, Miozzo M, Donghi R, Pilotti S, Cariani CT, Pastorini U *et al.* Deletions of 17p and p53 mutations in preneoplastic lesions of the

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

lung. *Cancer Research* 1992;**52**:6079-82.

363. Spring WB, Alden MW. Evaluation of needle biopsy in the diagnosis of prostatic carcinoma. *Can Med Assoc J* 1954;**70**:179-85.
364. Stackhouse GB, Sesterhenn IA, Bauer JJ, Mostofi FK, Connelly RR, Srivastava SK *et al.* p53 and bcl-2 immunohistochemistry in pretreatment prostate needle biopsies to predict recurrence of prostate cancer after radical prostatectomy. *Journal of Urology* 1999;**162**:2040-5.
365. Stamey TA, Kabalin JN, McNeal JE, Johnstone IM, Freiha FS, Redwine EA *et al.* Prostate specific antigen in the diagnosis and treatment of adenocarcinoma of the prostate. II. Radical prostatectomy treated patients. *Journal of Urology* 1989;**141**:1076-83.
366. Stamey, TA, Yang, N., Hay, A. R., McNeal, J. E., Freiha FS, and Redwine EA. Prostate-specific antigen as a serum marker for adenocarcinoma of the prostate. *New England Journal of Medicine* 317(15), 909-916. 1987.
367. Stamey, TA, Yemoto, C. M., McNeal, J. E., Sigbal, B. M., and Johnstone IM. Prostate cancer is highly predictable: a prognostic equation based on all morphological variables in radical prostatectomy specimens. *Journal of Urology* 163(4), 1155-1160. 2000.
368. Stapleton AM, Zbell P, Kattan MW, Yang G, Wheeler TM, Scardino PT *et al.* Assessment of the biologic markers p53, Ki-67, and apoptotic index as predictive indicators of prostate carcinoma recurrence after surgery. *Cancer* 1998;**82**:168-75.
369. Stattin P, Bergh A, Karlberg L, Nordgren H, Damber JE. p53 immunoreactivity as prognostic marker for cancer-specific survival in prostate cancer. *European Urology* 1996;**30**:65-72.
370. Stattin P, Damber JE, Karlberg L, Bergh A. Cell proliferation assessed by Ki-67 immunoreactivity on formalin fixed tissues is a predictive

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

factor for survival in prostate cancer. *Journal of Urology* 1997;**157**:219-22.

371. Stattin P, Damber JE, Modig H, Bergh A. Pretreatment p53 immunoreactivity does not infer radioresistance in prostate cancer patients. *International Journal of Radiation Oncology, Biology, Physics* 1996;**35**:885-9.
372. Stattin P, Westin P, Damber JE, Bergh A. Short-term cellular effects induced by castration therapy in relation to clinical outcome in prostate cancer. *British Journal of Cancer* 1998;**77**:670-5.
373. Stenmark-Askmal M, Stal O, Sullivan S, Ferraud L, Sun XF, Carstensen J *et al.* Cellular accumulation of p53 protein: an independant prognostic factor in stage 2 breast cancer. *European Journal of Cancer* 1994;**30**:175-80.
374. Stricker HJ, Jay JK, Linden MD, Tamboli P, Amin MB. Determining prognosis of clinically localized prostate cancer by immunohistochemical detection of mutant p53. *Urology* 1996;**47**:366-9.
375. Strohmeyer D, Rossing C, Bauerfeind A, Kaufmann O, Schlechte H, Bartsch *et al.* Vascular endothelial growth factor and its correlation with angiogenesis and p53 expression in prostate cancer. *Prostate* 2000;**45**:216-24.
376. Taftachi R, Ayhan A, Ekici S, Ergen A, Ozen H. Proliferating-cell nuclear antigen (PCNA) as an independent prognostic marker in patients after prostatectomy: a comparision of PCNA and Ki-67. *BJU Int.* 2005;**95**:650-4.
377. Takayama H, Shin M, Nonomura N, Okuyama A, Aozasa K. p53 mutations in prostatic intraepithelial neoplasia and concurrent carcinoma: analysis of laser capture microdissected specimens from non-transition and transition zones. *Japanese Journal of Cancer Research* 2000;**91**:941-7.

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

378. Talcott JA, Clark JA, Stark PC, Mitchell SP. Long-term treatment related complications of brachytherapy for early prostate cancer: a survey of patients previously treated. *Journal of Urology* 2001;**166**:494-9.
379. Tamboli P, Amin MB, Schultz DS, Linden MD, Kubus J. Comparative analysis of the nuclear proliferative index (Ki-67) in benign prostate, prostatic intraepithelial neoplasia, and prostatic carcinoma. *Modern Pathology* 1996;**9**:1015-9.
380. Tanaka M, Omura K, Watanabe Y, Oda Y, Nakanishi I. Prognostic factors of colorectal cancer:K-ras mutation, overexpression of the p53 protein, and cell proliferative activity. *Journal of Surgical Oncology* 1994;**57**:57-64.
381. Tanji N, Kikugawa T, Yokoyama M. Immunohistochemical study of cyclooxygenases in prostatic adenocarcinoma; relationship to apoptosis and Bcl-2 protein expression. *Anticancer Research* 2000;**20**:2313-9.
382. Tanji N, Yokoyama M, Sugamoto T, Takeuchi M, Terada N. Apoptosis in prostatic adenocarcinomas; a study of relationship to Ki-67 and Bcl-2 protein expression. *Anticancer Research* 1998;**18**:1111-6.
383. Taylor CR, Shi SR, Chaiwun B, Young L, Imam SA, Cote RJ. Strategies for improving the immunohistochemical staining of various intranuclear prognostic markers in formalin-paraffin sections: androgen receptor, estrogen receptor, progesterone receptor, p53 protein, proliferating cell nuclear antigen, and Ki-67 antigen revealed by antigen retrieval techniques. *Human Pathology* 1994;**25**:263-70.
384. Taylor CR, Shi SR, Chen C, Young L, Yang C, Cote RJ. Comparative study of antigen retrieval heating methods: microwave, microwave and pressure cooker, autoclave, and steamer. *Biotechnic & Histochemistry* 1996;**71**:263-70.
385. Theodorescu D, Broder SR, Boyd JC, Mills SE, Frierson HF, Jr. p53, bcl-2 and retinoblastoma proteins as long-term prognostic markers in localized carcinoma of the prostate. *Journal of Urology* 1997;**158**:131-7.

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

386. Thomas A, Giesler T, White E. p53 mediates bcl-2 phosphorylation and apoptosis via activation of the Cdc42/JNK1 pathway. *Oncogene* 2000;**19**:5259-69.
387. Thomas DJ, Robinson M, King P, Hasan T, Charlton R, Martin J *et al*. p53 expression and clinical outcome in prostate cancer. *British Journal of Urology* 1993;**72**:778-81.
388. Thomenius M, Wang N, Reineks E, Wang Z, Distelhorst C. Bcl-2 on the endoplasmic reticulum regulates bax activity by binding to BH3-only proteins. *J Biol Chem* 2003;**278**:6243-50.
389. Thompson SJ, Mellon K, Charlton RG, Marsh C, Robinson M, Neal DE. P53 and Ki-67 immunoreactivity in human prostate cancer and benign hyperplasia. *British Journal of Urology* 1992;**69**:609-13.
390. Thor AD, Moore DH, Edgerton SM, Kawasaki ES, Reihaus E, Lynch HT *et al*. Accumulation of p53 tumor suppressor gene protein: an independant marker of prognosis in breast cancers. *Journal of the National Cancer Institute* 1992;**84**:845-55.
391. Thor AD, Yandell DW. Prognostic significance of p53 over expression in node-negative breast carcinoma: preliminary studies support cautious optimism. *Journal of the National Cancer Institute* 1993;**85**:176-7.
392. Tinari N, Natoli C, Angelucci D, Tenaglia R, Fiorentino B, Di Stefano P *et al*. DNA and S-phase fraction analysis by flow cytometry in prostate cancer. Clinicopathologic implications. *Cancer* 1993;**71**:1289-96.
393. Tsuji M, Kanda K, Murakami Y, Kurokawa Y, Kanayama H, Sano T *et al*. Biologic markers in prostatic intraepithelial neoplasia: immunohistochemical and cytogenetic analyses. *Journal of Medical Investigation* 1999;**46**:35-41.
394. Tsuji M, Murakami Y, Kanayama H, Sano T, Kagawa S. Immunohistochemical analysis of Ki-67 antigen and Bcl-2 protein

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

- expression in prostate cancer: effect of neoadjuvant hormonal therapy. *British Journal of Urology* 1998;**81**:116-21.
395. Tsuzuki T, Tsunoda S, Sakaki T, Konishi N, Hiasa Y, Nakamura M. Alterations of retinoblastoma, p53, p16(CDKN2), and p15 genes in human astrocytomas. *Cancer* 1996;**78**:287-93.
 396. Uchida T, Wada C, Shitara T, Egawa S, Koshiha K. Infrequent involvement of p53 gene mutations in the tumourigenesis of Japanese prostate cancer. *British Journal of Cancer* 1993;**68**:751-5.
 397. Uzoaru I, Rubenstein M, Mirochnik Y, Slobodskoy L, Shaw M, Guinan P. An evaluation of the markers p53 and Ki-67 for their predictive value in prostate cancer. *Journal of Surgical Oncology* 1998;**67**:33-7.
 398. van der Kwast TH, Tetu B. Androgen receptors in untreated and treated prostatic intraepithelial neoplasia. *European Urology* 1996;**30**:265-8.
 399. Van Veldhuizen PJ, Sadasivan R, Garcia F, Austenfeld MS, Stephens RL. Mutant p53 expression in prostate carcinoma. *Prostate* 1993;**22**:23-30.
 400. Varley J. Germline TP53 mutations and Li-Fraumeni Syndrome. *Hum-Mutat* 2003;**21**:313-20.
 401. Vaux DL, Cory S, Adams JM. Bcl-2 gene promotes haemopoietic cell survival and cooperates with c-myc to immortalize pre-B cells. *Nature* 1988;**335**:440-2.
 402. Vesalainen S, Lipponen P, Talja M, Syrjanen K. Mitotic activity and prognosis in prostatic adenocarcinoma. *Prostate* 1995;**26**:80-6.
 403. Vis AN, Noordzij MA, Fitoz K, Wildhagen MF, Schroder FH, van der Kwast TH. Prognostic value of cell cycle proteins p27(kip1) and MIB-1, and the cell adhesion protein CD44s in surgically treated patients with prostate cancer. *Journal of Urology* 2000;**164**:2156-61.

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

404. Visakorpi T. Proliferative activity determined by DNA flow cytometry and proliferating cell nuclear antigen (PCNA) immunohistochemistry as a prognostic factor in prostatic carcinoma. *Journal of Pathology* 1992;**168**:7-13.
405. Visakorpi T, Kallioniemi OP, Heikkinen A, Koivula T, Isola J. Small subgroup of aggressive, highly proliferative prostatic carcinomas defined by p53 accumulation. *Journal of the National Cancer Institute* 1992;**84**:883-7.
406. Voeller HJ, Sugars LY, Pretlow T, Gelmann EP. p53 oncogene mutations in human prostate cancer specimens. *Journal of Urology* 1994;**151**:492-5.
407. Vojtesek B, Bartek J, Midgely C, Lane DP. An immunochemical analysis of the human nuclear phosphoprotein p53. New monoclonal antibodies and epitope mapping using recombinant p53. *Journal of Immunological Methods* 1992;**151**:237-44.
408. Wada H, Asada M, Nakazawa S, Itoh H, Kobayashi Y., Inoue T. *et al.* Clonal expansion of p53 mutant cells in leukaemia progression in vitro. *Leukaemia* 1994;**8**:53-9.
409. Walsh PC. Anatomic radical retropubic prostatectomy. *Campbells Urology* 1998;**7th edn**.
410. Wang LD, Hong JY, Qiu SL, Gao H, Yang CS. Accumulation of p53 protein in human esophageal precancerous lesions: a possible early biomarker for carcinogenesis. *Cancer Research* 1993;**53**:1783-7.
411. Weinstein MH. Digital image analysis of proliferative index: two distinct populations of high-grade prostatic intraepithelial neoplasia in close proximity to adenocarcinoma of the prostate. *Human Pathology* 1998;**29**:620-6.
412. Weir EG, Partin AW, Epstein JI. Correlation of serum prostate specific antigen and quantitative immunohistochemistry. *Journal of Urology* 2000;**163**:1739-42.

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

413. Wertz IE, Deitch AD, Gumerlock PH, Gandour-Edwards R, Chi SG, de V *et al.* Correlation of genetic and immunodetection of TP53 mutations in malignant and benign prostate tissues. *Human Pathology* 1996;**27**:573-80.
414. Westin P, Stattin P, Damber JE, Bergh A. Castration therapy rapidly induces apoptosis in a minority and decreases cell proliferation in a majority of human prostatic tumors. *American Journal of Pathology* 1995;**146**:1368-75.
415. Whitmore WF, Warner JA, Thompson IM Jr. Expectant management of localized prostatic cancer. *Cancer* 1991;**67**:1091-6.
416. Wijnhoven S, van Steeg H. Transgenic and Knock out mice for DNA repair functions in carcinogenesis and mutagenesis. *Toxicology* 2003;**193**:171-87.
417. Wills ML, Hamper UM, Partin AW, Epstein JI. Incidence of high-grade prostatic intraepithelial neoplasia in sextant needle biopsy specimens. *Urology* 1997;**49**:367-73.
418. Wilt TJ, Brawer MK. The Prostate Cancer Intervention Versus Observation Trial: a randomized trial comparing radical prostatectomy versus expectant management for the treatment of clinically localized prostate cancer. *Journal of Urology* 1994;**152**:1910-4.
419. Winter SF, Minna JD, Johnson BE, Takahashi T, Gazdar AF, Carbone DP. Development of antibodies in lung cancer patients appears to be dependant upon the type of p53 mutation. *Cancer Research* 1992;**52**:4168-74.
420. Wiseman S, Stoler D, Anderson G. The role of genomic instability in the pathogenesis of squamous cell carcinoma of the head and neck. *Surg Oncol Clin N Am* 2004;**13**:1-11.
421. Wu TT, Hsu Y, Lee J, Huang JK. The role of p53, bcl-2, and e-cadherin expression in predicting biochemical relapse for organ confined prostate cancer in Taiwan. *Journal of Urology* 2003;**170**:78-

Prostatic Intraepithelial Neoplasia and Prostate Cancer.

- 81.
422. Xie W, Wong YC, Tsao SW. Correlation of increased apoptosis and proliferation with development of prostatic intraepithelial neoplasia (PIN) in ventral prostate of the Noble rat. *Prostate* 2000;**44**:31-9.
423. Yaman O, Ozdiler E, Orhan D, Sak SD, Baltaci S, Tulunay O *et al*. Immunohistochemical determination of p53 protein in prostatic cancer and prostatic intraepithelial neoplasms. *Urologia Internationalis* 1997;**58**:199-202.
424. Yang E, Korsmeyer SJ. Molecular thanatopsis. *Blood* 1996;**88**:386-401.
425. Yang G, Stapleton AM, Wheeler TM, Truong LD, Timme TL, Scardino PT *et al*. Clustered p53 immunostaining: a novel pattern associated with prostate cancer progression. *Clinical Cancer Research* 1996;**2**:399-401.
426. Yasunaga Y, Shin M, Fujita MQ, Nonomura N, Miki T, Okuyama A *et al*. Different patterns of p53 mutations in prostatic intraepithelial neoplasia and concurrent carcinoma: analysis of microdissected specimens. *Laboratory Investigation* 1998;**78**:1275-9.
427. Zeng ZS, Sarkis AS, Zhang ZF, Klimstra DS, Charytonowicz E, Guillem JG *et al*. p53 nuclear overexpression: an independant predictor of survival in lymph node positive colorectal cancer patients. *Journal of Clinical Oncology* 1994;**12**:2043-50.